

149

DIRECTIONS FOR USE

APPLES & PEARS: To control European Red Mite, Clover Mite, and Two Spotted Mite, mix 2 pts. in 100 gals. water or 2 tsp. in 1 gal. water. To control Woolly Apple Aphid and Forbes Scale, mix 1 pt. in 100 gals. water or 1 tsp. in 1 gal. water. To control Green Apple Aphid and Rosy Apple Aphid, mix 1½ pts. in 100 gals. water or 1 tsp. in 1½ gal. water. Make two or more applications as needed for mites, one or more for aphids. At pre-pink or pink stage of bloom or when aphids appear. Do not use within 3 days of harvest. This product may injure foliage on some varieties of apples and pears.

TOMATOES: To control spiders and mites, mix 1½ pts. in 100 gals. water or 3 tsp. in 1 gal. water. To control aphids, mix 1 pt. in 100 gals. water or 3 tsp. in 1 gal. water. Make one or more applications as needed. Do not use within 3 days of harvest.

BEANS: To control Mexican Bean Beetle, Spider Mites, Leafhopper, and aphids, mix 2 pts. in 100 gals. water or 3 tsp. in 1 gal. water. Make one or more applications as needed. Do not use within 3 days of harvest.

PEAS & POTATOES: To control Pea Aphid on peas or aphids on potatoes, mix 2 pts. in 100 gals. water or 3 tsp. in 1 gal. water. Make one or more applications as needed. Do not use within 3 days of harvest. Do not use pea vines for forage.

CABBAGE, BROCCOLI, KALE, MUSTARD, & TURNIP: To control aphids, Imported Cabbage Worm, mix 2 pts. in 100 gals. water or 3 tsp. in 1 gal. water. Make full coverage applications. Repeat as necessary. Do not use within 3 days of harvest on broccoli and turnips, 7 days on others.

LIQUID FLY BAIT: To control House and Stable Flies in Barns, Feed Lots, Poultry Houses, Utility buildings and around food disposals, mix 5 ounces with 5 ounces of corn syrup and add to 2 gals. water. Paint or spray on desired surfaces.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and conditions of application equipment will vary, Seller makes no claims other than those stated on this label.

893-175

ACME®

60-50

malathion 50% spray

for flies and garden insects

ACTIVE INGREDIENTS:

Malathion* 50.00%

Xylene 38.51%

INERT INGREDIENTS 11.49%

TOTAL 100.00%

*O,O-Dimethyl dithiophosphate of diethyl mercaptosuccinate.

WARNING: Keep Out of Reach of Children
See Side Panels for Additional Warning Statements

NET CONTENTS ONE QUART

FLOWERS: Scale and in 1 gal. Mealy bug Plant Bug 100 gals. and Juniper water. To or 2½ tsp. in 50 when infested when crawling including of Crassul.

WARNING: Avoid contamination of

This product Do not apply in weather conditions. Apply this

This product residues. operative

DO NOT FLAME. D

EPA REG.

EPA EST.

ACME⁽²⁾**60-50****malathion
50% spray****for flies and garden insects****ACTIVE INGREDIENTS:**

Malathion* 50.00%

Xylene 38.51%

INERT INGREDIENTS 11.49%

TOTAL 100.00%

*O,O-Dimethyl dithiophosphate of diethyl mercaptosuccinate.

WARNING: Keep Out of Reach of Children
See Side Panels for Additional Warning Statements**NET CONTENTS ONE QUART****DIRECTIONS FOR USE**

FLOWERS, ORNAMENTALS, & SHRUBS: To control Oyster Shell Scale and Eucymus Scale, mix 1 pt. in 100 gals. water or 1 tsp. in 1 gal. water. To control aphids, Spider Mites, White Fly, Thrips, Mealy bugs, Japanese Beetle Adult, Four Lined Plant Bug, Tarnished Plant Bug, Rose Leafhopper, and Potato Leafhopper, mix 1½ pts. in 100 gals. water or 1½ tsp. in 1 gal. water. To control Bagworms and Juniper Scale, mix 2 pts. in 100 gals. water or 2 tsp. in 1 gal. water. To control Soft Brown Scale, mix 2½ pts. in 100 gals. water or 2½ tsp. in 1 gal. water. To control Pine Needle Scale, mix 2 pts. in 50 gals. water or 4 tsp. in 1 gal. water. Apply in summer when infestation occurs. Repeat in 10 to 12 days. For scales, apply when crawlers have settled on foliage. Do not use on certain ferns, including Boston, Maidenhair and Pteris, as well as some species of Crassula, because of possible injury to these ornamentals.

WARNING: Harmful if swallowed. Avoid breathing of spray mist. Avoid contact with skin. Wash thoroughly after use. Avoid contamination of feed or foodstuffs. Keep away from heat or open flames.

This product is toxic to fish. Keep out of lakes, streams, and ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME. DO NOT RE-USE CONTAINER. DESTROY WHEN EMPTY.

EPA REG. NO. 33955-394-AA


EPA EST. NO. 2217-KS-1

DISTRIBUTED BY

ACME DIVISION

-GORDON CORPORATION

KANSAS CITY, KANSAS 64118



2/17/71

DIRECTIONS FOR USE

APPLES & PEARS: To control European Red Mite, Clover Mite, and Two Spotted Mite, mix 2 tsp. in 1 gal. water. To control Woolly Apple Aphid and Forbes Scale, mix 1 tsp. in 1 gal. water. To control Green Apple Aphid and Rosy Apple Aphid, mix 1½ tsp. in 1 gal. water. Make two or more applications as needed for mites, one or more for aphids. At pre-pink or pink stage of bloom or when aphids appear. Do not use within 3 days of harvest. This product may injure foliage on some varieties of apples and pears.

TOMATOES: To control spiders and mites, mix 3 tsp. in 1 gal. water. To control aphids, mix 3 tsp. in 1 gal. water. Make one or more applications as needed. Do not use within 3 days of harvest.

BEANS: To control Mexican Bean Beetle, Spider Mites, Leafhopper, and aphids, mix 3 tsp. in 1 gal. water. Make one or more applications as needed. Do not use within 3 days of harvest.

PEAS & POTATOES: To control Pea Aphid on peas or aphids on potatoes, mix 3 tsp. in 1 gal. water. Make one or more applications as needed. Do not use within 3 days of harvest. Do not use pea vines for forage.

CABBAGE, BROCCOLI, KALE, MUSTARD, & TURNIP: To control aphids, Imported Cabbage Worm, mix 3 tsp. in 1 gal. water. Make full coverage applications. Repeat as necessary. Do not use within 3 days of harvest on broccoli and turnips, 7 days on others.

LIQUID FLY BAIT: To control House and Stable Flies in Barns, Feed Lots, Poultry Houses, Utility buildings and around food disposals, mix 5 ounces with 5 ounces of corn syrup and add to 2 gals. water. Paint or spray on desired surfaces.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and conditions of application equipment will vary, Seller makes no claims other than those stated on this label.

893-175



ACTIVE INGREDIENTS:

Malathion*	50.00%
Xylene	38.51%

INERT INGREDIENTS	11.49%
-------------------	--------

TOTAL	100.00%
-------	---------

*O,O-Dimethyl dithiophosphate of diethyl mercaptosuccinate.

WARNING:

Keep Out of Reach of Children

See Side Panels for Additional Warning Statements

NET CONTENTS ONE PINT

DIRECTIONS FOR USE

FLOWERS, ORNAMENTALS & SHRUBS: To control Euonymus Scale, mix 1 tsp. in 1 gal. water. Mites, White Fly, Thrips, Mealy bugs, Japanese Plant Bug, Tarnished Plant Bug, Rose Leafhopper, mix 1½ tsp. in 1 gal. water. To control Scale mix 2 tsp. in 1 gal. water. To control 2½ tsp. in 1 gal. water. To control Pine in 1 gal. water. Apply in summer when in 10 to 12 days. For scales, apply when foliage. Do not use on certain ferns, including Pteris, as well as some species of Craspedangium to these ornamentals.

WARNING: Harmful if swallowed. Avoid breathing vapors. Avoid contact with skin. Wash thoroughly after use. Do not use in or near water. Do not use in or near feed or foodstuffs. Keep away from children.

This product is toxic to fish. Keep out of lakes and streams. Do not apply where runoff is likely to occur. In adverse weather conditions favor drift from areas of application. Clean water by cleaning of equipment, or disposal of product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment. Protective information may be obtained from the Cooperative Agricultural Extension Service.

DO NOT USE, POUR, SPILL, OR STORE NEAR FOOD. DO NOT RE-USE CONTAINER. DESTROY WHEN EMPTY.

EPA REG. NO. 33955-394-AA

EPA EST. NO. 2217-KS-1

DISTRIBUTED BY

ACME DIVISION

GORDON CORP.

KANSAS CITY, KANSAS

SE

ed Mite, Clover Mite, and
ter. To control Woolly Apple
al. water. To control Green
1½ tsp. in 1 gal. water.
for mites, one or more for
m or when aphids appear.
product may injure foliage

nix 3 tsp. in 1 gal. water.
water. Make one or more
3 days of harvest.

Spider Mites, Leafhopper.
Make one or more applica-
ys of harvest.


d on peas or aphids on
one or more applications
harvest. Do not use pea

TURNIP: To control aphids,
al. water. Make full cover-
not use within 3 days of
others.

Stable Flies in Barns. Feed
around food disposals. mix
add to 2 gals. water. Paint

conforms to the ingredient
of use, such as weather,
ditions of application equip-
other than those stated on

175



ACTIVE INGREDIENTS:

Malathion*	50.00%
Xylene	38.51%

INERT INGREDIENTS	11.49%
-------------------	--------

TOTAL	100.00%
-------	---------

*O,O-Dimethyl dithiophosphate of diethyl mercapto-
succinate.

WARNING:

Keep Out of Reach of Children

See Side Panels for Additional Warning Statements

NET CONTENTS ONE PINT

DIRECTIONS FOR USE

FLOWERS, ORNAMENTALS & SHRUBS: To control Oyster Shell Scale and Euonymus Scale, mix 1 tsp. in 1 gal. water. To control aphids, Spider Mites, White Fly, Thrips, Mealy bugs, Japanese Beetle Adult, Four Lined Plant Bug, Tarnished Plant Bug, Rose Leafhopper, and Potato Leafhopper, mix 1½ tsp. in 1 gal. water. To control Bagworms and Juniper Scale mix 2 tsp. in 1 gal. water. To control Soft Brown Scale, mix 2½ tsp. in 1 gal. water. To control Pine Needle Scale, mix 4 tsp. in 1 gal. water. Apply in summer when infestation occurs. Repeat in 10 to 12 days. For scales, apply when crawlers have settled on foliage. Do not use on certain ferns, including Boston, Maidenhair and Pteris, as well as some species of Crassula, because of possible injury to these ornamentals.

WARNING: Harmful if swallowed. Avoid breathing of spray mist. Avoid contact with skin. Wash thoroughly after use. Avoid contamination of feed or foodstuffs. Keep away from heat or open flames.

This product is toxic to fish. Keep out of lakes, streams, and ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME. DO NOT RE-USE CONTAINER. DESTROY WHEN EMPTY.

EPA REG. NO. 33955-394-AA

EPA EST. NO. 2217-KS-1

DISTRIBUTED BY

ACME DIVISION

-GORDON CORPORATION

KANSAS CITY, KANSAS 64110

DIRECTIONS FOR USE

On all vegetables listed below, mix 3 teaspoonsful in one gallon water.

TOMATOES: To control spiders, mites, and aphids, spray as needed. Do not use within 3 days of harvest.

BEANS: To control Mexican Bean Beetle, Spider Mites, Leafhopper, and aphids, spray as needed. Do not use within 3 days of harvest.

PEAS & POTATOES: To control Pea Aphid on peas or aphids on potatoes, spray as needed. Do not use within 3 days of harvest.

CABBAGE, BROCCOLI, KALE, MUSTARD, & TURNIP: To control aphids and Imported Cabbage Worms, spray as needed. Do not use within 3 days of harvest on broccoli and turnips, 7 days on others.

FLOWERS, ORNAMENTALS, & SHRUBS: Mix 3 teaspoonsful in one gallon water. Make application when insects appear, and repeat as necessary to control aphids, mealy bugs, spider mites, white flies, and leafhoppers. To control scale on euonymus, juniper, and pine, spray in May. To control bagworm larvae, spray as soon as insects are seen and repeat in 10 to 14 days. Do not use on Boston or Maidenhair ferns, pteris, crassula, or canaerti juniper as injury may result.

EPA REC. NO. 33955-394-AA

EPA EST. NO. 2217-KS-1

893-175

ACME®

60-52

malathion 50% spray for garden insects

ACTIVE INGREDIENTS:

Malathion* 50.00%

Xylene 38.51%

INERT INGREDIENTS 11.49%

TOTAL 100.00%

*O,O-Dimethyl dithiophosphate of diethyl mercaptosuccinate.

WARNING:

Keep Out of Reach of Children
See Side Panels for Additional Warnings

NET CONTENTS 8 OZ ES

WARNING: Harmful if swallowed. Avoid breathing of spray mist. Avoid contact with skin. Wash thoroughly after use. Avoid contamination of feed or foodstuffs. Keep away from heat or open flames.

This product is toxic to fish. Keep out of lakes, streams, and ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues. Protective information may be obtained from your Co-operative Agricultural Extension Service.

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME. DO NOT RE-USE CONTAINER. DESTROY WHEN EMPTY.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and conditions of application equipment will vary, Seller makes no claims other than those stated on this label.

DISTRIBUTED BY

ACME DIVISION

GORDON CORPORATION

KANSAS CITY, MO. 64118

7414)


DIRECTIONS FOR USE

VEGETABLES: Mix 3 tsp. in 1 gal. water to control spiders, mites, and aphids on tomatoes; Mexican Bean Beetle, spider mites, Leafhoppers, and aphids on beans; Pea aphid on peas; aphids on potatoes; aphids and Imported Cabbage Worms on cabbage, broccoli, kale, mustard, and turnip. Spray as needed. Do not use within 7 days of harvest on cabbage, kale, and mustard, 3 days on others.

FLOWERS, ORNAMENTALS, & SHRUBS: Mix 3 tsp. in one gal. water. Make application when insects appear, and repeat as necessary to control aphids, mealy bugs, spider mites, white flies, and leafhoppers. To control scale on euonymus, juniper, and pine, spray in May. To control bagworm larvae, spray as soon as insects are seen and repeat in 10 to 14 days. Do not use on Boston or Maidenhair ferns, pteris, crassula, or canaerti juniper as injury may result.

893-175

EPA REG. NO. 33955-394-AA
EPA EST. NO. 2217-KS-1



ACTIVE INGREDIENTS:

Malathion*	50.00%
Xylene	38.51%

INERT INGREDIENTS

TOTAL	100.00%
-------	---------

*O,O-Dimethyl dithiophosphate
of diethyl mercaptosuccinate.

WARNING:

Keep Out of Reach of Children
See Side Panels for Additional Warnings

NET CONTENTS 4 OUNCES

WARNING: Harmful if swallowed. Avoid breathing of spray mist. Avoid contact with skin. Wash thoroughly after use. Avoid contamination of feed or foodstuffs. Keep away from heat or open flames.

This product is toxic to fish. Keep out of lakes, streams, and ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DO NOT USE, POUR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME. DO NOT RE-USE CONTAINER. DESTROY WHEN EMPTY

ACME D. S. CO.
GORDON CORPORATION