

acme[®]
fruit tree
spray
 controls:
INSECTS / DISEASES*

**CONTROLS INSECTS / DISEASES on Apples,
 Peaches, Plums, Prunes and Cherries**

**CAUTION - Keep Out of Reach of Children.
 Read cautions also on label.**

TWELVE OUNCES NET WEIGHT

ACCEPTED
 3955-408
 NOV 27 1974

acme fruit tree spray

DIRECTIONS: The contents of this package will make 24 gallons of commercial dose spray. For smaller quantities use *5 to 7 level* tablespoonfuls to each gallon of water or *4 to 6 level* teaspoonfuls to a quart. Mix well. Shake sprayer occasionally while spraying to keep wettable powder in suspension. A light thorough application as directed below is better than heavy irregular sprays.

APPLES, PEARS: To control scab, black rot, sooty blotch, Brooks spot, aphids, mites, codling moth, Forbes scale, curculio, apple maggot, pear slug, Japanese beetle, and leaf roller apply as follows: 1. When blossom stems separate and before blossoms open. 2. When 75% of the petals have fallen. 3. 10 days later. 4. 10 days to 2 weeks later. 5. Continue at 2-week intervals until 7 days before harvest. May injure Bosc pears in the East.

PEACHES, APRICOTS: To control brown rot, scab, mites, curculio, Oriental moth, Japanese beetle, and rose chafer apply as follows: 1. When 75% of petals have fallen. 2. When shucks begin to split. 3. 10 days later. 4. 10 days later. 5. 2 weeks later. 6. 21 days before harvest. 7. When brown rot continues to be a problem apply ACME WETTABLE DUSTING SULPHUR at 10-day intervals right up to harvest.

PLUMS AND PRUNES: To control mites, curculio, Japanese beetle, apply as follows: 1. When 75% of petals have fallen. 2. When shucks are falling. 3. 10 days later. 4. 2 weeks later. 5. Continue at 2-week intervals if necessary. 6. 21 days before harvest.

CHERRIES: To control leaf spot, brown rot, aphids, curculio, fruit fly, and slugs apply as follows: 1. When 75% of petals have fallen. 2. When shucks are falling from expanding fruit. 3. 2 weeks later. 4. When fruit is beginning to color but 21 days before harvest. 5. When disease continues to be a problem follow with ACME WETTABLE DUSTING SULPHUR weekly right up to harvest. 6. Spray immediately after harvest to maintain vigor of trees.

EPA REG. NO. 33955-408-AA
EPA EST. NO. 2217-KS-1

CAUTION: Harmful if swallowed. Avoid inhaling dust particles or spray mist. Avoid repeated or prolonged contact with skin, wash thoroughly. Keep out of reach of children. Avoid contamination of feed or foodstuffs. Do not apply to fruit closer to harvest than indicated for each crop. Remove residues at harvest by washing, wiping, brushing, or other effective means.

This product is toxic to fish. Keep out of lakes, streams, and ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label.

This product highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your cooperative Agricultural Extension Service.

NOTE: Where apple maggots are severe add 2 tablespoonfuls of ACME SEVIN 50W to at least two applications in July.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

SOLD FOR INSECTICIDE AND FUNGICIDE PURPOSES • DO NOT APPLY IN ANY MANNER NOT SPECIFIED ON THE LABEL • DO NOT REUSE CONTAINER. DESTROY WHEN EMPTY.

DISTRIBUTED BY
ACME DIVISION
PBI - GORDON CORPORATION
KANSAS CITY, KANSAS 66118

LD 255 Z14 525-474

228