

ACME®

33-62

fruit tree spray

controls:
insects / diseases*

**CONTROLS INSECTS/DISEASES on Apples,
Pears, Peaches, Plums, Prunes and Cherries.**

Active Ingredients

Methoxychlor Technical	12%
Malathion (O,O-Dimethyl phosphorothioate of dimethyl mercaptosuccinate)	5%
Captan (N-Hydroxymethyl-2-pyrone)	12%
Other Ingredients	70%
TOTAL	100%

*12% Methoxychlor Technical is equivalent to 10.56%
22, bis (p-methoxyphenyl) 1,1,1-trichloroethane and
1.44% other isomers and reaction products.

CAUTION: Keep Out of Reach of Children.

See back panel for additional cautions.

NET WEIGHT FOUR POUNDS

ACCEPTED

MAY 09 1975

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No.

33955-408

292

DIRECTIONS FOR USE

The contents of this package will make 50 gallons of commercial dose spray. For smaller quantities use 5 to 7 level tablespoonfuls to each gallon of water or 4 to 6 level teaspoonfuls to a quart. Mix well. Shake sprayer occasionally while spraying to keep wettable powder in suspension. A light thorough application as directed below is better than heavy irregular sprays.

APPLES, PEARS: To control scab, black rot, sooty blotch, Brooks spot, aphids, mites, codling moth, Forbes scale, curculio, apple maggot, pear slug, Japanese beetle, and leaf roller, apply as follows: 1. When blossom stems separate and before blossoms open. 2. When 75% of the petals have fallen. 3. 10 days later. 4. 10 days to 2 weeks later. 5. Continue at 2-week intervals until 7 days before harvest. May injure Bosc pears in the East.

PEACHES, APRICOTS: To control brown rot, scab, mites, curculio, Oriental moth, Japanese beetle, and rose chafer, apply as follows: 1. When 75% of petals have fallen. 2. When shucks begin to split. 3. 10 days later. 4. 10 days later. 5. 2 weeks later. 6. 21 days before harvest. 7. When brown rot continues to be a problem apply ACME WETTABLE DUSTING SULPHUR at 10-day intervals right up to harvest.

PLUMS AND PRUNES: To control mites, curculio, Japanese beetle, apply as follows: 1. When 75% of petals have fallen. 2. When shucks are falling. 3. 10 days later. 4. 2 weeks later. 5. Continue at 2-week intervals if necessary. 6. 21 days before harvest.

CHERRIES: To control leaf spot, brown rot, aphids, curculio, fruit fly, and slugs, apply as follows: 1. When 75% of petals have fallen. 2. When shucks are falling from expanding fruit. 3. 2 weeks later. 4. When fruit is beginning to color but 21 days before harvest. 5. When disease continues to be a problem follow with ACME WETTABLE DUSTING SULPHUR weekly right up to harvest. 6. Spray immediately after harvest to maintain vigor of trees.

CAUTION: Harmful if swallowed. Avoid inhaling dust particles or spray mist. Avoid contact with skin, wash thoroughly. Keep out of reach of children. Avoid contamination of feed or foodstuffs. Do not apply to fruit closer to harvest than indicated for each crop. Remove residues at harvest by washing, wiping, brushing, or other effective means.

This product is toxic to fish. Keep out of lakes, streams, and ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Apply this product only as specified on this label.

This product highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

NOTE: Where apple maggots are severe, add 2 tablespoonfuls of ACME SEVIN 50W to at least two applications in July.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, Seller makes no claims other than those stated on this label.

DO NOT REUSE EMPTY BAG. DESTROY BY BURNING (STAY OUT OF SMOKE).

EPA REG. NO. 33955-408-AA
525-475

EPA EST. NO. 2217-KS-1

