

PM 12 33955-408

194



ACCEPTED
 SEP 12 1986
 Under the Federal Insecticide, Fungicide, and Herbicide Act, as amended, for the pesticide registered under EPA Reg. No. 33955-408

*CONTROLS INSECTS/DISEASES on Apples, Pears, Peaches, Plums, Prunes and Cherries.

ACTIVE INGREDIENTS:

Methoxychlor Technical**	12.00%
Malathion (0,0-Dimethyl dithiophosphate of diethyl mercaptosuccinate)	6.00%
Captan***	11.35%
Related Derivatives	0.65%
INERT INGREDIENTS	70.00%
TOTAL	100.00%

**12% Methoxychlor Technical is equivalent to 10.56% 2,2-bis (p-methoxyphenyl)-1, 1,1-trichloroethane and 1.44% other isomers and reaction products.


***[N-trichloromethylthio-4-cyclohexene-1, 2-dicarboximide].

KEEP OUT OF REACH OF CHILDREN

D A N G E R

See Side Panels for Statement of Practical Treatment and Additional Precautionary Statements.

NET WEIGHT TWO POUNDS



284

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

DANGER: Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Wear goggles or face shield when handling. Avoid contact with skin and clothing. Remove and separately launder clothing before reuse.

Statement of Practical Treatment

IF IN EYES: Wash thoroughly with water for at least 15 minutes and contact a physician if necessary.

IF ON SKIN: In case of contact wash with soap and water.

IF INHALED: Move to fresh air and call emergency medical care.

IF SWALLOWED: Seek immediate medical attention if ingestion has occurred.

ENVIRONMENTAL HAZARDS

Fish: This pesticide is toxic to fish. Do not apply directly to water or wetlands, swamps, bogs, marshes, and potholes. Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes.

Bees: This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

Endangered Species Restrictions: The use of this product in a manner which may harm endangered species or adversely modify their habitat is prohibited by Federal Law. The following endangered species and the general location of their habitats have been identified by county:

- Slackwater Darter, Alabama (Lauderdale, Limestone and Madison)
Tennessee (Lawrence and Wayne)
- Pecos gambusia, New Mexico (Chaves and Eddy)
Texas (Reeves, Jeff Davis and Pecos)
- Freshwater Mussels, Tennessee (Hancock, Claiborne, Hawkins and Sullivan)
Virginia (Smyth, Scott, Lee, Washington & Grayson)
- Commanche Springs Pupfish, Texas (Reeves and Jeff Davis)

Prior to use of this product in the above counties you must determine that these species are not located in, or immediately adjacent to, the areas to be treated with captan. Consult your regional U.S. Fish and Wildlife Service Office (Endangered Species Specialist) or State Wildlife Agency, if there is any doubt as to the areas where the outdoor use of captan is prohibited.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL:
 Store in original container in a locked storage area inaccessible to children and pets. For disposal, securely wrap original container in several layers of newspaper and discard in trash. Do not reuse empty container. Do not contaminate water, food, or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Wear natural or synthetic rubber, vinyl, or plastic gloves impermeable to captan when using. Do not use indoors. When using outdoors, wear long pants and long-sleeved shirt and apply with the wind to your back. Wash nondisposable gloves thoroughly with soap and water before removing. Clothing worn while handling this product must be laundered separately from other clothing before reusing.

Do not rotate a foliar treated crop with crops other than those with registered captan uses.

Do not allow persons to enter treated areas within 4 days following application unless protective clothing is worn. Conspicuously post reentry information at site of application.


Do not apply to fruit closer to harvest than indicated for each crop. Remove residues at harvest by washing, wiping, brushing, or other effective means.

The contents of this package will make 24 gallons of commercial dose spray. For smaller quantities use 5 to 7 level tablespoonfuls to each gallon of water or 4 to 6 level teaspoonfuls to a quart. Mix well. Shake sprayer occasionally while spraying to keep wettable powder in suspension. A light thorough application as directed below is better than heavy irregular sprays.

APPLES, PEARS: To control scab, black rot, sooty blotch, Brooks spot, aphids, mites, codling moth, Forbes scale, curculio, apple maggot, pear slug, Japanese beetle, and leaf roller, apply as follows: 1. When blossom stems separate and before blossoms open. 2. When 75% of the petals have fallen. 3. 10 days later. 4. 10 days to 2 weeks later. 5. Continue at 2-week intervals until 7 days before harvest. May injure Bosc pears in the East.

PEACHES, APRICOTS: To control brown rot, scab, mites, curculio, Oriental moth, Japanese beetle, and rose chafer, apply as follows: 1. When 75% of petals have fallen. 2. When shucks begin to split. 3. 10 days later. 4. 10 days later. 5. 2 weeks later. 6. 21 days before harvest. 7. When brown rot continues to be a problem apply ACME WETTABLE DUSTING SULPHUR at 10-day intervals right up to harvest.

PLUMS AND PRUNES: To control mites, curculio, Japanese beetle, apply as follows: 1. When 75% of petals have fallen. 2. When shucks are falling. 3. 10 days later. 4. 2 weeks later. 5. Continue at 2-week intervals if necessary. 6. 21 days before harvest.



4034

CHERRIES: To control leaf spot, brown rot, aphids, curculio, fruit fly, and slugs, apply as follows: 1. When 75% of petals have fallen. 2. When shucks are falling from expanding fruit. 3. 2 weeks later. 4. When fruit is beginning to color but 21 days before harvest. 5. When disease continues to be a problem follow with ACME WETTABLE DUSTING SULPHUR weekly right up to harvest. 6. Spray immediately after harvest to maintain vigor of trees.

NOTE: Where apple maggots are severe, add 2 tablespoonfuls of ACME SEVIN 50W to at least two applications in July.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatability with other chemicals, and condition of application equipment will vary. Seller makes no claims other than those stated on this label.

525/586

EPA Reg. No. 33955-408-AA
EPA EST. NO. 2217-KS-1

