

ACME FRUIT TREE SPRAY  
Controls Insects/Diseases on Apples, Peaches,  
Plums, Prunes and Cherries.

ACTIVE INGREDIENTS:

Methoxychlor Technical* . . . . .	12.00%
Malathion: O,O-dimethyl phosphorodithioate of diethyl mercaptosuccinate . . . . .	6.00%
Captan** . . . . .	11.74%
Related Derivatives. . . . .	0.26%
INERT INGREDIENTS . . . . .	70.00%
	TOTAL 100.00%

\*12% Methoxychlor Technical is equivalent to 10.56% 2,2-bis  
(p-methoxyphenyl)-1,1,1-trichloroethane and 1.44% other isomers  
and reaction products.

\*\* (N-trichloromethylthio-4-cyclohexene-1,2-dicarboximide).

KEEP OUT OF REACH OF CHILDREN


DANGER

See Back Panel for Statement of Practical Treatment  
and Additional Precautionary Statements.

NET WEIGHT TWELVE OUNCES

525/389

AP



EPA REG. NO. 33955-408

EPA EST. NO. 2217-KS-X

Mfd. by ACME DIVISION  
PBI/GORDON CORPORATION  
KANSAS CITY, KS 66118

294

**STOP!** READ THE ENTIRE LABEL FIRST.  
OBSERVE ALL PRECAUTIONS AND  
FOLLOW DIRECTIONS CAREFULLY.

**PRECAUTIONARY STATEMENTS**

**Hazards to Humans and Domestic Animals**

**DANGER:** Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Wear goggles or face shield when handling. Avoid contact with skin and clothing. Remove contaminated clothing and separately launder before reuse.

**Statement of Practical Treatment**

**IF IN EYES:** Flush with plenty of water. Call a physician.

**IF ON SKIN:** In case of contact, wash with soap and water.

**IF INHALED:** Remove victim to fresh air. If not breathing give artificial respiration preferably mouth-to-mouth. Get medical attention.

**IF SWALLOWED:** Drink promptly a large quantity of milk, egg whites, gelatin solutions, or if these are not available, drink large quantities of water. Avoid alcohol.

**Environmental Hazards**

This is toxic to fish, aquatic invertebrates, and aquatic life stages of amphibians. Do not apply directly to water or wetlands (i.e., swamps, bogs, marshes, and potholes). Drift and runoff may be hazardous to aquatic organisms in areas near the application site. Do not clean equipment or dispose of equipment washwaters in a manner that will contaminate water resources. This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

**DIRECTIONS FOR USE**

It is a violation Federal law to use this product in a manner inconsistent with its labeling.

**STORAGE & DISPOSAL**

**STORAGE:** Store in original container in a locked storage area inaccessible to children and pets.

**PESTICIDE DISPOSAL:** Securely wrap original container in several layers of newspaper and discard in trash.

**CONTAINER DISPOSAL:** Do not reuse empty container. Rinse thoroughly before discarding in trash.

**BEST AVAILABLE COPY**

Do not apply this product through any type of irrigation system.

Wear goggles or face shield when handling. Remove contaminated clothing and separately launder clothing before reuse. Wear chemical resistant gloves when using. Do not use indoors. When using outdoors, wear long pants and long-sleeved shirt and apply with the wind to your back. Wash nondisposable gloves thoroughly with soap and water before removing.

Do not apply to fruit closer to harvest than indicated for each crop. Remove residues at harvest by washing, wiping, brushing, or other effective means.

Mix well. Shake sprayer occasionally while spraying to keep wettable powder in suspension. A light thorough application as directed below is better than heavy irregular sprays. With hand sprayers, occasionally stir or shake the mixture to keep particles in suspension.

On small or dwarf trees 5 to 6 feet high, apply approximately 1 to 2 pints of dilute spray per tree. For midsize trees, approximately 1 to 2 quarts per tree and for large trees apply 1 to 3 gallons per tree.

APPLES: To control scab, black rot, sooty blotch, Brooks spot, aphids, mites, Codling moth, Forbes scale, curculio, apple maggot, pear slug, Japanese beetle, and leaf roller, for apples mix at the rate of 6 1/3 level tbsps. per gallon of water and apply as follows:

1. When blossom stems separate and before blossoms open.
2. When 75% of the petals have fallen.
3. 10 days later.
4. 10 days to 2 weeks later.
5. Continue at 2-week intervals until 7 days before harvest. May injure Bosc pears in the East.

PEACHES, APRICOTS: To control brown rot, scab, mites, curculio, Oriental moth, Japanese beetle, and rose chafer, for peaches mix at the rate of 6 1/3 level tbsps. per gallon of water, for apricots mix at the rate of 4 2/3 tbsps. per gallon of water and apply as follows:

1. When 75% of petals have fallen.
2. When shucks begin to split.
3. 10 days later.
4. 10 days later.
5. 2 weeks later.
6. 21 days before harvest.
7. When brown rot continues to be a problem apply ACME WETTABLE DUSTING SULPHUR at 10-day intervals right up to harvest.

BEST AVAILABLE COPY

PLUMS (FRESH PRUNES): To control mites, curculio, Japanese beetle, for plums mix at the rate of 4 2/3 tbsps. per gallon of water and apply as follows:

1. When 75% of petals have fallen.
2. When shucks are falling.
3. 10 days later.
4. 2 weeks later.
5. Continue at 2-week intervals if necessary.
6. 21 days before harvest.

CHERRIES: To control leaf spot, brown rot, aphids, curculio, fruit fly, and slugs, for cherries mix at the rate of 3 level tbsps. per gallon of water and apply as follows:

1. When 75% of petals have fallen.
2. When shucks are falling from expanding fruit.
3. 2 weeks later.
4. When fruit is beginning to color but 21 days before harvest.
5. When disease continues to be a problem follow with ACME WETTABLE DUSTING SULPHUR weekly right up to harvest.
6. Spray immediately after harvest to maintain vigor of trees.

NOTE: Where apple maggots are severe, add 2 tablespoonfuls of ACME SEVIN 50W to at least two applications in July.

LIMITED WARRANTY: Manufacture warrants that the chemical composition conforms to the ingredient statement given on the label and that this product is suited for the labeled use when applied according to label directions. Because of widely varying use conditions, it is impossible to eliminate all risks even when label directions are followed.

Exclusion of Other Warranties and Remedies. Except where such disclaimers and exclusions are specifically prohibited by applicable law. THE FOREGOING IS THE ONLY GUARANTEE OR WARRANTY APPLICABLE TO THIS PRODUCT AND IS GIVEN EXPRESSLY AND IN LIEU OF ALL OTHER WARRANTIES EXPRESSED OR IMPLIED, OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, AND ALL SUCH WARRANTIES WHICH EXCEED OR DIFFER FROM SAID LIMITED WARRANTY ARE DISCLAIMED BY MANUFACTURER, and, upon Manufacturer's compliance with said limited warranty, BUYER SHALL HAVE NO REMEDY AGAINST MANUFACTURER FOR ANY TYPE OF DAMAGE OR LOSS, and, IN NO EVENT SHALL MANUFACTURER BE LIABLE FOR CONSEQUENTIAL DAMAGE OR LOSS.