

33955-450

11/01/1974

1/2

Best Available Copy

ACCEPTED
33955-450
 NOV 1 1974

UNDER FEDERAL PESTICIDE
 FUNGICIDE AND WEEDKILLER ACT
 FOR ECONOMIC PERSON REGISTERED
 UNDER NO. _____ SUBJECT
 TO ATTACHED COMMENTS.

CHINCHBUG
CONTROL

ACME®

TWO POUNDS
NET WEIGHT

Controls

Japanese Beetles, Chinchbugs, Mexican Bean Beetles, Squash Bugs, and certain other insect pests of fruit, vegetables, ornamentals, and turf.

CAUTION: Keep out of reach of children
See back panel for additional cautions.

ACTIVE INGREDIENTS:	
Carbaryl (1-naphthyl N-methylcarbamate)	50.0%
INERT INGREDIENTS:	
	50.0%
Total	
	100.0%

*Trade Mark Union Carbide Corporation for 1-naphthyl N-methylcarbamate, U.S. Patent 2,993,478.

DIVISION OF
 ACME DIVISION
GORDON CORPORATION
 KANSAS CITY, KANSAS 64116

Best available copy

2/2

DIRECTIONS FOR USE

ACNE SEVIN[®] 50W is a wettable spray powder. Measure in level tablespoons. Keep spray mixture well agitated. Do not mix with lime, Bordeaux Mixture or other alkaline materials. It is compatible with most fungicides and miticides. Do not mix with oil base sprays.

TUPE: Fall army worms, soil web worms, or lawn worms, ants, earwigs, fleas, millipedes, mosquitoes, and chinch-bugs, use one cupful in 15 to 20 gallons of water per 500 square feet of lawn. Repeat as needed especially for chinchbug. Also, treat along curbs, and edges of sidewalks. This same dosage sprayed on flower beds, has given some control of slug damage.

ORNAMENTALS: Use 2 tablespoons per gallon for Japanese beetles, flea beetles, Japanese weevils, leatherjackets, elm leaf beetles, scarab beetles, flower beetles, leaf miners, plant hoppers, grasshoppers, ground crickets, tree crickets, box elder bugs and tent caterpillars. Apply when pests appear. Repeat as needed.

APPLES AND PEARS: Use two tablespoons per gallon for codling moth, pear psylla, apple maggot, plum curculio, fruit tree leaf rollers, Japanese beetles, apple sprays and San Jose scale. Do not apply before second or third spray for codling moth. Apply at 10 day intervals in July for apple maggot. May be used up to one day before picking. This tag makes 100 gallons of spray.

ALMONDS, WALNUTS, PECAN, CHERRIES: Use two to four tablespoons per gallon for peach leaf curl, fruit tree leaf roller, San Jose scale, codling moth, fruit tree worm and excrement. Higher dosage. Do not apply after buds split.

PEACHES, APRICOTS, NECTARINES AND ALMONDS: Use one tablespoon per gallon for Oriental fruit moth, plum curculio, peach twig borer, leaf rolling bugs, San Jose scale, Japanese beetles. May be used one day before harvest or earlier, 3 days on apricots and almonds.

SLUG DAMAGE: Use two tablespoons per gallon up to one day before picking for leaf rollers and meadow webbug control. Make three to five applications as needed. Do not apply more than 200 gallons per acre per application.

CHERRIES, PLUMS AND PEACHES: Use 2 tablespoons per gallon in this tag with 10 gallons of water for Cherry Fruit Fly, Fruit Worm, Peach Twig Borer, Mealy Plum Borer, Plum Curculio, Red Banded Leaf Roller, and Japanese Beetles. Do not apply closer than one day before picking.

NOTE ON VEGETABLES: Do not apply more than 100 gallons per acre per application. See specific tags for beans, peas, etc. Use two tablespoons per gallon for fly, aphid, leaf miner, webworm, etc. 10 days. For Mexican bean beetle, leatherjackets, flea beetles, slink bugs, and earwigs, use up to one day before picking. Do not apply more than 100 gallons per acre per application.

CABBAGE, BROCCOLI, CARNIFLOWER, BRUSSELS SPROUTS: Use two to four tablespoons per gallon for flea beetle, Harlequin bug, corn earworm, cabbage worm, and diamond borer. Stop spraying 3 days before picking.

CORN: Use two to four tablespoons per gallon for corn earworm, fall army worm, and grasshopper. Apply up to 10 days before harvest. Repeat at 10 day intervals. Do not apply more than 200 gallons per acre per application.

CUCUMBER, MELONS, AND SQUASH: Use two to four tablespoons per gallon for cucumber beetle, squash vine borer, and squash bug. Do not apply more than 100 gallons per acre per application.

PEAS: Use 2 tablespoons per gallon for Army worms, grasshoppers and leatherjackets. May be used up to one day of picking. Do not feed beetle larvae to stock.

TOMATO, PEPPERS, EGGPLANTS: Use 2 tablespoons per gallon for tomato fruit worm, flea beetles, Fall army worms, slink bugs and leatherjackets. Apply up to day of picking.

CAUTION: Harmful if swallowed or inhaled! Do not take internally. Do not breathe dust. Avoid breathing of spray. Contact with skin may be harmful. Wash thoroughly after use. Wear regular long sleeved work clothing and change to clean clothing daily. Avoid contamination of feet and foodstuffs. Keep children and pets off treated areas until the insecticide is washed into the soil and grass is thoroughly dry.

NOTE TO PHYSICIAN:—Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidote. Keep out of lakes, streams, and ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on treated plants. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DO NOT REUSE EMPTY BAG. DESTROY BY BURNING (stay out of smoke). Buy or discard residues in a safe place away from water supplies.

EPA REG. NO. 33955-450-AA

EPA EST. NO. 2217-KS-1

10-295-07

520-474