

ACME® SEVIN® 50 W INSECTICIDE

A wettable powder for the control of Japanese beetle, Chinch bug, Mexican bean beetle, Squash bug, and certain other insect pests of vegetables, fruit, shrubs, trees.

ACTIVE INGREDIENT:

Carbaryl(1-naphthyl N-methylcarbamate)..... 50%

INERT INGREDIENTS..... 50%


TOTAL 100%

SEVIN® is the registered trademark of Rhône-Poulenc for carbaryl insecticide.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for Precautionary Statements and Statement of Practical Treatment



NET WEIGHT ONE POUND

620/ APXXXXXX
 EPA Reg. No. 33955-450
 EPA Est. No. 2217-KS-1
 Manufactured By ACME DIVISION
 PBI/GORDON CORPORATION
 KANSAS CITY, MISSOURI 64101

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: May be harmful if swallowed. Avoid breathing vapors (dust or spray mist). Avoid contact with eyes, skin or clothing. Wash thoroughly after handling.

- User Safety Requirements:**
- ◆ Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when mixing and applying this product.
 - ◆ Wear a hat and eye protection when making overhead applications.
 - ◆ Remove clothing immediately if pesticide soaks clothing. Change clothing as soon as possible after use.
 - ◆ Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.
 - ◆ Do not allow children or pets to contact treated area until sprays have dried.

Statement of Practical Treatment

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting.

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

Note to Physician: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine sulfate is antidotal. Do not use 2-PAM, opiates, or cholinesterase inhibiting drugs.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic invertebrates. ~~For terrestrial uses, Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.~~ Do not contaminate water when disposing of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL

STORAGE: Store in original container in a locked storage area inaccessible to children or pets. Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Discard container in trash.

GENERAL INFORMATION AND USE PRECAUTIONS

Acme Sevin 50W[®] Insecticide is a wettable spray powder. Measure in level tablespoonfuls. Keep spray mixture well agitated. Do not mix with lime, Bordeaux Mixture, or other alkaline material. This product is compatible with most fungicides and miticides. Do not mix with oil base sprays.

- ◆ Apply when air is calm to avoid drift and contact with eyes and skin.
- ◆ Start applying at the farthest corner of the treatment area and work backward to avoid contact with wet surfaces. Allow spray to dry in treated areas before reentering.
- ◆ For trees taller than 10 feet, consider hiring a licensed professional.
- ◆ Spray thoroughly to wet upper and lower leaf surfaces, stems and branches.
- ◆ Do not repeat application more than once a week.
- ◆ Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl (Sevin) label(s).

TREES AND SHRUBS: Mix 2 tablespoons of product per gallon of water. Apply when pests appear. Repeat as needed. For trees taller than 10 feet, consider hiring a licensed professional. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.

Trees & Shrubs	Pests
Evergreens:	Bagworm
Elm:	Elm leaf beetle, Elm leaf aphid, scale crawlers.
Oak:	Spring cankerworm, leaf rollers.
Rose:	Rose slug, leaf rollers, rose aphid, exposed thrips, Japanese beetle.
Azaleas:	Lace bug, mealybug, blister beetle.
Boxelder:	Boxelder bug.
Flowering Cherry:	Leaf roller, black cherry aphid, Japanese beetle, brown soft scale, tent caterpillar, orange tortrix.

APPLES AND PEARS: Mix 2 tablespoons of product per gallon of water to control codling moth, pear psylla, apple maggot, plum curculio, fruit tree leaf rollers, Japanese beetle, apple aphid, and San Jose scale. Do not apply before second cover spray to avoid fruit thinning. Make applications at 10 to 14 day intervals throughout the season. Apply at 10 day intervals in July to control apple maggot. May be used up to one day before picking. For trees taller than 10 feet, consider hiring a licensed professional. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.

PEACHES AND APRICOTS: Use 1 tablespoon of product per gallon of water to control oriental fruit moth, plum curculio, peach twig borer, "cat-facing" bug, San Jose scale, and Japanese beetle. May be used up to one day before harvest on peaches; 3 days on apricots. For trees taller than 10 feet, consider hiring a licensed professional. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.

GRAPES: Use 1 tablespoon of product per gallon of water to control leafhoppers, grape berry moth, and Japanese beetle. Make first application before bloom, repeat on 14 to 20 day schedule. Make one application just before berries touch in the clusters. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.

STRAWBERRIES: Use 2 tablespoons of product per gallon of water up to one day before picking to control leaf rollers and meadow spittlebug. Make three to five post-bloom summer sprays, as needed. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.

CHERRIES, PLUMS, AND PRUNES: Use 2 tablespoons of product per gallon of water to control cherry fruit fly, fruitworm, peach twig borer, mealy plum aphids, plum curculio, red banded leaf roller, and Japanese beetle. Do not apply closer than one day before picking. For trees taller than 10 feet, consider hiring a licensed professional. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.

BEANS: Use 2 tablespoons of product per gallon of water to control Mexican bean beetle, leafhoppers, flea beetle, stink bugs, lima bean pod borer. -Spray until all foliage is wet. Repeat in 7 to 10 days. May be used up to harvest. Do not feed vines to dairy or meat animals. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.

CABBAGE, BROCCOLI, CAULIFLOWER, AND BRUSSELS SPROUTS: Use 2 tablespoons of product per gallon of water to control flea beetle, harlequin bug, corn earworm, cabbageworm, but not cabbage looper. Treat weekly. Stop spraying 3 days before cutting. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.

CORN: Use 2 tablespoons of product per gallon of water to control corn earworm, corn borer, fall armyworm, and grasshoppers. Repeat at 2-to-3 7 day intervals to control corn earworm, or until silks are dry. Do not feed treated fodder to animals. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.

CUCUMBERS, MELONS, AND SQUASH: Use 2 tablespoons of product per gallon of water to control squash bug, melon worm, cucumber beetle, and pickleworms. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.

PEAS: Use 2 tablespoons of product per gallon of water to control armyworm, grasshoppers, and leafhoppers. May be used up to one day before picking. Do not feed treated vines to stock. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.

POTATOES: Use 2 tablespoons of product per gallon of water to control Colorado potato beetle, leafhoppers, and flea beetle. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.

TOMATOES, PEPPERS, AND EGGPLANT: Use 2 tablespoons of product per gallon of water to control tomato fruitworm, flea beetle, fall armyworm, stink bugs, and leafhoppers. Apply up to one day before picking. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.

LIMITED WARRANTY AND DISCLAIMER.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.