

Label Copy No: LSC-14133

Date: 5/23/67

Product: S-W Dimite

Package: Can

Size: Gallon(s)

(See Panel)

SHERWIN-WILLIAMS PRODUCTS

 D I M I T E

 25% EMULSIFIABLE

A Specific Control for Certain
Mites and Red Spiders

ACCEPTED

JUL 27 1967

UNDER THE FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. 577-43 SUBJECT
TO ATTACHED COMMENTS.

33955-451

ACTIVE INGREDIENTS:

4,4'-Dichloro alpha-methyl-benzhydrol . . .	27.0%
Xylene	61.0%
Inert Ingredients	11.9%
Total	100.0%

18 PT. - CAUTION: KEEP OUT OF REACH OF CHILDREN - 12 PT.
See Back Panel For Other Cautions - 6 PT.

 D I M I T E kills by residual and contact action.
The residual action may last as long as
one month or more.

** GALLON (S)

U.S. STANDARD
MEASURE

*Trademark

(Back Panel)

SUGGESTIONS FOR USE

DMITE 25% Emulsifiable is formulated to mix readily with water. This material kills by residual and contact action. It does not appreciably reduce beneficial insect populations. The residual activity may last as long as a month or more.

DMITE 25% Emulsifiable is compatible with most commonly used insecticides and fungicides except those of a highly acid nature.

To control European red mite, Scheenii mite, Pacific mite, 2-spotted mite, and red spider on evergreens and ornamentals, use 1 pint of DMITE 25% Emulsifiable per 100 gallons water (1 teaspoonful per gallon water). Make first application as soon as the mites appear and repeat if necessary. In many areas one application may be sufficient to keep mites under control throughout the season.

To control 2-spotted mites on sugar beets, apply 1-2 quarts DMITE per acre in sufficient water to give adequate coverage with the particular spray equipment being used. Apply not later than 60 days before harvest. Do not feed sugar beet tops to livestock. Do not apply more than once per season.

To control 2 spotted mites on dry beans, apply at above dosage but not later than 15 days before harvest. Do not use on green or bush beans. Do not feed straw to livestock. Do not apply more than once per season.

To control 2-spotted mites on seed crops, apply at above dosage. Make first application when mites appear and repeat as necessary. Do not graze seed crops or feed crop waste to livestock.

For early season control of European Red mites on apples, use 1 pint DIMITE in 100 gallons of water and apply in the pink spray. Do not apply when fruit is formed.

CAUTION: Harmful if taken internally. This solution, if brought into repeated or prolonged contact with the skin, may cause toxic symptoms. Avoid inhalation and skin contact. In case of spillage on the skin, wash immediately with soap and water. Avoid contamination of foodstuffs.

Toxic to fish. Keep out of lakes, ponds and streams.

WARNING--FLAMMABLE: Keep away from heat, sparks, and open flame. Keep container closed when not in use. Use with adequate ventilation.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

SOLD FOR MEDICINE PURPOSES

THE SHERWIN-WILLIAMS COMPANY
General Offices: Cleveland, Ohio 44101
Made in U.S.A.

USDA Registration No. 577-451

LSC-14103 **

**Insert month and year of current printing e.g.: 5-67

**One Gallon or
Five Gallons

