

Label Copy No: LS-24538

Date: 1/22/67

Product: SW Standard Arsenate of Lead
(Deflocculating)

Invoice: 1001

Size: 4#

(See Panel)

(For use in Florida only)

SW STANDARD

ARSENATE OF LEAD

Contains Deflocculating Compound

CONTAINS DEFLOCCULATING COMPOUND

THIS PRODUCT CONTAINS:

<u>ACTIVE INGREDIENT:</u>	Arsenate of Lead . . .	90%
<u>INERT INGREDIENTS:</u>	10%
	Total	100%

Total arsenic (expressed as metallic) not less than 20.7%. Water soluble arsenic (expressed as metallic) not more than 0.5%. Lead (expressed as metallic) 87%.

PREPARATION

1. Preparation: See label for directions - 0.17 lb.

R
E
D

ANTIDOTE: CALL A PHYSICIAN OR DRUGGIST. Do not attempt to give fluids to an unconscious person. Give a tablespoonful of salt in a glass of warm water and repeat until vomit fluid is clear. Then give 2 tablespoonfuls of epsom salts or milk of magnesia in water and plenty of milk and water. The victim lie down and keep quiet.

WARNING: Poisonous if swallowed. Avoid contact with eyes, skin, or clothing. Wash thoroughly after using. Do not re-use dust or spray mist. Avoid drift onto adjacent crops and pasture land. Turn all containers, as such, as emptied and stay out of smoke. If on clothing, remove and wash. Keep away from domestic animals, and feedstuffs.

NAME AND ADDRESS

SW STANDARD

1001

Label


SW STANDARD

ARSENATE OF LEAD

Contains Deflocculating Compound

Label

For Spraying Deciduous Fruit Trees, such as apples, pears, cherries, and plums, to control codling moth, cunkerworm, curculionid, apple maggot, and other leaf-rolling insects, use from 2 to 3 pounds of S-W Standard Arsenate of Lead to 100 gallons of water.



Apply at intervals every 1 to 2 days as needed as long as local conditions warrant. Use on tender foliage. Use peach and Japanese plum use from 1 to 1.5 pounds of S-1 Standard Arsenate of Lead combined with 2 to 3 pounds of hydrated lime and 6 pounds of S-1 Standard Sulfate to 100 gallons of water. Do not graze livestock in sprayed orchards.

To control the apple worm, use 4 pounds S-1 Standard Arsenate of Lead in 50 gallons of water and apply at the rate of from 1 to 50 gallons of spray solution to the acre.

For Use on Grapefruit -- Florida only - As a maturity spray on grapefruit, use at the rate of from 1.5 to 2.5 lbs. per 100 gallons for white varieties and from .5 to 1.0 lbs. for pink and red varieties. Apply within 1 to 2 weeks after bloom. The lower dosage is used for a 1:1 ratio of solids to acids in mid-season, the higher dosage is used for a 1:1 ratio in the early season. Do not spray while mature grapefruit from the previous season are present.

CAUTION: Do not apply to tobacco or fruit crops within 30 days of harvest. Remove remaining residue at harvest by washing, wiping, or other effective means. Residues must not be eaten from fruit or tobacco and can be harmful if eaten or consumed.

This material is toxic to wildlife. Do not contaminate any body of water or other natural habitat.

EMPTY CONTAINER: - When empty rinse and cover into tank or pump hole that can be filled. Turn this container at once--stay out of smoke. Carry all boxes or residue.

WARRANTY: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer accepts all risks of any such use.

The Arsenate of Lead has been applied for control to prevent its accidental misuse. The physical and chemical properties are in no way affected.

SELLER'S RESPONSIBILITY

Buyer's Responsibility - 1-1
1-1-2

13-211-1
1-1

Information on this product is available within the U.S. at 1-1-1