

300
17
95261
1

ACELP Division
PBI/Gordon Corporation
1717 West 12th Street
P.O. Box 4050
Kansas City, MO 64101

Attention: S. Johnson

Re: Acelp

Subject: Acelp Amino & Vegetable Dust
EPA Registration No. 35955-456
Your letter dated March 4, 1983

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Herbicide Act, as amended, is acceptable. A stamped copy is enclosed for your records.


Submit (5) copies of the final printed labeling changing the wording in the Bee precaution to "... in the area or if plants or weeds in the treatment area are in bloom."

Sincerely yours,

df
Henry M. Jacoby
Product Manager (21)
Fungicide-Herbicide Branch
Registration Division (25-757)

TS-767:Jacoboy:DCR-17116:WANG-0191C:LN:raven:479-2013:5/5/83

CONCURRENCES							
SYMBOL							
SURNAME							
DATE							



ACCEPTED
with COMMENTS
in EPA Letter Dated:

MAY 25 1983

United States
Environmental Protection Agency
Washington, D.C. 20460
EPA Reg. No.
100-100-0100

Contains ASEVIN, ZINEB & SULFUR.

ACTIVE INGREDIENTS:

Sulfur.....	20.0%
Zineb (zinc ethylene bisdithiocarbamate).....	5.25%
*Carbaryl (1-naphthyl N-methyl- carbamate).....	4.0%
INERT INGREDIENTS.....	70.75%

KEEP OUT OF REACH OF CHILDREN
CAUTION
See side panel for additional
directions and precautions.

12 pt. bold
18 pt. bold

ASEVIN is the registered name of
Chloro-carbonyl-Dimethyl-piperidate
insecticide.

NET WEIGHT 100 LBS (45.45 KG)
NET WEIGHT 50 LBS (22.73 KG)
NET WEIGHT 25 LBS (11.36 KG)

STOP! READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND FOLLOW
DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

-Hazards to Humans & Domestic Animals-

CAUTION: May be harmful if swallowed or inhaled. May cause irritation of eyes, nose, and throat. Avoid breathing dust. Avoid contact with eyes, skin, or clothing. When using, wear protective clothing, i.e. long sleeves, long pants, hat, gloves, covering for face and eyes. Wash thoroughly after use and launder clothing before reuse. Remove any residues at harvest by brushing, washing or peeling. Do not feed treated crops, pods, or husks to dairy animals or animals being readied for slaughter.

-Environmental Hazards-

This product is toxic to fish. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Bee Precaution: This product is highly toxic to bees exposed to direct treatment or residues on crops. Do not dust when bees are active in the area or plants in full bloom. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

12 pt.
bold

→ STORAGE & DISPOSAL (for the 10 ounce squeeze duster label)

Store away from children, pets, and foodstuffs. This squeeze duster is an applicator/container package. The container may be refilled and used again and again, but only with Acme Tomato & Vegetable Dust. WHEN NO LONGER WANTED OR NO LONGER SERVICEABLE, WRAP AND PLACE IN TRASH COLLECTION.

12 pt.
bold

→ STORAGE & DISPOSAL (for the one pound shaker canister label)

Store away from children, pets, and foodstuffs. DO NOT REUSE EMPTY CONTAINER. WRAP AND PLACE IN TRASH COLLECTION.

12 pt.
bold

→ STORAGE & DISPOSAL (for the 4 lb. bag label)

Store away from children, pets, and foodstuffs. DO NOT REUSE EMPTY BAG. WRAP AND PLACE IN TRASH COLLECTION.

585/383

EPA REG. NO. 33955-456
EPA EST. NO. 2217-KS-1

Distributed by
Acme Division
PBI/GORDON CORPORATION
Kansas City, Kansas 66119

Limited Warranty. Manufacturer warrants that the chemical composition conforms to the ingredient statement given on the label and that this product is suited for the labeled use when applied according to label directions. Because of widely varying use conditions, it is impossible to eliminate all risks even when label directions are followed.

Exclusion of Other Warranties and Remedies. Except where such disclaimers and exclusions are specifically prohibited by applicable law, THE FOREGOING IS THE ONLY GUARANTEE OR WARRANTY APPLICABLE TO THIS PRODUCT AND IS GIVEN EXPRESSLY AND IN LIEU OF ALL OTHER WARRANTIES EXPRESS OR IMPLIED, OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, AND ALL SUCH WARRANTIES WHICH EXCEED OR DIFFER FROM SAID LIMITED WARRANTY ARE DISCLAIMED BY MANUFACTURER, and, upon Manufacturer's compliance with said limited warranty, BUYER SHALL HAVE NO REMEDY AGAINST MANUFACTURER FOR ANY TYPE OF DAMAGE OR LOSS, and, IN NO EVENT SHALL MANUFACTURER BE LIABLE FOR CONSEQUENTIAL DAMAGE OR LOSS.

HOW TO USE

(for the 10 ounce squeeze duster label)

Lift cap and squeeze sidewalls to dust. This product is ready-to-use and is packaged in a refillable applicator/container. This container may be refilled and used again and again but only with Acme Tomato & Vegetable Dust. A 4 lb. bag is available for refills. Simply remove the bottom plug and refill. Do not use teaspoons or tablespoons or other utensils for other purposes once used with pesticides.

HOW TO USE

(for the one pound shaker-top canister label)

Shake from the shaker top or pry off the top and apply with a suitable dusting gun. Do not use teaspoons or tablespoons or other utensils for other purposes once used with pesticides.

HOW TO USE

(for the 4 lb. bag label)

Apply with a suitable dusting gun or use this package to refill the 10 ounce squeeze duster applicator/container this product is also available in. Remove the bottom plug of the squeeze duster and refill. Do not use teaspoons or tablespoons or other utensils for other purposes once used with pesticides.

HOW TO USE

(continuation of copy for all three labels)

Make application in morning or evening when it is cool. Treat upper and lower surfaces of all foliage. Do not apply in excess. Start applications at first sign of insects or disease. Repeat as necessary, generally at 7 day intervals, oftener in rainy weather. Do not use closer to harvest than indicated following each crop.

(continuation of copy for 10 ounce label only)

Ten ounces will treat 375 to 750 feet of row depending on plant size and degree of infestation.

(continuation of copy for one pound label only)

One pound will treat 600 to 1200 feet of row depending on plant size and degree of infestation.

(continuation of copy for 4 lb. label only)

Four pounds will treat 2400 to 4800 feet of row depending on plant size and degree of infestation.

(continuation of copy for all three labels)

Light application should be made in seed beds if plants show signs of disease or insects.

FOR CONTROL OF

Time to Stop Use
Before Harvest

BEANS Mexican bean beetle, leafhoppers, flea beetles, corn earworm, lygus and stinkbugs, powdery mildew, rust and Anthracnose

7 days

bold
bold

CARROTS Leafhoppers, flea beetles, spittlebugs, early and late blight

7 days

bold

SWEET CORN Corn earworms (See Notes 5 & 6), fall army worm, grasshoppers, Helminthosporium leaf blight (See Notes 1 and 2)

day of harvest

bold

CABBAGE AND CAULIFLOWER Flea beetles, cabbage loopers, imported cabbageworms and Diamond Back moths (See Note 7), harlequin bugs, corn earworm leaf spot

7 days

bold

BRUSSELS SPROUTS and BROCCOLI Flea beetles, cabbage worms, harlequin bugs, corn earworm, leaf spot

7 days

bold

CUCUMBERS Pickleworms, squash bugs, flea beetles, cucumber beetles, Anthracnose, powdery mildews, Alternaria blight (See Note 3)

5 days

bold

MELONS AND SQUASH Pickleworms, squash bugs, flea beetles, cucumber beetles, Anthracnose, powdery mildews, Alternaria blight (See Note 3)

10 days

bold

LETTUCE Leafhoppers, army worms, stinkbugs, downy mildew (See Note 4)

10 days

bold

PEPPERS Tomato fruitworm, flea beetles, Fall army worm, leafhoppers, lygus bug, Anthracnose

5 days

bold

EGGPLANT Lace bugs, tomato hornworm, mites, Anthracnose, cercospora leaf spot, early and late blight

5 days

bold

TOMATOES Early and late blight, Septoria leaf spot, Anthracnose, mites, tomato fruitworm, flea beetles, tomato hornworm, stinkbugs

5 days

bold

NOTE 1 Do not feed forage to dairy animals or animals being finished for slaughter

NOTE 2 Apply to corn silks every 3 days until dry for earworm control

NOTE 3 Do not use closer than 7 days on Summer squash. Do not apply to melons or cucumbers when temperatures exceed 80° F or when vines are wet

NOTE 4 Do not use on leaf lettuce closer than 14 days before picking

NOTE 5 Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population

NOTE 6 Apply to silks as soon as they appear. Make 3 to 4 applications at 2-3 day intervals as infestation warrants. Re apply immediately after rain.

NOTE 7 Where the cabbage looper is the principal problem use an alternate pesticide specifically recommended for it. For control of small cabbage loopers after edible parts start to form, apply at 5 to 7 day intervals

bold
bold

bold