

300
17
153517
3

FBI/Gordon Corporation
P.O. Box 4090
Kansas City, MO 64101-9984

SEP 17 1985

Attention: Mr. Stephen Johnson

Gentlemen:

Subject: Acme Tomato & Vegetable Dust
EPA Registration No. 33955-456
Your Application Dated March 18, 1985

The labeling submitted in response to this Agency's letter of August 6, 1984, regarding labeling requirements in accordance with the Carbaryl Registration Standard issued March 30, 1984, is acceptable provided that the following revisions are made:

1. Revise replanting statement to read "labels," deleting label(s)."
2. Add to label the following reentry interval statement: "Do not reenter treated crop area prior to 24 hours after application." (see enclosed A-117).
3. The warranty statements must be revised in accordance with enclosed EPA Memorandum dated July 31, 1978.

A stamped copy of the label is enclosed for your records.

Submit five (5) copies of your final printed label before you release the product for shipment. Refer to the A-79 enclosure for further description of final printed labeling.

Sincerely yours,

Henry M. Jacoby
Product Manager (21)
Fungicide-Herbicide Branch
Registration Division (TS-767)

Enclosures (4)

RD/PFB; JOB: 88242; Jacoby; RD-68; KENDY; Co: 898-1270; 9/11/85; Del: 9/20/85; DKD

SYMBOL ▶							
SURNAME ▶							
DATE ▶							



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

— Hazards to Humans & Domestic Animals —
CAUTION: May be harmful if swallowed or inhaled. May cause irritation of eyes, nose, and throat. Avoid breathing dust. Avoid contact with eyes, skin, or clothing. When using, wear protective clothing, i.e. long sleeves, long pants, hat, gloves, covering for face and eyes. Wash thoroughly after use and launder clothing before reuse. Remove any residues at harvest by brushing, washing or peeling. Do not feed treated crops, pods, or husks to dairy animals or animals being readied for slaughter.

— Environmental Hazards —
 This product is toxic to fish. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE
 It is a violation of federal law to use this product in a manner inconsistent with its labeling.

STORAGE: Store in original container in a locked, dry storage area inaccessible to children or pets. Do not store with food or feedstuffs. Keep container tightly closed.

DISPOSAL: If can contains unused material, securely wrap in original container in several layers of newspaper and discard in trash. Do not reuse empty can. Discard in trash.

GENERAL INFORMATION (How to Use):
 Shake from this shaker-top can or pry off the top and apply with a suitable dusting gun. Do not use teaspoons or tablespoons or other food utensils for other purposes after used with pesticides.

ONE POUND WILL TREAT 600 to 1,200 FEET OF ROW depending on plant size and degree of infestation.

Make application in morning or evening when it is cool. Treat upper and lower surfaces of all foliage. Do not apply in excess. Start applications at first sign of insects or disease. Repeat as necessary, generally at 7 day intervals, oftener in rainy weather. Do not use closer to harvest than indicated following each crop. Light application should be made in seed beds if plants show signs of disease or insects.

Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl label(s). This product contains carbaryl insecticide.

EPA REG. NO. 33055-456
 EPA EST. NO. 2217-KS-1 585/285

ACCEPTED
 with COMMENTS
 in EPA Letter Dated:



SEP 17 1985

Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 33055-456

ACME® 83-86

tomato & vegetable dust

controls certain insects and diseases of vegetables

Contains *Sevin, Zineb & Sulfur.

ACTIVE INGREDIENTS:

Sulfur	20.0%
Zineb (zinc ethylene bisdithiocarbamate)	5.2%
*Carbaryl (1-naphthyl N-methyl-carbamate)	3.0%
INERT INGREDIENTS	71.8%
TOTAL 100.0%	

KEEP OUT OF REACH OF CHILDREN

CAUTION

See side panels for additional precautionary statements.

*SEVIN is the registered trademark of Union Carbide Corporation for carbaryl insecticide.

NET WEIGHT ONE POUND

FOR CONTROL OF

BEANS: Mexican bean beetle, beetles, corn earworm, fly, powdery mildew, rust and

CARROTS: Leafhoppers, flea early and late blight

SWEET CORN: Corn earworm, Fall army worm, grasshopper, leaf blight (See Notes 1 & 2)

CABBAGE and CAULIFLOW: cabbage loopers, imported Diamond Back moths (See Note 3), corn earworm, leaf spot

BRUSSELS SPROUTS and beetles, cabbage worms, h corn earworm, leaf spot

CUCUMBERS: Pickleworms, beetles, cucumber beetles, mildews, Alternaria blight

MELONS and SQUASH: Pick bugs, flea beetles, cucurbit Anthracnose, powdery mildew (See Note 3)

LETTUCE: Leafhoppers, army downy mildew (See Note 4)

PEPPERS: Tomato fruitworm, army worm, leafhoppers, h Anthracnose

EGGPLANT: Lace bugs, tomato Anthracnose, cercospora blight

TOMATOES: Early and late spot, Anthracnose, mites, beetles, tomato hornworm

NOTE 1. Do not feed forage finished for slaughter

NOTE 2. Apply to corn silks control.

NOTE 3. Do not use clover. Do not apply to me exceed 80° F or w

NOTE 4. Do not use on le picking.

NOTE 5. Application of this during the pollen population.

NOTE 6. Apply to silks as a cations at 2-3 day apply immediately

NOTE 7. Where the cabba an alternate pest For control of ar start to form, appl

Limited Warranty. Manufacturer forms to the ingredient statements suited for the labeled use when of widely varying use conditions when label directions are followed. Exclusion of Other Warranties. Claims and exclusions are stated. FOREGOING IS THE ONLY WARRANTY TO THIS PRODUCT AND IS OTHER WARRANTIES EXPRESS OR FITNESS FOR A PARTICULAR PURPOSE WHICH EXCEED OR DIFFER FROM THIS STATEMENT ARE DISCLAIMED BY MANUFACTURER with said limited warranty. BUYER MANUFACTURER FOR ANY EVENT SHALL MANUFACTURER BE LIABLE FOR DAMAGE OR LOSS.

to dairy animals or animals being reared for slaughter
 — Environmental Hazards —
 This product is toxic to fish. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.
 This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE
 It is a violation of federal law to use this product in a manner inconsistent with its labeling.

ORAGE: Store in original container in a locked, dry storage area inaccessible to children or pets. Do not store with food or food utensils. Keep container tightly closed.
NOTE: This container is an applicator package and may be refilled again, but only with ACME TOMATO & VEGETABLE DUST.
DISPOSAL: If package contains unused material, securely wrap in several layers of newspaper and discard in trash. When package is no longer serviceable, or no longer wanted, discard empty container in trash.

GENERAL INFORMATION (How to Use):
 Shake from this shaker-top can or pry off the top and apply with a suitable dusting gun. Do not use teaspoons or tablespoons or other food utensils for other purposes after used with pesticides.

TEN OUNCES WILL TREAT 375 TO 750 FEET OF ROW depending on plant size and degree of infestation.

Make application in morning or evening when it is cool. Treat upper and lower surfaces of all foliage. Do not apply in excess. Start applications at first sign of insects or disease. Repeat as necessary, generally at 7 day intervals, often in rainy weather. Do not use closer to harvest than indicated following each crop. Light application should be made in seed beds if plants show signs of disease or insects.

Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl label(s). This product contains carbaryl insecticide.

Vegetable dust

controls certain insects and diseases of vegetables

Contains *Sevin, Zineb & Sulfur.

ACTIVE INGREDIENTS:

Sulfur	20.0%
Zineb (zinc ethylene bisdithiocarbamate)	5.2%
*Carbaryl (1-naphthyl N-methyl-carbamate)	3.0%
INERT INGREDIENTS	71.8%
TOTAL	100.0%

KEEP OUT OF REACH OF CHILDREN
CAUTION
 See side panels for additional precautionary statements.

*SEVIN is the registered trademark of Union Carbide Corporation for carbaryl insecticide.

NET WEIGHT 10 OUNCES

BRUSSELS SPROUTS and BROCCOLI Flea beetles, cabbage worms, harlequin bugs, corn earworm, leaf spot	7 days
CUCUMBERS Pickleworms, squash bugs, flea beetles, cucumber beetles, Anthracnose, powdery mildews, Alternaria blight (See Note 3)	5 days
MELONS and SQUASH Pickleworms, squash bugs, flea beetles, cucumber beetles, Anthracnose, powdery mildews, Alternaria blight (See Note 3)	10 days
LETTUCE: Leafhoppers, army worms, stinkbugs, downy mildew (See Note 4)	10 days
PEPPERS: Tomato fruitworm, flea beetles, Fall army worm, leafhoppers, lygus bug, Anthracnose	5 days
EGGPLANT: Lace bugs, tomato hornworm, mites, Anthracnose, cercospora leaf spot, early and late blight	5 days
TOMATOES: Early and late blight, Septoria leaf spot, Anthracnose, mites, tomato fruitworm, flea beetles, tomato hornworm, stinkbugs	5 days

- NOTE 1** Do not feed forage to dairy animals or animals being finished for slaughter.
- NOTE 2** Apply to corn silks every 3 days until dry for earworm control.
- NOTE 3** Do not use closer than 7 days on summer squash. Do not apply to melons or cucumbers when temperatures exceed 80° F or when vines are wet.
- NOTE 4** Do not use on leaf lettuce closer than 14 days before picking.
- NOTE 5** Application of this product to theassel region of corn during the pollen shed period will seriously reduce bee population.
- NOTE 6** Apply to silks as soon as they appear. Make 3 to 4 applications at 2-3 day intervals as infestation warrants. Re-apply immediately after rain.
- NOTE 7** Where the cabbage looper is the principal problem use an alternate pesticide specifically recommended for it. For control of small cabbage loopers after edible parts start to form, apply at 5 to 7 day intervals.

EPA REG NO 33955-456
 EPA EST NO 2217-KS-1

585 285



ACCEPTED with COMMENTS to EPA Letter 7/2/85

SEP 17 1985

Under the Federal Insecticide, Fungicide, and Rodenticide Act
 registered pesticide
 registered under F.F.C. Reg. No. 33955-456

Limited Warranty. Manufacturer warrants that the chemical composition conforms to the ingredient statement given on the label and that this product is suited for the labeled use when applied according to label directions. Because of widely varying use conditions, it is impossible to eliminate all risks even when label directions are followed.

Exclusion of Other Warranties and Remedies. Except where such disclaimers and exclusions are specifically prohibited by applicable law, THE FOREGOING IS THE ONLY GUARANTEE OR WARRANTY APPLICABLE TO THIS PRODUCT AND IS GIVEN EXPRESSLY AND IN LIEU OF ALL OTHER WARRANTIES EXPRESS OR IMPLIED, OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, AND ALL SUCH WARRANTIES WHICH EXCEED OR DIFFER FROM SAID LIMITED WARRANTY ARE DISCLAIMED BY MANUFACTURER, and, upon Manufacturer's compliance with said limited warranty BUYER SHALL HAVE NO REMEDY AGAINST MANUFACTURER FOR ANY TYPE OF DAMAGE OR LOSS, and, IN NO EVENT SHALL MANUFACTURER BE LIABLE FOR CONSEQUENTIAL DAMAGE OR LOSS.