

33455-462

9-27-1974

1/2

FOUR POUNDS

ACME

NET WEIGHT



Controls:

Mexican bean beetles, Japanese beetles, Squash bugs and certain other insect pests of fruit, tobacco, vegetables, ornamentals, and animals.

ACTIVE INGREDIENT:

Carbaryl (*1-naphthyl N-methyl-carbamate)	5.0%
---	------

INERT INGREDIENTS	95.0%
-------------------------	-------

TOTAL	100.0%
-------	--------

*Trade Mark Union Carbide Corporation for 1-naphthyl N-methylcarbamate. U.S. Patent 2,903,478.

CAUTION: KEEP OUT OF REACH OF CHILDREN

See back panel for other cautions.

DISTRIBUTED BY
ACME DIVISION
PBI - GORDON CORPORATION
KANSAS CITY, KANSAS

SEP 27 1974

INDEXED
FEDERAL INSECTICIDE
RODENTICIDE ACT
ECONOMIC POISON REGISTERED
UNDER NO. SUBJECT
TO ATTACHED COMMENTS

33955-462

DIRECTIONS FOR USE

Read the Cautions! Make application in morning or evening when it is cool. Treat upper and lower surfaces of all foliage. Do not apply in excess. Start application at first sign of insects. Repeat as necessary, generally at 7-day intervals, oftener in rainy weather. Do not use closer to harvest than indicated following each crop. Light application should be made in seed beds if plants show signs of insects.

FOR CONTROL OF

Time to Stop Use Before Harvest

Beans: Mexican bean beetle, leafhoppers, flea beetles, corn earworm, lygus and stinkbugs.....	No Limit
Carrots: Leafhoppers, flea beetles and spittlebugs.....	No Limit
Corn: Corn earworms, Fall army worm. (See Notes 1, 2 & 4).....	Day of Harvest
Cabbage and Cauliflower: Flea beetles, imported cabbage worms, diamond-back moth larvae, harlequin bugs, corn earworm.....	3 Days
Brussels Sprouts and Broccoli: Flea beetles, imported cabbage worms, diamond-back moth larvae, harlequin bugs, corn earworm.....	3 Days
Cucumbers, Melons and Squash: Pickleworms, squash bugs, flea beetles, cucumber beetles. (See Note 3).....	Day of Harvest
Lettuce: Leafhoppers, army worms, stinkbugs.....	Head 3 Days
Okra: Stinkbugs, corn earworm.....	No Limit
Potatoes: Colorado potato beetle, flea beetle, leafhopper.....	No Limit
Peppers: Tomato fruitworm, flea beetles, Fall army worm, leafhoppers, lygus bug.....	Day of Harvest
Eggplant: Lace bugs, tomato hornworm.....	Day of Harvest
Tomatoes: Tomato fruitworm, flea beetle, tomato hornworm, stinkbugs.....	Day of Harvest
Strawberries: Meadow spittlebug, leaf roller and strawberry weevil.....	1 Day
Tobacco: Tobacco flea beetles, apply 1 to 2 lbs. per 100 square yards when plants come up. Repeat in 10-14 days. Again just before transplanting from seed bed.....	No Limit

- Note 1. Do not feed forage to dairy animals or animals being finished for slaughter.
- Note 2. Apply to corn silks every 3 days until dry for earworm control.
- Note 3. Do not use on watermelon in Florida. Do not apply to melons or cucumbers when temperatures exceed 80° F or when vines are wet.
- Note 4. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population.

Cats and Dogs: Fleas and ticks on cats and dogs. Rub into skin, avoid eyes. Dust quarters weekly. Do not treat kittens or puppies under 4 weeks of age.

Chickens: Fleas, lice, mites, and fowl ticks. One pound is enough to treat 100 chickens. It can also be used on pheasants, pigeons, and partridge. Dust the chicken house and roosts for mites and fowl ticks with one pound per 40 square feet. Direct dust to vent and fluff areas. Also treat head and neck for stick-tight fleas. Repeat in 4 weeks if needed. Force dust into cracks and crevices. Do not dust birds within 7 days of slaughter. Avoid dusting of nests or eggs. Do not use on pet birds.

CAUTION: Harmful if swallowed or inhaled. Do not take internally. Avoid breathing of dust. Contact with skin may be harmful. Wash hands before eating and take shower or bath after working with this insecticide. Wear regular long sleeved work clothing and change to clean clothing daily.

NOTE TO PHYSICIAN—Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

KEEP OUT OF REACH OF CHILDREN AND ANIMALS

Do not treat kittens or puppies under 4 weeks of age

33955-462

This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Keep out of lakes, streams, and ponds. Do not contaminate water by cleaning of equipment, or disposal of wastes.

CONTAINER DISPOSAL: Do not reuse empty bag. Destroy by burning (stay out of smoke). Bury or discard residues in a safe place.

EPA REG. NO. 33955-462-AA
LD 304 FB

EPA EST. NO. 2217-KS-1
611-974