

ACME[®]

SEVIN[®] 5% DUST

FOR RESIDENTIAL USE ONLY

Controls Mexican bean beetle, armyworm, squash bug, and grasshoppers on fruits and vegetables.

ACTIVE INGREDIENT:

Carbaryl (1-naphthyl N-methylcarbamate) 5%

INERT INGREDIENTS: 95%

TOTAL

ACCEPTED
OCT 19 2004
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 33955-462

SEVIN[®] is the registered trademark of Rhône-Poulenc Nederland B.V.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for Precautionary Statements and First Aid.

NET WEIGHT: ONE POUND or FOUR POUNDS

611/ AP-----
EPA REG. NO. 33955-462
EPA Est. No. 2217-KS-2
MANUFACTURED BY:



Telephone: 1-800-821-7925

2/0



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with eyes, skin or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

User Safety Requirements:

- ◆ Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product.
- ◆ Change clothing as soon as possible after use.
- ◆ Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.
- ◆ Do not allow children or pets to contact treated area until dust has settled.

First Aid

If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.

NOTE TO PHYSICIAN: Carbaryl is a N-methyl carbamate insecticide which is a cholinesterase inhibitor. Specific antidote consists of atropine. Narcotics and other sedatives should not be used. Drugs like 2-PAM (pyridine-2-aldoxime methiodide) are NOT recommended.

ENVIRONMENTAL HAZARDS:

This pesticide is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production.

For use on plants intended for home fruit and vegetable gardens, or for aesthetic purposes.

3/6

STORAGE & DISPOSAL

STORAGE: Store in original container in a locked storage area inaccessible to children or pets.

CONTAINER DISPOSAL:

If empty: Do not reuse this container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

GENERAL INFORMATION & USE PRECAUTIONS:

- ◆ Start application at first sign of insects. Repeat at 7 day intervals.
- ◆ Do not use closer to harvest than indicated following each crop.
- ◆ Wearing household latex or rubber gloves, use a dust applicator such as a shaker can, dust gun, puffer or other bulbous duster, rotary duster or other appropriate duster as available. Dust lightly to cover both upper and lower leaf surfaces with a thin, even film of dust.
- ◆ Apply when air is calm to avoid drift and contact with eyes and skin.
- ◆ Start applying at the farthest corner of the treatment area and work backward to avoid contact with dusted surfaces. Allow dust to settle in treated areas before reentering.
- ◆ For plants above chest height, consider using a liquid carbaryl product.
- ◆ Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl (Sevin) labels.

- INSECT CONTROL -

Vegetables and Small Fruit	Insects	Maximum Application Rate, Ozs. of Product/1,000 Sq.Ft.	Retreatment Interval, Days	Preharvest Interval (PHI), Days
Beans (fresh & dried - snap, navy, & kidney); Pea (fresh & dried); Cowpea; Lentil	Mexican bean beetle, bean leaf beetle, cucumber beetle, flea beetle, Japanese beetle, grasshoppers, leafhoppers, velvetbean caterpillar, western bean cutworm, armyworm, corn earworm, tarnished plant bug, and stink bug.	11 ounces per 1,000 sq.ft.	7 days	PHI = 3 days (succulents) PHI = 21 days (dried) PHI = 14 days (forage) PHI = 21 days (hay)
Carrots	leafhoppers, flea beetle and spittlebug.	15 ounces per 1,000 sq.ft.	7 days	PHI = 7 days
Brussel sprouts; Broccoli; Cabbage; Cauliflower; Chinese cabbage; Collards; Kale; Kohlrabi	flea beetle, harlequin bug, armyworm, cabbage caterpillar, grasshoppers, corn earworm, imported cabbageworm, and diamond-back moth larvae	15 ounces per 1,000 sq.ft.	7 days	PHI = 3 days (Brussel sprouts, broccoli, cabbage, kohlrabi, cauliflower) PHI = 14 days (Chinese cabbage, kale, collards)
Sweet Corn Only (See footnotes 1 & 2)	corn earworm, corn rootworm (adults), European corn borer, fall armyworm, flea beetle, Japanese beetle, sap beetle, grasshoppers, and leafhoppers.	15 ounces per 1,000 sq.ft.	3 days	PHI = 2 days (ears) PHI = 14 days (forage) PHI = 48 days (fodder)
Cucumber; Melon; Pumpkin; Squash; Okra (See footnote 3)	pickleworm, melonworm, cucumber beetle, flea beetle, leafhoppers, grasshoppers, squash bug, stinkbug and corn earworm	7 ounces per 1,000 sq.ft.	7 days	PHI = 3 days

0/4

Continued....

Vegetables and Small Fruit	Insects	Maximum Application Rate, Ozs. of Product/1,000 Sq.Ft.	Retreatment Interval, Days	Preharvest Interval (PHI), Days
Potato	Colorado potato beetle, flea beetle, leafhoppers, European corn borer, fall armyworm, lace bug, tomato fruitworm, tomato hornworm, grasshoppers, tarnished plant bug, and stink bug.	15 ounces per 1,000 sq.ft.	7 days	PHI = 7 days
Tomato; Eggplant; Pepper	Colorado potato beetle, flea beetle, leafhoppers, European corn borer, fall armyworm, lace bug, tomato fruitworm, tomato hornworm, grasshoppers, tarnished plant bug, and stink bug.	15 ounces per 1,000 sq.ft.	7 days	PHI = 3 days
Lettuce (head & leaf); Endive; Celery; Dandelion; Parsley; Spinach; Swiss Chard; Hanover Salad, Rhubarb; Garden Beets (tops only); Turnips (tops only)	flea beetle, harlequin bug, leafhoppers, armyworm, cabbage caterpillar, corn earworm, spittlebug, grasshoppers, tarnished plant bug and stink bug.	15 ounces per 1,000 sq.ft.	7 days	PHI = 14 to 15 days
Strawberry	meadow spittlebug, strawberry leaf roller, and strawberry weevil	15 ounces per 1,000 sq.ft.	7 days	PHI = 7 days

FOOTNOTE:

- 1) Do not feed forage to dairy animals or animals being finished for slaughter.
- 2) Application of this product to the corn tassel during the pollination period will seriously reduce bee population.
- 3) Do not use on watermelon in Florida. Do not apply to melon or cucumber when temperatures exceed 80°F or when vines are wet.

5/6

6/10

LIMITED WARRANTY AND DISCLAIMER.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.