

Label Copy No: LD-30828

Date: 7-28-69

Product: Acme Rose and Flower Spray

Package: Aerosol Can

Size: 13-3/4 Oz.

N. Y. F. D. C. of A. No. 2302
Do Not Puncture, Incinerate
or Store Above 110°F.

CAUTIONS: Harmful if
swallowed. Avoid breathing
spray mist. May be irritating to
skin. Avoid placing fingers into
spray when pressing nozzle.
Remove from skin by washing
with soap and water. Keep con-
tainer away from children at all
times. Do not place on stove or
store in direct sunlight, or where
temperature is over 110° F. as
can may explode. Do not spray
on plastics or asphalt tile. Do
not spray on pets including fish
and birds.

SOLD FOR INSECTICIDE PURPOSES

*MGK 264 Synergist U.S. Pat. 2,476,512
**Equivalent to 0.264% 2,2-Bis(p-methoxy-
phenyl)-1,1,1-trichloroethane.
***U.S. Patent 2,349,772
****Karatane T.M. Rohm & Haas Co.

Made in U.S.A. for
ACME INSECTICIDE DIVISION
The Sherwin-Williams Co.
8250 St. Antin Avenue
Detroit, Michigan 48211

LD-30828

NET WEIGHT

13 3/4 OZ.

USDA Reg. No. 577-406

**

ACME

ROSE & FLOWER SPRAY

69-11

**Controls Rose and Flower Insects
Plus Fungicides for Mildew
and Black Spot Prevention.**

ACTIVE INGREDIENTS:

Pyrethrins	0.020%
N-octyl bicycloheptene dicarboximide*	0.300%
Rotenone	0.100%
Other Cube Resins	0.200%
Methoxychlor Technical**	0.300%
Dichlorone*** (2,3-Dichloro-1,4-Naphthoquinone)	0.120%
← 2-(1-Methylheptyl)-4,6-dinitrophenyl crotonate****	0.113%
Other related nitro phenols chiefly 1-methylheptyl-4,6-dinitrophenol	0.012%
Petroleum distillate	0.115%

INERT INGREDIENTS:

TOTAL 100.00%

CAUTION: KEEP OUT OF REACH OF CHILDREN

See left panel for other cautions.

WARNING: FLAMMABLE
KEEP FROM HEAT OR FLAME
CONTENTS UNDER PRESSURE

HOW TO USE: Read all the
directions first! Remove cap.
Hold can upright with nozzle
opening away from you. Keep
fingers out of stream. Hold at
least a foot away from foliage
to prevent freezing. If you wet
the foliage, you are too close.
Do not over-spray. Keep the
nozzle sweeping. Do not use
when foliage is wet. Do not spray
in extreme mid-day temperature
or make application to plants
that lack water. Make repeat
applications weekly. Be careful
on tender new tip growth. For
control of powdery mildew, black
spot, aphids, lace bugs, mites,
ants, white flies and leafhoppers
as well as certain caterpillars.
Direct spray contact will control
Japanese beetles, flower thrips
and rose chafers.

May be used on Roses, Azaleas,
African Violets, Asters, Begonias,
Marigolds, Delphinium, Gladio-
lus, Zinnias and other ornamen-
tals. Do not use on Cyclamen or
Nasturtiums or vegetables.

ACCEPTED WITH COMMENTS