

(Front Panel)

Net Contents

ACCEPTED
 FEB 25 1965
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC POISON REGISTER
 EP UNDER NO. 111

5 Gallons
U.S. Measure

ACME

MCPP HERBICIDE

33-755-413

FOR CONTROL OF CRICKWEED AND OTHER BRADYRIS VEEIDU
IN GOLF GREENS, FAIRWAYS, AND LAWNS.

Active Ingredients:

* Potassium Salt of 2 (4 Chloro
2 methylphenoxy) propionic acid 31.1%

Inert Ingredients:

68.9%

TOTAL 100.0%

* Equivalent to 26.1% 2 (4 Chloro 2 methylphenoxy)
propionic acid.

(18 pt. Type)

CAUTION: Keep out of reach of children
(See other panels for additional cautions)

(12 pt. Type)

Made for

ACME INSECTICIDE DIVISION
ACME QUALITY PAINTS, INC.
DETROIT, MICHIGAN 48211

(Side Panels)

DIRECTION FOR USE: Acme MCPP Herbicide will not injure bent grasses, established lawns, or turf grasses when used as directed. Controls mouse ear chickweed, clover, plantains, Black Medic and Knotweed in turf grasses. Apply when weeds are young and growing vigorously. Do not apply to bent in the heat of the summer. Do not mow for several days before or after treatment. Do not water for one day following treatment.

ESTABLISHED GRASS (PARKWAY AND LAWN): Apply 1.5 to 2.25 fluid ounces per 1000 square feet in 2 1/2 gallons of water (4 to 6 pints in at least 25 gallons of water per acre). Do not spray during long dry spells.

NOTE: If dandelions are a problem, add 2,4-D at the usual rate except on bent grasses where the rate should not exceed 4 oz. acid equivalent per acre.

ESTABLISHED GREEN: Apply 1.5 fluid ounces per 1000 square feet in 2 1/2 gallons of water (4 pints in at least 25 gallons of water per acre) for control of mouse ear chickweed, black medic, plantains and Knotweed.

CAUTIONS: Keep out of reach of children. Keep in original container in storage. Destroy empty container. Do not allow spray, spray mist, visible or invisible vapors, or air containing mist or vapors to contact plants other than those being sprayed. If 2,4-D or MCPP susceptible plants are growing within a distance of 1/2 mile from area being sprayed, make application only when there is a light breeze away from the susceptible plants. Spray drift can be reduced by using high volume sprays under low pressure, coarse droplet nozzles and boom nozzle drops. Susceptible crops include grapes, tomatoes, tobacco, sugar beets, sunflowers, beans, cauliflower and cabbages, boxelder trees, as well as many other field, vegetables, flowers ornamental and fruit crops. Do not contaminate streams, lakes, ponds, irrigation water, or water used for livestock or domestic purposes. Do not store open containers near feeds, seeds, fertilizer or pesticides including insecticides, fungicides, and weed killers other than those containing 2,4-D or brush killers. Sprayer which has contained MCPP should not be used to apply other pesticides to 2,4-D or MCPP susceptible crops. Avoid contact with skin, eyes, and clothing. Wash hands after use.

WARRANTY: Seller's guarantee shall be limited to the terms of the label, and subject thereto the buyer assumes any risk to persons or property arising out of use or handling and accepts the product on these conditions.

Sold for Herbicide Purposes

Product made in Canada

LD _____

USDA Reg. No. 577-_____

Revised per USDA letter Dec. 17, 1964

LPH