

10% Chlordane Granules

3WT USEC NITE

**For Control of Ants,
Sod Webworms (Lawn Moth), and
Grubs in Turf and Flower Beds.**

ACTIVE INGREDIENT:

*Technical Chlordane	10.0%
INERT INGREDIENTS	90.0%
TOTAL	100.0%

*Equivalent to 6% Octachloro 4,7-methanotetrahydroindane and 4% related compounds.

CAUTION

KEEP OUT OF REACH OF CHILDREN
SEE BACK PANEL FOR OTHER CAUTIONS

NET WEIGHT FIVE POUNDS

33955-485

PBI - **GORDON CORPORATION**
DISTRIBUTED BY
ACME DIVISION
KANSAS CITY KANSAS 66118

Made in U.S.A.

Acme® 10% Chlordane Granules

DIRECTIONS FOR USE

LAWNS: To control ants, chiggers, earwigs, sow bugs, cutworms, sod webworms (lawn moths), and white grubs. For ants apply 1 lb. per 1,000 square feet or contents of this bag on 5,000 square feet. Repeat as necessary to maintain control. For better control of earwigs, apply the product along the edges of cement walks, around base of trees and along edges of building foundations. For chiggers, white grubs, Japanese beetle larvae, cutworms, and sod webworms use 2 lbs. per 1,000 square feet. For better control of chiggers and sod webworms (lawn moths) in established lawns, water the lawn before applying insecticide and do not water again for 2-3 days. For better control of subsurface feeding cutworms, white grubs and Japanese beetle larvae, lawn should be watered after application of insecticide. For new lawns, work 2 lbs. per 1,000 square feet into top 2-3 inches of soil while preparing the seed bed. Birds feeding on treated areas may be harmed.

Suggested Spreader Settings for 2 lbs./1,000 square feet:

These are approximate. Each individual spreader may require some adjustment for the proper dosage rate. Use crisscross application. For 1 lb. per 1,000 square feet apply in one direction only.

Scott 38-5	5	Asgrow	5
Sears	5½	Lawn Beauty	3
Wards	4	Jackson	5
	Vaughans	4	

Try to maintain uniform walking speed. Shut off cart while turning.

FLOWER BEDS AND TRANSPLANTS: Work into soil at 2 lbs. per 1,000 square feet as for new turf. Sprinkle around new transplants as set out to control cutworms. Sprinkle along walls, rock borders, and sidewalk edges where ants and sowbugs tend to hide. If ants are observed around trees and shrubs subject to aphid attack, apply a barrier around base of plants (does not control aphids — spray with Acme Malathion 50% Spray for these). Make application with a tin can or sifter box.

BULBS: Protects bulbs against thrips. Before storing gladiolus, tulips sprinkle with granules and stir to coat. Remove soil and dried tops beforehand. If stored in peat moss or vermiculite mulch, sprinkle a small amount of granules over each layer as bulbs are packed.

CAUTION: This material, if swallowed, can cause toxic symptoms. Avoid breathing of dust. Avoid contamination of feed or foodstuffs. Do not use around food crops. Wash with soap and water after using. Keep children and pets off treated areas until material has been watered into soil and grass is dry.

This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams, and ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

DO NOT REUSE EMPTY BAG. DESTROY BY BURNING (STAY OUT OF SMOKE) and bury in a safe place away from water supplies.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment, will vary, Seller makes no claims other than those stated on this label.

bugs, cutworms, sod webworms (lawn moths), and white grubs. For ants apply 1 lb. per 1,000 square feet or contents of this bag on 5,000 square feet. Repeat as necessary to maintain control. For better control of earwigs, apply the product along the edges of cement walks, around base of trees and along edges of building foundations. For chiggers, white grubs, Japanese beetle larvae, cutworms, and sod webworms use 2 lbs. per 1,000 square feet. For better control of chiggers and sod webworms (lawn moths) in established lawns, water the lawn before applying insecticide and do not water again for 2-3 days. For better control of subsurface feeding cutworms, white grubs and Japanese beetle larvae, lawn should be watered after application of insecticide. For new lawns, work 2 lbs. per 1,000 square feet into top 2-3 inches of soil while preparing the seed bed. Birds feeding on treated areas may be harmed.

Suggested Spreader Settings for 2 lbs./1,000 square feet:

These are approximate. Each individual spreader may require some adjustment for the proper dosage rate. Use crisscross application. For 1 lb. per 1,000 square feet apply in one direction only.

Scott 38-5	5	Asgrow	5
Sears	5½	Lawn Beauty	3
Wards	4	Jackson	5
	Vaughans	4	

Try to maintain uniform walking speed. Shut off cart while turning.

EPA Reg. No. 33955-485-AA EPA Est. No. 2217-KS-1

Use to control cutworms. Sprinkle along walks, rock borders, and sidewalk edges where ants and sowbugs tend to hide. If ants are observed around trees and shrubs subject to aphid attack, apply a barrier around base of plants (does not control aphids — spray with Acme Malathion 50% Spray for these). Make application with a tin can or sifter box.

BULBS: Protects bulbs against thrips. Before storing gladiolus, tulips sprinkle with granules and stir to coat. Remove soil and dried tops beforehand. If stored in peat moss or vermiculite mulch, sprinkle a small amount of granules over each layer as bulbs are packed.

CAUTION: This material, if swallowed, can cause toxic symptoms. Avoid breathing of dust. Avoid contamination of feed or foodstuffs. Do not use around food crops. Wash with soap and water after using. Keep children and pets off treated areas until material has been watered into soil and grass is dry.

This product is toxic to fish, birds, and other wildlife. Keep out of lakes, streams, and ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

DO NOT REUSE EMPTY BAG. DESTROY BY BURNING (STAY OUT OF SMOKE), and bury in a safe place away from water supplies.

NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment, will vary, Seller makes no claims other than those stated on this label.

LD 321 G7 536-674

