

ACME WEED-KILLER (EPA), one quart / label copy (continued)



[second side label]

For best results, treat weeds while they are young and actively growing. Woody plants and deep rooted perennials should be resprayed as soon as any regrowth occurs. Dense, tall weed stands should be treated every 7 to 10 days and any skips be sprayed. Depending on density of vegetation, one gallon of spray may cover approximately 150 square feet. Do not over spray as material that falls on the ground is not active. Two light applications just to the growth point is more effective than one heavy spraying.

ADDITION FOR ARSENIC

In case of accident, call a physician immediately. If swallowed, give a tablespoon of salt in a glass of warm water and repeat until vomit fluid is clear. Then give two tablespoonfuls of Epsom salts or Milk of Magnesia in water and plenty of milk and water. Have victim lie down and keep quiet. Do not give fluids to an unconscious person.

CAUTION

Harmful if swallowed. Avoid contact with skin. Avoid breathing spray mist. Wash hands after use. Do not store near food or feed. Wash clothes worn while spraying.

Do not allow children or pets on treated areas. Do not allow domestic animals to graze treated areas. Do not contaminate water for domestic use or irrigation.

Rinse equipment thoroughly after use. Do not sprinkle immediately after application.

DISPOSAL OF USE: when container is empty, rinse with water at least twice. Add this water as part of final dilution. Punch holes in top and bottom so container cannot be reused. Discard in safe place or bury.

WARRANTY: offer warrants that this product conforms to the ingredient statement on the label. No conditions of use such as weather, compatibility with other chemicals, and condition of application equipment will void other warranties unless they offer more than those stated on the label.

ACCEPTED

FEB 19 1975

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. **34572-1**

GENERAL INFORMATION:

Selectar 6 is a postemergence herbicide for use in turf. Selectar 6 contains a surfactant (wetting agent) in proper ratio for best results. It is not necessary to add additional surfactant to the spray tank. Local conditions and recommendations vary; consult your local agricultural experiment station or extension service weed specialists for conditions in your area.

MIXING INSTRUCTIONS:

Selectar 6 herbicide must be thoroughly dissolved. Fill the spray tank with half the required amount of water and, with pump or agitator operating, add the recommended amount of Selectar 6 herbicide and continue filling the tank with the balance of water.

If you prefer, you can premix Selectar 6 herbicide by dissolving the required amount of Selectar 6 in 3 to 5 gallons of water in a clean, openhead bucket. This premix should then be poured in the partially filled spray tank and the tank filled with the balance of water. Mix thoroughly before using.

Do not store spray solution in tank overnight. Clean application equipment thoroughly after use by flushing with water in a safe place.

APPLICATION METHODS:

Apply Selectar 6 with a low volume, properly calibrated sprayer having satisfactory pumping and bypass action. Adjust nozzles to keep spray off susceptible foliage and in a manner to allow maximum coverage of weeds. Adequate coverage is very important for effective weed control. Apply during warm weather when weeds are in an active stage of growth.

SELECTAR NO. 6
Kills Nutgrass
(Nutsedge) Sandbur

ACTIVE INGREDIENT:
Monosodium Acid
Methanearsonate* 48.35%

INERT INGREDIENT: . . 51.65%

TOTAL 100.00%

Total Arsenic, all in water soluble form expressed as elemental 22.38%

*Product contains 6.0 pounds MSMA per gallon.

CAUTION:
KEEP OUT OF REACH OF CHILDREN
See side panel for additional cautions.
READ ENTIRE LABEL BEFORE USING

RHODES, INC.
Lincoln, Nebraska

Net Contents
E.P.A. Reg. No. 34572-1
E.P.A. (Est.) No. 34572-NB-1

DIRECTIONS FOR USE:

ORNAMENTAL TURFGRASS:
Selectar 6 can be used for selective control of smooth and large crabgrass, bahiagrass, barnyardgrass, chickweed, dalliagrass, nutsedge, sandbur and wood sorrel with little or no injury to well-established, actively growing turfgrasses. Mow turfgrass to height of 1 to 1 1/4 inches before treatment. Mix 1 fluid ounce (2 tablespoons) of Selectar 6 in 5 gallons of water for application to 1,000 square feet. Application should be uniform and thorough to adequately wet all undesirable plants. Two or more repeat treatments at 14-day intervals may be necessary. Make applications during warm weather when temperature is between 80 and 90 F. Do not water turf for at least 24 Hr. after application. Turfgrasses may be temporarily discolored. Bermudagrass, bluegrass, and zoysiagrass have shown tolerance to Selectar 6 properly applied. Injury may result if applied to bentgrasses and fescues. Do not apply to St. Augustinegrass, Carpetgrass, centipedegrass or to Dichondra. Do not reseed until 2 weeks after last application.

CAUTION:

KEEP OUT OF REACH OF CHILDREN. Harmful if swallowed. Avoid contact with skin and eyes. In case of contact flush eyes with plenty of water. Avoid breathing spray mist. Wash hands after using. Keep away from children and domestic animals and do not allow them on treated areas until material has been washed into soil. Avoid storage near food and feedstuffs.

DO NOT feed treated forage to livestock or graze treated areas.

DO NOT spray or allow drift onto vegetables, ornamentals or other desirable plants.

DO NOT contaminate waters used for domestic purposes or by animals, including wildlife and aquatic life, or water used for irrigation purposes.

DO NOT reuse empty container. Return to drum reconditioner, or destroy by perforating, crushing and burying or discarding in a safe place.

DO NOT contaminate water by cleaning or equipment, or disposal of wastes. Apply this product only as specified on this label.

HI-YIELD LAWN WEED KILLER

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH LABEL CAUTIONS, PRECAUTIONS AND DIRECTIONS. KEEP PESTICIDE IN ORIGINAL CONTAINER. DO NOT PUT CONCENTRATE OR DILUTE INTO FOOD OR DRINK CONTAINERS.

GENERAL INFORMATION: HI-YIELD LAWN WEED KILLER is a most effective hormone type herbicide that helps control most broadleaf and vining type weeds in home lawns, golf courses and similar turf areas.

WEEDS THIS PRODUCT CONTROLS

Henbit	Kochia	Self-heal
Beggarticks	Lambsquarters	Shepherdspurse
Blessed Thistle	Moneywort	Stinging Nettle
Cats-ear	Mustards	St. Johnswort
Chicory	Pennywort	Thistle, Canadian
Cocklebur	Pepper-grasses	Velvet Weed
Dock, curly	Pigweed	Vervains
Fleabane	Butter Cup	Wild Carrot
Galinsoga	Poison Hemlock	Wild Garlic
Ground Ivy	Buckhorn Plantain	Wild Onion
Morning Glory	Puncturevine	Wintercress
Gum Weed	Ragweed, Common	Witchweed
Hawkweed	Ragweed, Giant	Yarrow
		Yellow Rocket

DIRECTIONS:

Regular Grasses: Mix 8 fl. ozs. in 6 gallons of water. Thoroughly apply to 1,900 sq. ft. (approximately 30' x 63') of lawn or turf area. Heavily infested lawns may require repeat applications to kill missed weeds. Best results are obtained when weeds are young and actively growing. Do not apply during periods of drouth. Do not apply within dripline of desirable trees and shrubs. Repeat in 10-14 days if necessary. Do not water for 24 hours.

St. Augustine: Mix 2 fl. ozs. in 3 gallons of water. Apply to 950 sq. ft. (approximately 30' x 32') of lawn area. Repeat in 10-14 days if necessary. Do not water for 24 hours.

Precaution: May cause possible, temporary injury to St. Augustine. Vapors from this product may injure susceptible plants in the immediate vicinity. Coarse sprays are less likely to drift.

ENVIRONMENTAL CAUTION: May be harmful if swallowed or absorbed through the skin. May cause irritation to eyes and skin. Do not breathe spray mist. Avoid contact with skin, eyes and clothing. In case of contact, flush eyes with plenty of water for at least 15 minutes. Get medical attention. Wash thoroughly after handling. Do not store near fertilizers, seeds, insecticides or fungicides. Do not contaminate food or foodstuffs. Avoid contamination of water intended for irrigation and domestic use. Do not contaminate water by cleaning of equipment, or disposal of wastes.

Direct spray at undesirable plants. Do not use same spray equipment for applying other materials to plants as injury will result. If lawn begins to turn brown temporarily, water heavily. Do not use on Centipede grass, Dichondra or on lawns or turf where desirable Bentgrass or Clovers are present. Do not use on newly seeded lawns or turf. Avoid spray drift to susceptible plants, as this product may injure cotton, peas, beans, grapes, ornamentals and similar broadleaf plants.

Container Disposal: Do not reuse container for any purpose. Destroy when empty.

Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

HI-YIELD CHEMICAL COMPANY

BONHAM, TEXAS

5M-1-115

EPA Reg. No. 34911-9-AA