



ONE POUND

ACME®

NET WEIGHT

CAUTION!

May be harmful if swallowed or inhaled. May cause irritation of eyes, nose and throat. Avoid breathing any dust. Avoid contact with eyes and skin. Wash after using. Do not store near feed or food-stuffs. Remove residues at harvest by brushing, washing or peeling. Do not feed treated crops, pods or vines to dairy animals being finished for slaughter.

Toxic to honeybees. Do not apply to open blossoms or spray in late afternoon. Warn local beekeepers well in advance to move hives to a safe distance until one week after final application.

Do not contaminate any body of water. Do not use in any manner not specified on this label.

DO NOT REUSE CONTAINER.

DESTROY WHEN EMPTY BY BURNING.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

SOLD FOR INSECTICIDE AND FUNGICIDE PURPOSES

DISTRIBUTED BY
ACME DIVISION
GORDON CORPORATION
KANSAS CITY, KANSAS 64116

Made in U.S.A.

EPA REGISTRATION No. 33955-511AA



RED RIVER
VEGETABLE SPRAY
73-44

COMBINED FUNGICIDE AND
INSECTICIDE SPRAY POWDER
FOR CERTAIN VEGETABLES

ACTIVE INGREDIENTS:

- Carbaryl* (1-naphthyl-N-methylcarbamate) 25%
- Zineb (Zinc ethylene bisdithiocarbamate) 38%

INERT INGREDIENTS: 37%
TOTAL 100%

* Sevin, T.M. Union Carbide Corp. for 1-naphthyl-N-methylcarbamate.
U. S. Patent 2,903,478

CAUTION: KEEP OUT OF REACH OF CHILDREN

See left panel for other cautions

Use 1 pound
gallons per ac
spoonfuls per
Use reduced
avoid injury.

TOMATO, EGGS
and late blight
fruitworm, flea
plants are
before harvest

CUCUMBERS.
to 2 gallons of
plants start to
infestation. Ma
To control sc
beetles, downy
and gummy st
Do not use o
before harvest.

BEANS: Mexi
beetles, corn
and Anthracno
insects. Do not
CAUTIONS.

CABBAGE AND
cabbage worm
bug, corn earw
soon as transp
before harvest

EPA EST. NO.

ACCEPTED

AUG 5 1974

UNDE... FEDERAL INSECTICIDE
FUNGICIDE AND RODENTICIDE ACT
FOR ECONOMIC POISON REGISTERED
ED UNDER NO. ... SUBJECT
TO ATTACHED COMMENTS.

33955-511

UTION!

swallowed or inhaled. May
res. nose and throat. Avoid
oid contact with eyes and skin.
not store near feed or food-
ues at harvest by brushing.
o not feed treated crops. pods
animals being finished for

Do not apply to open blos-
-afternoon. Warn local bee-
nce to move hives to a safe
nk after final application.
e any body of water. Do not
specified on this label.

REUSE CONTAINER.
EN EMPTY BY BURNING.

ants that the product conforms
ption and is reasonably fit for
n the label when used in ac-
ons under normal conditions of
arranty nor any other warranty
fitness for a particular purpose.
tends to the use of this product
structions, or under abnormal
conditions not reasonably fore-
buyer assumes the risk of any

IDE AND FUNGICIDE PURPOSES

DISTRIBUTED BY
E DIVISION
I CORPORATION
CITY, KANSAS 66118

Made in U.S.A.
RATION No. 33955-511AA

ONE POUND

ACME

NET WEIGHT

**RED RIVER
VEGETABLE SPRAY**

73-44

**COMBINED FUNGICIDE AND
INSECTICIDE SPRAY POWDER
FOR CERTAIN VEGETABLES**

ACTIVE INGREDIENTS:

- Carbaryl* (1-naphthyl-N-
methylcarbamate) 25%
- Zineb (Zinc ethylene
bisdithiocarbamate) 38%

- INERT INGREDIENTS: 37%
- TOTAL 100%

*Sevin T.M. Union Carbide Corp. for 1-naphthyl-N-methylcarbamate.
U. S. Patent 2,903,478

CAUTION: KEEP OUT OF REACH OF CHILDREN

See left panel for other cautions

Use 1 pound per 12½ gallons of water. Spray 100
gallons per acre or enough to wet all foliage (5 table-
spoonfuls per gallon). Repeat at 5 to 7-day intervals.
Use reduced dosage on cucumbers and melons to
avoid injury.

TOMATO, EGGPLANTS, PEPPERS: To control early
and late blight, Anthracose, Septoria leaf spot, tomato
fruitworm, flea beetle and leafhopper. Start when
plants are 6 inches high. May be used up to 5 days
before harvest. Wash to remove residues.

CUCUMBERS, MELONS, SQUASH: Use 5 tablespoons
to 2 gallons of water (4 lbs./100 gals./acre). Begin when
plants start to run or at first sign of disease or insect
infestation. May be used up to 5 days before harvest.
To control squash bug, melon worms, cucumber
beetles, downy mildew, Anthracose, Alternaria blight
and gummy stem blight, spray undersides of leaves.
Do not use on summer squash closer than 10 days
before harvest.

BEANS: Mexican bean beetles, leafhoppers, flea
beetles, corn earworm, stinkbugs, downy mildew, rust
and Anthracose. Spray at first sign of disease or
insects. Do not use closer than 7 days of harvest. See
CAUTIONS.

CABBAGE AND CAULIFLOWER: Flea beetle, imported
cabbage worm, diamond back moth larvae, harlequin
bug, corn earworm, Alternaria, downy mildew; spray as
soon as transplants are set out. Stop spraying 10 days
before harvest.

LD 348 C5
553-474

EPA EST. NO. 2217-KS-1

one week after final application.
 contaminate any body of water. Do not
 inner not specified on this label.
DO NOT REUSE CONTAINER.
DESTROY WHEN EMPTY BY BURNING.
 This warrant warrants that the product conforms
 to the description and is reasonably fit for
 the uses stated on the label when used in ac-
 cording to the directions under normal conditions of
 use. This warranty nor any other warranty
 of merchantability or fitness for a particular purpose,
 whether implied, extends to the use of this product
 in violation of label instructions, or under abnormal
 conditions or under conditions not reasonably fore-
 seeable, and buyer assumes the risk of any

FOR INSECTICIDE AND FUNGICIDE PURPOSES

DISTRIBUTED BY
ACME DIVISION
ORDON CORPORATION
 KANSAS CITY, KANSAS 66118

Made in U.S.A.
 REGISTRATION No. 33955-511AA

COMBINED FUNGICIDE AND INSECTICIDE SPRAY POWDER FOR CERTAIN VEGETABLES

ACTIVE INGREDIENTS:

Carbaryl* (1-naphthyl-N-
 methylcarbamate) 25%

Zineb (Zinc ethylene
 bisdithiocarbamate) 38%

INERT INGREDIENTS:..... 37%

TOTAL 100%

* Sevin T.M. Union Carbide Corp. for 1 naphthyl N-methyl-carbamate.
 U. S. Patent 2,903,478

CAUTION: KEEP OUT OF REACH OF CHILDREN

See left panel for other cautions

CUCUMBERS:
 To 2 gallons of
 plants start to
 infestation. M
 To control s
 beetles, dcwm
 and gummy s
 Do not use c
 before harvest

BEANS: Mexi
 beetles, corn
 and Anthrax
 insects. Do no
CAUTIONS:

CABBAGE AND
cabbage worm
bug, corn earw
 soon as transp
 before harvest

EPA EST. NO.

