



ACCEPTED
JUN 8 1983
Federal Insecticide,
Fungicide and Rodenticide Act,
Registration for the pesticide
33853-511

COMBINED FUNGICIDE
AND INSECT SPRAY
POWDER FOR CERTAIN
VEGETABLES

ACTIVE INGREDIENTS	
Zineb (Zinc ethylene bisdithiocarbamate)	38%
*Carbaryl (1-naphthyl-N methylcarbamate)	25%
INERT INGREDIENTS	37%
TOTAL 100%	

*SEVIN is the registered trademark of Union Carbide Corporation for carbaryl insecticide.

KEEP OUT OF REACH OF CHILDREN
CAUTION
See side panels for additional
precautionary statements.

← 12 pt. bold
← 18 pt. bold

NET WEIGHT 6 OUNCES

STOP! READ THE ENTIRE LABEL FIRST.
OBSERVE ALL PRECAUTIONS AND FOLLOW
DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

-Hazards to Humans & Domestic Animals-

CAUTION: May be harmful if swallowed or inhaled. May cause irritation of eyes, nose, and throat. Avoid breathing dust. Avoid contact with eyes, skin, or clothing. When using, wear protective clothing, i.e. long sleeves, long pants, hat, gloves, covering for face and eyes. Wash thoroughly after use and launder clothing before reuse. Remove any residues at harvest by brushing, washing or peeling. Do not feed treated crops, pods, or husks to dairy animals or animals being readied for slaughter.

-Environmental Hazards-

This product is toxic to fish. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Bee Precaution: This product is highly toxic to bees exposed to direct treatment or residues on crops. Do not dust when bees are active in the area or plants in full bloom. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

12 pt. bold → STORAGE & DISPOSAL

Store away from children, pets, and foodstuffs. DO NOT REUSE EMPTY CONTAINER. WRAP AND PLACE IN TRASH COLLECTION.

553/383

EPA REG. NO. 33955-511
EPA EST. NO. 2217-KS-1

Distributed By
Acme Division
PPI/GORDON CORPORATION
Kansas City, Kansas 66118

BEST DOCUMENT AVAILABLE

NEED NOT BE USED FOR ANY OTHER PURPOSES, AND ALL SUCH WARRANTIES EXCEPT OR DIFFER FROM SAID LIMITED WARRANTY ARE DISCLAIMED BY MANUFACTURER, and, upon Manufacturer's compliance with said limited warranty, BUYER SHALL HAVE NO REMEDY AGAINST MANUFACTURER FOR ANY TYPE OF DAMAGE OR LOSS, and, IN NO EVENT SHALL MANUFACTURER BE LIABLE FOR CONSEQUENTIAL DAMAGE OR LOSS.

HOW TO USE

Use 5 tbs./gal. water. Spray until all foliage is wet. Repeat at 5 day intervals to protect new foliage. Note reduced dosage for use on cucumbers and melons to avoid injury. Do not use teaspoons or tablespoons or other utensils for other purposes once used with pesticides. Note the time intervals to stop use before harvest.

CROPS

TOMATO, EGGPLANTS, PEPPERS: To control early and late blight Anthracnose, Septoria leaf spot, tomato fruitworm, flea beetle and leafhopper. Start when plants are 6 inches high. May be used up to 5 days before harvest. Wash to remove residues.

CUCUMBERS, MELONS, SQUASH: Use 5 tablespoons to 2 gallons of water. Begin when plants start to run or at first sign of disease or insect infestation. May be used up to 5 days before harvest. To control squash bug, melon worms, cucumber beetles, downy mildew, Anthracnose, Alternaria blight and gummy stem blight, spray undersides of leaves. Do not use on summer squash closer than 10 days before harvest.

BEANS: Mexican bean beetles, leafhoppers, flea beetles, corn earworm, stinkbugs, downy mildew, rust and Anthracnose. Spray at first sign of disease or insects. Do not use closer than 7 days of harvest.

CABBAGE AND CAULIFLOWER: Flea beetle, imported cabbage worm, diamond back moth larvae, harlequin bug, corn earworm, Alternaria, downy mildew. spray as soon as transplants are set out. Stop spraying 10 days before harvest.

USE FOR ALL CROPS

*undersides of leaves
is bold-faced type.*