1OF2 Pri 21 K: G, # 139-- 511

300 <u>15 3562</u> 11

PBI/Gordon Corporation P.O. Box 4090 Kansas City, MO 64101-9984

SEP | 7 1985

Attention: Mr. Stephen Johnson

Gentlemen:

Ń

1

Subject: Acme Red River Vegetable Spray EPA Registration No. 33955-511 Your Application Dated March 18, 1985

The labeling submitted in response to this Agency's letter of August 6, 1984, regarding labeling requirements in accordance with the Carbaryl Registration Standard issued March 30, 1984, is acceptable provided that the following revisions are made:

- 1. Revise replanting statement to read "labels," deleting "label(s)."
- 2. The warranty statements must be revised in accordance with enclosed EPA Memorandum dated July 31, 1978.

A stamped copy of the label is enclosed for your records.

Submit five (5) copies of your final printed label before you release the product for shipment. Refer to the A-79 enclosure for further description of final printed labeling.

Sincerely yours,



Henry M. Jacoby Product Manager (21) Fungicide-Herbicide Branch Registration Division (TS-767)

Enclosures (3)

RD/FHB:JOB:88242:Jacoby:RD-68:Kendrick&Co:898=1270:9/11/85:Del:9/20/85:DKD CONCURRENCES						
SYMBOL .						
SURNAME						
	1					
EPA Form 1320-1 (4-81)	· ·	·	$\sim$		OFFIC	CIAL FILE COPY



#### READ THE ENTIRE LABEL FIRST. **OBSERVE ALL PRECAUTIONS AND FOLLOW** DIRECTIONS CAREFULLY. .

## PRECAUTIONARY STATEMENTS

### - Hazards to Humans & Domestic Animals -

CAUTION: May be harmful if swallowed or inhaled. May cause irritation of eyes, nose, and throat. Avoid breathing dust. Avoid contact with eyes, skin, or clothing. When using, wear protective clothing, i.e. long sleeves, long pants, hat, gloves, covering for face and eyes. Wash thoroughly after use and launder clothing before reuse. Remove any residues at harvest by brushing, washing or peeling. Do not feed treated crops, pods, or husks to dairy animals or animals being readied for slaughter.

#### – Environmental Hazarda —

This product is toxic to fish. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds it bees are visiting the treatment area.

## **DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

## STORAGE & DISPOSAL

STORAGE: Store in original container in a locked, dry storage area inaccessible to children or pets. Do not store with food or feedstuffs. Keep container tightly closed.

DISPOSAL: If can contains unused material, wrap in several tayers of newspaper and discard in Irash. Do not reuse empty container, Discard in trash.

#### HOW TO USE

Use 5 tbs /gal. water. Spray until all foliage is wet. Repeat at 5 day intervals to protect new foliage. Note reduced dosage for use on cucumbers and melons to avoid injury. Do not use teaspoons or tablespoons or other utensils for other purposes once used with pesticides. Note the time intervals to stop use before harvest. Subsequent planting of food and feed crops in treated soll must be made with crops on this or other carbaryl (Sevin) label(s).

		• • •	
۰ ب		•	-
*	÷ -	-	
	•	- -	· · ·
•		• -	-
*	-	-	
	_		
		•	· · · · · ·

		-		-			•		-	•••	-
٠		٠		٠		•	-		-		•
	٠	-		• •	-		. <del>.</del>		•		-
۳		٠	-	-		•	•	-	-		-
•		~	٠	-			-	•	۰.		•
	٠	٠		• •					-		•



## COMBINED FUNGICIDE AND INSECT SPRAY POWDER FOR CERTAIN VEGETABLES

ACTIVE INGREDIENTS:	
Zineb (Zinc ethylene	
bisdithiocarbamate)	38%
*Carbaryl (1-naphthyl-N-	
methylcarbamate)	25%
INERT INGREDIENTS	37%
TOTAL	100%
'SEVIN is the registered trademark of	Union

Carbide Corporation for carbaryl insecticide.



NET WEIGHT 6 OUNCES WHE COMMENTS EPA Letter Dated

# SEP17 1985

Winder the Federal Insecticide, Fingicide, and Rodenticide Act amended, for the resticide enclosed under EF. .eg. No. 73955-51

## CROPS

TOMATO, EGGPLANTS, PEPPERS Anthracnose, Septoria leaf spot, toma hopper. Start when plants are 6 inche before hervest. Wash to remove resi

CUCUMBERS, MELONS, SQUASH: water. Begin when plants start to run infestation. May be used up to 5 days bug, melon worms, cucumber beetles, naria blight and gummy stem blight, s use on summer squash closer than

BEANS: Mexican bean beetles, leafh stinkbugs, downy mildew, rust and / disease or insects. Do not use close

CABBAGE AND CAULIFLOWER: FI diamond back moth larvae, harlequ downy mildew; spray as soon as tran 10 days before harvest.

Limited Warranty, Manufacturer wan conforms to the ingredient statement g uct is suited for the labeled use when a Because of widely varying use condit risks even when label directions are f

Exclusion of Other Warranties and claimers and exclusions are specifical FOREGOING IS THE ONLY GUARA BLE TO THIS PRODUCT AND IS GIV ALL OTHER WARRANTIES EXPRES BILITY OR FITNESS FOR A PARTIC WARRANTIES WHICH EXCEED O WARRANTY ARE DISCLAIMED BY ufacturer's compliance with said limite NO REMEDY AGAINST MANUFAC AGE OR LOSS, and, IN NO EVE LIABLE FOR CONSEQUENTIAL DA

EPA REG. NO. 33955-511 EPA EST. NO. 2217-KS-1

