

UNDER THE FEDERAL INSECTICIDE  
 FUNGICIDE AND REPELLENT ACT  
 REGISTERED PRODUCT  
 TRADE NAME  
 SUBJECT

**ACME**

**PHALTAN 75W  
 FUNGICIDE**

**Controls certain diseases  
 of flowers, roses,  
 fruit and vegetables**

**ACTIVE INGREDIENT:**  
 Folpet [N-(Trichloromethylthio)phthalimide\*] ..... 75%  
**INERT INGREDIENTS:** ..... 25%  
 TOTAL 100%

U.S. Patent Nos. 2,553,770; 2,553,771  
 \*Phaltan is a registered trademark for phthalimide fungicide

**CAUTION: KEEP OUT OF REACH OF CHILDREN**  
 SEE LEFT SIDE PANEL FOR OTHER CAUTIONS

**NET WEIGHT 11 oz.**

**SOUR CHERRIES (RED TART):** Leaf spot and brown rot, use 2 to 4 teaspoonfuls per gallon. Apply as fruit buds separate, again just before bloom opens (popcorn stage), in full bloom and at 7-10 day intervals, with a final application after fruit is picked. Do not use on sweet cherries, may cause injury.

**CAUTION:** Avoid inhalation of dust or spray mist. Avoid prolonged or repeated contact with skin. Avoid contamination of feed or foodstuff. Do not store near herbicides. Do not reuse container. Destroy when empty.

**NOTICE:** Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

Made in U.S.A.  
**ACME INSECTICIDE DIVISION**  
**The Sherwin-Williams Co.**  
 8250 St. Aubin Ave., Detroit, Mich. 48211

USDA REG. NO. 577-517 LD-357B1 11-69 22954

**DIRECTIONS FOR USE**

May be mixed with most common insecticides and miticides. Do not combine with hydrated lime or other alkaline materials. Do not mix with or use in sprays immediately after oil sprays. Do not combine with emulsifiable concentrates, mixture may "curdle"! Shake container prior to measuring. Do not exceed dosages given. Use level teaspoonfuls.

**ASTERS, CHRYSANTHEMUMS, MARIGOLD, PHLOX, SNAPDRAGON AND ZINNIAS:** Alternaria leafspot, powdery mildew, Anthracnose, use 4 teaspoonfuls per gallon of water. Treat weekly during rapid new growth. Spray every 3 to 4 days during damp weather when disease is more severe.

**CARNATION, DIANTHUS:** Rust, Alternaria leafspot, use 4 teaspoonfuls per gallon about two weeks after planting. Repeat every 3 to 4 days for two weeks.

**ROSES:** Blackspot, Powdery Mildew, Rust, use 4 teaspoonfuls per gallon of water. Start as new foliage appears. Repeat at weekly intervals throughout the season. Spray as soon as possible following heavy rains. Do not combine with emulsifiable formulations, injury may result.

**CANTALOUPE, MELONS, CUCUMBERS AND SQUASH:** Downy mildew, Powdery mildew and Anthracnose, use 4 to 8 teaspoonfuls per gallon. Start as soon as first true leaves appear. Repeat weekly during growing season. Spray to wet all parts of leaves.

**CELERY:** spot, use as transp intervals. D

**LETTUCE** gallon. Ma are establ

**ONIONS** blotch, us soon as p tops ripen

**TOMATOE** gallon. Sta harvest.

**STRAWBE** spoonfuls Repeat we

**APPLES:** and black all fruit a buds are f vals until cious and first cover

**RASPBERR** mold, spu Spray to w Repeat we

# ACME

## PHALTAN 75W FUNGICIDE

**Controls certain diseases  
of flowers, roses,  
fruit and vegetables**

**ACTIVE INGREDIENT:**

Folpet [N-(Trichloromethylthio)phthalimide\*] ..... 75%

**INERT INGREDIENTS:** ..... 25%

TOTAL 100%

U.S. Patent Nos. 2,553,770; 2,553,771

\* Phaltan is a registered trademark for phthalimide fungicide

**CAUTION: KEEP OUT OF REACH OF CHILDREN**

SEE LEFT SIDE PANEL FOR OTHER CAUTIONS

**NET WEIGHT**

**11 oz.**

### DIRECTIONS FOR USE

May be mixed with most common insecticides and miticides. Do not combine with hydrated lime or other alkaline materials. Do not mix with or use in sprays immediately after oil sprays. Do not combine with emulsifiable concentrates, mixture may "curdle"! Shake container prior to measuring. Do not exceed dosages given. Use level teaspoonfuls.

**ASTERS, CHRYSANTHEMUMS, MARIGOLD, PHLOX, SNAPDRAGON AND ZINNIAS:** Alternaria leafspot, powdery mildew, Anthracnose, use 4 teaspoonfuls per gallon of water. Treat weekly during rapid new growth. Spray every 3 to 4 days during damp weather when disease is more severe.

**CARNATION, DIANTHUS:** Rust, Alternaria leafspot, use 4 teaspoonfuls per gallon about two weeks after planting. Repeat every 3 to 4 days for two weeks.

**ROSES:** Blackspot, Powdery Mildew, Rust, use 4 teaspoonfuls per gallon of water. Start as new foliage appears. Repeat at weekly intervals throughout the season. Spray as soon as possible following heavy rains. Do not combine with emulsifiable formulations, injury may result.

**CANTALOUPE, MELONS, CUCUMBERS AND SQUASH:** Downy mildew, Powdery mildew and Anthracnose, use 4 to 8 teaspoonfuls per gallon. Start as soon as first true leaves appear. Repeat weekly during growing season. Spray to wet all parts of leaves.

**CELERY:** Early blight, late blight, Cercospora leaf spot, use 4 teaspoonfuls per gallon. Start as soon as transplants are growing. Repeat at weekly intervals. Do not apply within 7 days of harvest.

**LETTUCE:** Downy mildew, use 4 teaspoonfuls per gallon. Make application weekly as soon as plants are established.

**ONIONS AND GARLIC:** Downy mildew, purple blotch, use 8 teaspoonfuls per gallon. Spray as soon as plants are well up. Repeat weekly until tops ripen.

**TOMATOES:** Anthracnose, use 4 teaspoonfuls per gallon. Start at first bloom, continue weekly until harvest.

**STRAWBERRIES:** Leaf spot, fruit rot, use 4 teaspoonfuls per gallon. Start spraying at first bloom. Repeat weekly until fruit is harvested.

**APPLES:** Scab, bitter rot, sooty blotch, fly speck and black rot, use 4 teaspoonfuls per gallon. Spray all fruit and foliage until wet. Start when flower buds are first apparent. Repeat at 10-14 day intervals until harvest. May cause russetting on Delicious and Stayman Winesap if used in petal fall and first cover sprays.

**RASPBERRIES, BLACKBERRIES:** Fruit rot, gray mold, spur blight, use 4 teaspoonfuls per gallon. Spray to wet all foliage. Start when first fruit set. Repeat weekly until harvest.

9 22954