



[33955-519



ACME



**RED SPIDER AND
MITE SPRAY**

CONTAINS KELTHANE*

**Controls Clover Mites, Red Spiders on
Roses, Evergreens, Trees and Flowers**

ACTIVE INGREDIENT:

*1-bis(chlorophenyl)
2,2,2-trichloroethanol

INERT INGREDIENTS:

16.5%
83.5%
TOTAL 100.0%

*Kelthane Registered TM Robin and Haas Co.

**CAUTION. KEEP OUT OF REACH OF CHILDREN
SEE LEFT PANEL FOR OTHER CAUTIONS**

ONE PINT

U. S. MEASURE

DIRECTIONS FOR USE

Read cautions and directions carefully. Use 2 teaspoonfuls per gallon. Apply at first sign of mites. Spray all leaf surfaces. Repeat in 10 days or as necessary to maintain control. Use a 5X magnifying glass to examine mottled or yellowish leaves.

CLOVER MITES: Spray in early Spring around windows, walls and foundations. Also spray shrubs and edge of lawn. Plants such as ivy should be thoroughly sprayed.

ROSES, EVERGREENS, ORNAMENTALS: On tender foliage reduce dosage to 1 teaspoonful per gallon and repeat in 5 to 7 days. Greenhouse plants are more susceptible to injury than those outdoors.

STRAWBERRIES: Several mites attack this plant. For Cyclamen mites use 4 teaspoonfuls per gallon. Use as high pressure as possible to force spray into growing buds. Spray as soon as growth starts. Repeat 2 or 3 times. Do not spray closer than 7 days before picking.

VEGETABLES: For two spotted mites and tomato russet mites spray beans, melons, cucumbers, tomatoes and peppers. Do not use on eggplants. Stop spraying beans 7 days before picking. 7 days before harvest on other crops.

GRAPES, RASPBERRIES: Apply just after bloom or when mites appear. Repeat as necessary. Do not use on grapes within 7 days of harvest or on raspberries within 7 days.

APPLES, PEARS, PEACHES, PLUMS: Apply at petal fall through second cover spray to control European red and clover mites and prevent buildup of two spotted mite. Repeat during season, if necessary, but no closer than 30 days since the last application. Stop spraying peaches and apricots 14 days before harvest, apples, pears, plums 7 days before harvest.

CAUTION: Harmful if swallowed. Avoid inhalation of spray, mist, skin contact and contamination of feed or foodstuff. Wash with soap and water after spraying or other contact.

This product toxic to fish. Keep out of any body of water. Destroy container when empty. Do not reuse for any purpose.

NOTICE: Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

ACME INSECTICIDE DIVISION

The Sherwin-Williams Co.

8250 St. Aubin Ave., Detroit, Mich. 48211

Made in U.S.A.

U.S. PAT. NO. 2,811,111

LD 3547