

# acme methoxychlor 50% wettable

## DIRECTIONS FOR USE:

### Tree Fruit:

#### Apples and Pears

To control codling moth, plum curculio, apple maggot, tent caterpillar and Japanese beetle. Use the stronger dosage where curculio is a problem.

#### Apricots, Peaches, Nectarines, Plums

To control plum curculio and Japanese beetle, use 8 tsp./gallon. Start at petal fall or when adults appear.

#### Cherries

To control cherry fruitworm, cherry fruit fly and rose chafer, use 5 tsp./gallon. Use 8 tsp./gallon for plum curculio and Japanese beetle.

### Bush Fruit:

#### Blackberries, loganberries, raspberries

To control rose chafer and strawberry weevil, use 8 tsp./gallon on foliage when insects appear.

#### Strawberries

To control leaf tier, flea beetle, spittlebug, strawberry leaf beetle and strawberry weevil, use 8 tsp./gallon. Apply on foliage as needed.

#### Grapes

To control grape berry moth, grape leafhopper and rose chafer, use 5 tsp./gallon. Start at petal fall and repeat. Japanese beetle, use 8 tsp./gallon and apply as soon as insects appear.

### Vegetable Gardens:

#### Asparagus

To control asparagus beetle, use 5 tsp./gallon. Wash to remove residues.

#### Beans, blackeyed peas

To control corn earworm, flea beetle, Mexican bean beetle, potato leafhopper, use 8 tsp./gallon. Do not feed vines to animals.

#### Beets

To control blister beetle and flea beetle, use 8 tsp./gallon. Spray foliage.

#### Cucumbers, squash, melons

To control cucumber beetle, use 8 tsp./gallon on foliage as soon as insects appear.

If dosage reduced to 1/2 in last two sprays.

#### Eggplant, peppers

To control flea beetle, use 8 tsp./gallon on foliage as needed.

#### Potatoes

To control leafhoppers, use 10 tsp./gallon on foliage as needed.

#### Tomatoes

To control blister beetle, flea beetle and tomato fruitworm, use 8 tsp./gallon. Apply on foliage as needed.

When disease is a problem, combine with Acme Zineb 75W Fungicide or Acme Wettable Dusting Sulphur.

## Time to Stop Use Before Harvest

7 days

21 days

7 days

3 days

3 days

14 days

3 days

3 days

7 days (Roots),  
14 days (Tops)

7 days

1 day

1 day

No Limit

7 days

## DIRECTIONS FOR USE:

With the lid on, shake the can prior to measuring. Apply at 1 to 1 1/2 lbs. per 50 gallons of water — 5 to 8 teaspoonfuls per gallon. Apply so as to completely wet all fruit and foliage. Do not apply more than 500 gallons per acre — 5 lbs. actual Methoxychlor. Make application as soon as insects appear, at petal fall on fruit. Repeat at 7-14 day intervals to maintain control. Do not apply closer to harvest than shown below for specific crops.

To control house flies, stable flies, and aid in residual control of mosquitoes, use the contents of this can in 5 quarts of water (5% actual). Apply 1 to 2 gallons per 1000 sq. ft. or to run-off point. Spray around porches, windows and outside walls of farm buildings where insects tend to congregate. Do not use this dosage on crops. Around barns, remove all animals and cover feed and water supplies before spraying.

**CAUTION:** Wash thoroughly after handling. Avoid breathing dust while measuring. Do not feed treated crop residues to animals if sprayed within 7 days of harvest. Avoid contaminating feed and foodstuffs.

### Environmental Cautions:

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this product.

This product may kill fish and other aquatic organisms. Avoid spraying or spillage that will contaminate any body of water. Apply only as specified on this label.

This product is toxic to bees and should not be applied when bees are actively visiting the area.

**NOTICE:** Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

**SOLD FOR INSECTICIDE PURPOSES.**


**DO NOT REUSE CONTAINER. DESTROY WHEN EMPTY.**

Made in U.S.A. for:

**The Sherwin-Williams Co.** CLEVELAND, OHIO 44101

LD366 C1 10-73

EPA No. 577-528-AA



33955  
528

to Start or  
before harvest

7 days

14 days

7 days

3 days

3 days

14 days

3 days

3 days

ays (Roots)  
ays (Top)

7 days

1 day

1 day

No limit

7 days

**DIRECTIONS FOR USE**

Shake the lid on, shake the can prior to measurement. Apply at 1 to 1 1/2 lbs. per 50 gallons of water (5 to 8 teaspoons per gallon). Apply so as to completely wet all fruit and foliage. Do not apply more than 500 gallons per acre. 5 lbs. actual Methoxychlor. Make application as soon as insects appear at petal fall on fruit. Repeat at 7-14 day intervals to maintain control. Do not apply closer to harvest than shown below for specific crops.

To control house flies, stable flies, and aid in residual control of mosquitoes, use the contents of this can in 5 quarts of water (5 lbs. actual). Apply 1 to 2 gallons per 1000 sq. ft. or to run-off point. Spray around porches, windows and outside walls of farm buildings where insects tend to congregate. Do not use this dosage on crops. Around barns, remove all animals and cover feed and water supplies before spraying.

**CAUTION** Wash thoroughly after handling. Avoid breathing dusts. Do not feed treated crop residues to animals. Do not spray within 7 days of harvest. Avoid contaminating feed and feedstuffs.

**Environmental Cautions:**

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water by cleaning equipment or disposing of wastes. Apply to outdoor plants as specified on this product.

To spray trees, shrubs and other aquatic organisms. Avoid spraying or spraying that will contaminate any body of water. Apply as specified on this label.


This product is toxic to birds and other animals. Do not feed to animals. Do not allow access to treated areas.

Do not use for purposes that the product is not intended for. Do not use for purposes not specifically stated on the label. Do not use for purposes not specifically stated on the label, when the label is clearly marked with directions under normal conditions of use. No warranty, express or implied, extends to the use of this product for any purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or for purposes not intended, or for purposes not intended, or for purposes not intended, or for purposes not intended, or for purposes not intended.

**USE FOR INSECTICIDE PURPOSES  
DO NOT REUSE CONTAINER. DESTROY WHEN EMPTY**

Made in U.S.A. for  
**The Sherwin-Williams Co.** CLEVELAND, OHIO 44101

© 1966 Sherwin-Williams Co.



**\*For Control of Certain Flies, Mosquitoes and Certain Chewing Insects on Shade Trees, Fruit Trees and Vegetables.**

**Active Ingredient:**  
Methoxychlor\*\*, Technical..... 50.0%  
**Inert Ingredients:**..... 50.0%  
TOTAL..... 100.0%

\*\*Equivalent to 44% 2,2-bis(p-methoxyphenyl)-1,1,1-trichloroethane and 6% other isomers and reaction products.

**CAUTION: Keep Out Of Reach Of Children.  
Read caution elsewhere on label.**


**ONE POUND NET WEIGHT**

33955-527

# acme pestroy 25% methoxychlor

DEFINITION

*[Faint, illegible text, likely a definition or description of the product.]*



Control of many of the most serious  
pests of the home and garden.

Keep out of reach of children.  
Use only as directed.