## DIRECTIONS FOR USE STOP! READ THE LABEL BEFORE USE!

Shake well before measuring. Keep container closed to prevent caking of contents. Best results are obtained by thinning the measured dosage with water in a separate container before adding to the sprayer. Protect from freezing as bottle may break. Treat at first sign of insect damage and repeat in 7 to 10-days intervals, or as necessary to maintain control. Spray all plant surfaces thoroughly.

STRAWBERRIES: Use 4 tsp./gal. water up to 1 day before picking to control leaf rollers, strawberry weevil, and meadow spittle bug.

ROSES: Use 4 tsp./gal. water to control Exposed thrips. Rose slug, Japanese beetle, Rose aphids and leaf rollers.

riowers & shrubs: Use 4 tsp./gal. water to control Japanese beetle, boxelder bust tent caterpillars, lace bugs, leaf rollers, hoppers.

bagworms, Elm leaf beetles, birchleaf miners, boxelder bugs, tent caterpillars, periodical cicadas, and leafhoppers.

LAWNS: Mow lawn and water before spraying. Some use 1 pint in 15 gals. water, and apply to 250 sq. ft. of turf to control chinch bugs sod webworms (lawn moths), fall armyworms, comparison, earwigs, fleas, and ants.



An insecticide for control of certain insect pests of shade trees, lawns, ornamentals, fruit trees & vegetables.

his product contains 2 lbs. actual Carbaryl per gallon.

CAUTION: Keep Out of Reach of Children

See Side Panels for Additional Cautions.

Trade Mark UNION CARBIDE CORPORATION for 1-naphthylimethylcarbamate. U.S. Patent Numbers 2,903,478 & 3,009,855.

**NET CONTENTS ONE QUART** 

VEGETABLES: Lima and green beans—use 4 tsp./gal. to control Mexican bean beetle, leaf hoppers, flea beetles, and tarnished plant bugs. May be used up to 1 day before harvest. Do not feed vines to dairy or meat animals. Tomato eggplant, and peppers—use 4 tsp./gal. to control tomato fruitworms, stinkbugs, flea beetles, leaf-hoppers, and tall armyworms. Apply up to day of picking. Cuclimbers, melons, and squash—use 4 tsp./gal. to control squash bugs, melon worms, cucumber beetles, and flea beetles. Do not apply to wet foliage or to watermelons in Florida or injury may result. Potatoes—use 4 tsp./gal. to control Colorado potato beetle leaf-hoppers, and flea beetles.

APPLES & PEARS: To control codling moth, plum curculio, apple maggot, and full tree leat roller, use 4 tsp./gal. Do not apply before second cover spray to avoid fruit thinning. May be used up to 1 day before picking.

PEACHES & APRICOTS: To control oriental fruit moth, plum curculio, and cat facing bugs use 4 tsp./gal. May be used up to 1 day before harvest on peaches, 3 days before harvest on apricots.

GRAPES: To control grape berry moth leafhoppers, and red-banded leaf rollers use 6 tsp./gal. May be used up to day of harvest.

ADULT MOSQUITOES: Spray, shrubbery, hedges flower beds, and other places where mosquitoes may rest during the day, using 4 tsp./gal. of spray.

CAUTION: Harmful if swallowed or inhaled. Avoid breathing of spray mist. Avoid skin contact. Wash face and hands after spraying. Avoid contamination of food and feedstuffs. Keep out of reach of children and pets. Do not use in any manner not specified in this label.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal.

environmental cautions: Keep out of lakes, streams, and ponds. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.

This product is highly toxic to bees exposed to cirect treatment or residues on plants. Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not spray when plants are in full bloom or when bees are active in the area.

DO NOT RELISE CONTAINER. DESTROY WHEN EMPTY.

EPA REG. NO. 33955-533-AA

EPA EST. NO. 2217-KS-1

838-275

pbi/gordon
corporation
KANBAR CITY, KANBAR 66118

#C-2-17