

PM 19

33955-533

PA 175

ACME® SEVIN® BRAND CARBARYL INSECTICIDE LIQUID DURA-SPRAY®

An insecticide for the control of certain pests affecting vegetables,
fruit, citrus fruit, nuts, shade trees, fruit trees, lawn and ornamentals

- KEEP FROM FREEZING -

ACTIVE INGREDIENT:

Carbaryl (1-naphthyl N-methylcarbamate).....	23.4%
INERT INGREDIENTS.....	76.6%
TOTAL	100.0%

This Product Contains 2 lbs. carbaryl per gallon.


SEVIN® is the registered trademark of Rhône-Poulenc for carbaryl insecticide.

KEEP OUT OF REACH OF CHILDREN

CAUTION

See back panel for Precautionary Statements and Statement of Practical Treatment

NET CONTENTS ONE-HALF GALLON



838/ APXXXXXX

EPA Reg. No. 33955-533

EPA Est. No. 2217-KS-1

Manufactured By ACME DIVISION

PBI/GORDON CORPORATION

KANSAS CITY, MISSOURI 64101

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: May be Harmful if swallowed. Avoid breathing vapors (dust or spray mist). Avoid contact with eyes, skin or clothing. Wash thoroughly after handling.

- User Safety Requirements:**
- ◆ Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when mixing and applying this product.
 - ◆ Wear a hat and eye protection when making overhead applications.
 - ◆ Remove clothing immediately if pesticide soaks clothing. Change clothing as soon as possible after use.
 - ◆ Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.
 - ◆ Do not allow children or pets to contact treated area until sprays have dried.

Statement of Practical Treatment

IF SWALLOWED: Induce vomiting and seek medical attention immediately.

IF IN EYES OR ON SKIN: Flush eyes thoroughly with water, wash skin thoroughly with soap and water.

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. If person is unconscious, do not give anything by mouth and do not induce vomiting.

IF IN EYES: Flush eyes with plenty of water. Call a physician if irritation persists.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

NOTE TO PHYSICIAN: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine sulfate is antidotal. Do not use 2-PAM, opiates, or cholinesterase inhibiting drugs.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to aquatic and estuarine invertebrates. For terrestrial uses, Do not apply directly to water, or, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwaters. Do not apply when weather conditions favor drift away from target area.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or ornamental gardens or parks, or on golf courses or lawns and grounds.

STORAGE & DISPOSAL

STORAGE: Keep from freezing. Store in original container in a cool, dry and locked storage area inaccessible to children or pets. Do not store with food or feed. Do not store in areas where temperatures frequently exceed 100°F. Keep from freezing as bottle may break.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse container. Rinse thoroughly before discarding in trash.

- GENERAL -

- ◆ Shake well before measuring. Keep container closed to prevent caking of contents. Best results are obtained by thinning the measured dosage with water in a separate container before adding to the sprayer.
- ◆ Treat at first sign of insect damage and repeat in 7 to 10 day intervals, or as necessary to maintain control. **Spray all plant surfaces thoroughly.**
- ◆ Apply when air is calm to avoid drift and contact with eyes and skin.
- ◆ Start applying at the farthest corner of the treatment area and work backward to avoid contact with wet surfaces. Allow spray to dry in treated areas before reentering.
- ◆ For trees taller than 10 feet, consider hiring a licensed professional.
- ◆ Spray thoroughly to wet upper and lower leaf surfaces, stems and branches.
- ◆ Do not repeat applications more than once a week.
- ◆ Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl label(s).

- VEGETABLES -

-You may use up to the day of harvest on:

Beans (Green or Lima), Carrots, Corn, Eggplant, Peas, Peppers, Potatoes & Tomatoes: Mix 4 to 6 teaspoons of product per gallon of water to control bean leaf beetle, flea beetle, Japanese beetle, leafhoppers, thrips, Western cutworm, armyworm, chinch bug, corn earworm, corn rootworm adults, European corn borer, fall armyworm, Colorado potato beetle, tomato fruitworm, tomato hornworm, tomato pinworm, tarnished plant bug, stink bugs. To control corn earworm, apply to silks as soon as they appear. Make 3 to 4 applications at 2 to 3 day intervals as infestation warrants. Application of this product to the tassel region of corn during the pollen shed period will seriously reduce bee population. **Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.**

Cucumbers, Melons, Pumpkin & Squash (Summer and Winter): Mix 4 teaspoons of product per gallon of water to control pickleworm, melonworm, cucumber beetle, flea beetle, leafhoppers, squash bugs. **Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.**

-You may use up to 3 days before harvest on:

Beets (Roots only), Broccoli, Brussels Sprouts, Cabbage, Cauliflower, Lettuce (Head only), Parsnips, Radish and Rutabagas: Mix 4 to 6 teaspoons of product per gallon of water to control flea beetle, harlequin bug, leafhoppers, lygus bug, and spittlebugs, armyworm, corn earworm, fall armyworm, imported cabbageworm, stink bugs, and tarnished plant bug. **Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.**

NOTE: If beet tops are to be used for food, and, if used on Leaf Lettuce, do not use within 14 days of harvest.

-You may use up to 14 days before harvest on:

Beets (Tops only), Endives, Lettuce (Leaf only), Mustard Greens and Spinach: Mix 4 to 6 teaspoons of product per gallon of water to control armyworm, flea beetle, harlequin bug, leafhoppers, imported cabbageworm, lygus bug, and stink bugs. **Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.**

4/5

- FRUIT -

-You may use up to the day of harvest on:

Grapes: Mix 4 to 8 teaspoons of product per gallon of water to control European fruit lecanium, grape leafroller, grape leafhopper, Western grapeleaf skeletonizer, Western yellow-striped armyworm, cutworm, grape berry moth, red-banded leafroller. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.

-You may use up to 1 day before harvest on:

Apples, Cherries, Peaches, Pears, and Plums: Mix 4 teaspoons of product per gallon of water to control apple aphid, apple rust mite, apple mealybug, apple maggot, forbes scale, green fruitworm, black cherry aphid, brown soft scale, cherry fruitworm, European earwig, Japanese beetle, lecanium scales, pear psylla, pear leaf blister mite, pear rust mite, periodical cicada, plum curculio, redbanded leafroller, San Jose scale, tarnished plant bug, woolly apple aphid, codling moth, fruitree leafroller, lesser peachtree borer, peach twig borer, tussock moth, mealy plum aphid, prune leafhopper, variegated leafroller. For trees taller than 10 feet, consider hiring a licensed professional. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.

Blueberries, Cranberries & Strawberries: Mix 8 teaspoons of product per gallon of water on strawberries and 4 teaspoons of product per gallon on blueberries and cranberries. Uniformly spray each gallon of finished spray over 160 square feet to control cutworm, blueberry maggot, cherry fruitworm, cranberry fruitworm, cranberry fireworm, Japanese beetle, leafhoppers, European fruit lecanium, flea beetle, meadow spittlebug, strawberry leafroller, strawberry weevil. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.

-You may use up to 3 days before harvest on:

Apricots and Nectarines: Refer to apples, cherries, peaches, pears, and plums shown above. For trees taller than 10 feet, consider hiring a licensed professional. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.

-You may use up to 5 days before harvest on:

Citrus (Grapefruit, Lemons, Limes, Oranges, Tangerines): Mix 4 teaspoons of product per gallon of water to control avocado leafroller, California orangedog, citrus cutworm, citrus root weevil, fruitree leafroller, orange tortrix. Do not apply less than 10 gallons dilute spray per mature tree. May be mixed with petroleum oils commonly used on citrus. For trees taller than 10 feet, consider hiring a licensed professional. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.

- NUTS -

-You may use up to the day of harvest on:

Filberts, Pecans & Walnuts: Mix 4 teaspoons of product per gallon of water to control filbert aphid, filbert leafroller, filbertworm, pecan leaf phylloxera, pecan nut casebearer, pecan spittlebug, pecan weevil, walnut caterpillar, codling moth, calico scale, European fruit lecanium. For trees taller than 10 feet, consider hiring a licensed professional. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.

575

- ORNAMENTALS -

Roses, Flowers, Shrubs and Shade Trees: Do not use on Boston Ivy. Mix 4 teaspoons of product per gallon of water to control ants, armyworm, bagworm, boxelder bugs, Japanese beetle, blister beetle, Elm leaf beetle, flea beetle, willow leaf beetle, leafhoppers, tent caterpillar, June beetle, gypsy moth, leafrollers, birch leafminer, mealybugs, exposed thrips, stink bugs, oak leafminer, periodical cicada, and ticks. Apply dilute sprays to obtain thorough coverage of both upper and lower leaf surfaces. To control scale insects, treat trunks, stems and twigs in addition to plant foliage. The use of spreader-sticker may improve residual control. For trees taller than 10 feet, consider hiring a licensed professional. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.

Lawns and Ornamental Turf: For best results, mow lawn and water before spraying. Mix 1 pint of product in 15 gallons of water and spray over 1,250 square feet to control ants, armyworm, bluegrass billbug, centipedes, chiggers, cutworm, earwigs, fall armyworm, fleas, grasshoppers, June beetle, leafhoppers, millipedes, mosquitoes, sod webworm (lawn moth), sowbugs, springtails, ticks, chinch bugs.

LIMITED WARRANTY AND DISCLAIMER. The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.