

33955-533

9-10-2004

Please read instructions on reverse before completing form.

Form Approved. OMB No. 2070-0060. Approval Expires 2-28-95



United States
Environmental Protection Agency
Washington, DC 20460

- Registration
- Amendment
- Other

OPP Identifier Number

Application for Pesticide - Section I

1. Company/Product Number 33955-533	2. EPA Product Manager Meredith Laws, Chief	3. Proposed Classification <input checked="" type="checkbox"/> None <input type="checkbox"/> Restricted
4. Company/Product (Name) Acme Sevin Brand Carbaryl Insecticide Liquid Dura-Spray	PM# 57 Insecticide-Rodenticide Branch	
5. Name and Address of Applicant (Include ZIP Code) PBI/Gordon Corporation Post Office Box 014090 Kansas City, Missouri 64101 Check if this is a new address <input type="checkbox"/>	6. Expedited Review. In accordance with FIFRA Section 3(c)(3) (b)(i), my product is similar or identical in composition and labeling to: EPA Reg. No. _____ Product Name _____	

Section - II

<input type="checkbox"/> Amendment - Explain below.	<input type="checkbox"/> Final printed labels in response to Agency letter dated _____
<input type="checkbox"/> Resubmission in response to Agency letter dated _____	<input type="checkbox"/> "Me Too" Application.
<input checked="" type="checkbox"/> Notification - Explain below.	<input type="checkbox"/> Other - Explain below.

NOTIFICATION

SEP 10 2004

Explanation: Use additional page(s) if necessary. (For section I and Section II.)

Notification per Pesticide Registration Notice (PRN) 98-10:

We ask to add the following ad claim: "Kills Over 100 Insects on Listed Vegetables, Fruits, Ornamentals and Lawns".

This notification is consistent with the provisions of PR Notice 98-10 and EPA regulations at 40 CFR 152.46, and no other changes have been made to the labeling or Confidential Statement of Formula of this product. I understand that it is a violation of 18 U.S.C Sec. 1001 to willfully make any false statement to the EPA. I further understand that if this notification is not consistent with the terms of PR Notice 98-10 and 40 CFR 152.46, this product may be in violation of FIFRA and I may be subject to enforcement action and penalties under Section 12 and 14 of FIFRA.

Section - III

1. Material This Product Will Be Packaged In:					
Child-Resistant Packaging <input type="checkbox"/> Yes* <input type="checkbox"/> No	Unit Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	Water Soluble Packaging <input type="checkbox"/> Yes <input type="checkbox"/> No	2. Type of Container <input type="checkbox"/> Metal <input type="checkbox"/> Plastic <input type="checkbox"/> Glass <input type="checkbox"/> Paper <input type="checkbox"/> Other (Specify) _____		
* Certification must be submitted		If "Yes" Unit Packaging wgt. No. per container	If "Yes" Package wgt. No. per container		
3. Location of Net Contents Information <input type="checkbox"/> Label <input type="checkbox"/> Container		4. Size(s) Retail Container		5. Location of Label Directions <input type="checkbox"/> On Label <input type="checkbox"/> On Labeling accompanying product	
6. Manner in Which Label is Affixed to Product <input type="checkbox"/> Lithograph <input type="checkbox"/> Other _____ <input type="checkbox"/> Paper glued <input type="checkbox"/> Stenciled					

Section - IV

1. Contact Point (Complete items directly below for identification of individual to be contacted, if necessary, to process this application.)					
Name James L. Kunstman, Ph.D.	Title Director of Regulatory Services	Telephone No. (Include Area Code) 816-460-6292			
Certification I certify that the statements I have made on this form and all attachments thereto are true, accurate and complete. I acknowledge that any knowingly false or misleading statement may be punishable by fine or imprisonment or both under applicable law.					6. Date Application Received (Stamped)
2. Signature 	3. Title Director of Regulatory Services				
4. Typed Name James L. Kunstman, Ph.D.	5. Date August 23, 2004				

ACME SEVIN® BRAND CARBARYL INSECTICIDE LIQUID DURA-SPRAY®

FOR RESIDENTIAL USE ONLY

AN INSECTICIDE FOR THE CONTROL OF LISTED PESTS AFFECTING
VEGETABLES, FRUIT, CITRUS FRUIT, NUTS, SHADE TREES, FRUIT
TREES, LAWN AND ORNAMENTALS.

- KEEP FROM FREEZING -

ACTIVE INGREDIENT:	
Carbaryl (1-naphthyl N-methylcarbamate)	23.4%
INERT INGREDIENTS:	76.6%
TOTAL	100.0%

This product contains 2 lbs. carbaryl per gallon.

KEEP OUT OF REACH OF CHILDREN


CAUTION

See back panel for Precautionary Statements and First Aid.

NET CONTENTS: ONE-HALF GALLON, ONE QUART, ONE PINT

838/ AP082304
 EPA REG. NO. 33955-533
 EPA Est. No. 2217-KS-1
MANUFACTURED BY:

An Employee-Owned Company
 1217 West 12th Street
 Kansas City, Missouri 64101
 Telephone: 1-800-821-7925



3/11



READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION: Harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with eyes, skin or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

User Safety Requirements:	
◆	Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when mixing and applying this product.
◆	Wear a hat and eye protection when making overhead applications.
◆	Remove clothing immediately if pesticide soaks clothing. Change clothing as soon as possible after use.
◆	Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.
◆	Do not allow children or pets to contact treated area until sprays have dried.

First Aid

If swallowed:	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give anything by mouth to an unconscious person.
If in eyes:	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. • Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 - 20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice.
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact 1-877-800-5556 for emergency medical treatment advice.</p> <p>NOTE TO PHYSICIAN: Carbaryl is a N-methyl carbamate insecticide which is a cholinesterase inhibitor. Specific antidote consists of atropine. Narcotics and other sedatives should not be used. Drugs like 2-PAM (pyridine-2-aldoxime methiodide) are NOT recommended.</p>	

ENVIRONMENTAL HAZARDS:

This pesticide is toxic to aquatic and estuarine invertebrates. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater. Do not apply when weather conditions favor drift away from target area.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes or ornamental gardens or parks, or on golf courses or lawns and grounds.

4/11

STORAGE & DISPOSAL

STORAGE: Keep from freezing. Store in original container in a cool, dry and locked storage area inaccessible to children or pets. Do not store with food or feed. Do not store in areas where temperatures frequently exceed 100°F. Keep from freezing as bottle may break.

CONTAINER DISPOSAL:

If empty: Do not reuse this container. Place in trash or offer for recycling if available.

If partly filled: Call your local solid waste agency or 1-800-CLEANUP for disposal instructions. Never place unused product down any indoor or outdoor drain.

INSECT CONTROL:

Apply when insects or their damage appears. To maintain control repeat at 7 to 10 day intervals unless a shorter interval is specified below. Where a dosage range is indicated use the lower rate on young plants and early instars, and use the higher rate on mature plants, advanced instars, and adults. Thorough and uniform spray coverage is essential for effective control.

- ◆ Shake well before measuring. Keep container closed to prevent caking of contents. Best results are obtained by diluting measured amount of product with water in a separate container before adding to the sprayer.
- ◆ Apply when air is calm to avoid drift and contact with eyes and skin.
- ◆ Start applying at the farthest corner of the treatment area and work backward to avoid contact with wet surfaces. Allow spray to dry in treated areas before reentering.
- ◆ For trees taller than 10 feet, consider hiring a licensed professional.
- ◆ Spray thoroughly to wet upper and lower leaf surfaces, stems and branches.
- ◆ Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl label(s).

- INSECT CONTROL -

Vegetables, Berries, Citrus Fruit, Stone Fruits, Pome Fruits & Nuts	Insects	Maximum Application Rate, Tsp. of Product/ 1 Gal. of Water	Retreatment Interval, Days	Preharvest Interval, Days	Comments
Beans (green & lima); Peas	bean leaf beetle, flea beetle, Japanese beetle, leafhoppers, thrips, Western cutworm, armyworm, chinch bug, corn earworm, corn rootworm (adults), European corn borer, fall armyworm, Colorado potato beetle, tomato fruitworm, tomato hornworm, tomato pinworm, tarnished plant bug, and stink bug.	4 - 6 tsp./gallon	7 days	PHI = 3 days (succulents) PHI = 21 days (dried) PHI = 14 days (forage) PHI = 21 days (hay)	Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.
Carrots; Beets (roots only); Parsnip; Radish; Rutabaga	flea beetle, harlequin bug, leafhoppers, lygus bug, spittlebug, armyworm, corn earworm, fall armyworm, imported cabbageworm, tarnished plant bug, and stink bug.	4 - 6 tsp./gallon	7 days	PHI = 7 days	Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.
Brussel sprouts; Broccoli; Cabbage; Cauliflower	flea beetle, harlequin bug, leafhoppers, lygus bug, spittlebug, armyworm, corn earworm, fall armyworm, imported cabbageworm, tarnished plant bug, and stink bug.	4 - 6 tsp./gallon	7 days	PHI = 3 days	Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.
Sweet Corn Only	bean leaf beetle, flea beetle, Japanese beetle, leafhoppers, thrips, Western cutworm, armyworm, chinch bug, corn earworm, corn rootworm (adults), European corn borer, fall armyworm, Colorado potato beetle, tomato fruitworm, tomato hornworm, tomato pinworm, tarnished plant bug, and stink bug.	4 - 6 tsp./gallon	3 days	PHI = 2 days (ears) PHI = 14 days (forage) PHI = 48 days (fodder)	To control corn earworm, apply to silks when earworms appear. Make 3 to 4 applications at 3 day intervals as infestation warrants. Application of this product to the corn tassel during pollination will seriously reduce bee population. Spray thoroughly to wet upper and lower leaf surfaces, and stems.

11/5

Continued...

Vegetables, Berries, Citrus Fruit, Stone Fruits, Pome Fruits & Nuts	Insects	Maximum Application Rate, Tsp. of Product/ 1 Gal. of Water	Retreatment Interval, Days	Preharvest Interval, Days	Comments
Cucumber; Melon; Pumpkin; Squash (summer & winter)	pickleworm, melonworm, cucumber beetle, flea beetle, leafhoppers, and squash bug.	4 - 6 tsp./gallon	7 days	PHI = 3 days	Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.
Potato	bean leaf beetle, flea beetle, Japanese beetle, leafhoppers, thrips, Western cutworm, armyworm, chinch bug, corn earworm, corn rootworm (adults), European corn borer, fall armyworm, Colorado potato beetle, tomato fruitworm, tomato hornworm, tomato pinworm, tarnished plant bug, and stink bug.	4 - 6 tsp./gallon	7 days	PHI = 7 days	Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.
Tomato; Eggplant; Pepper	bean leaf beetle, flea beetle, Japanese beetle, leafhoppers, thrips, Western cutworm, armyworm, chinch bug, corn earworm, corn rootworm (adults), European corn borer, fall armyworm, Colorado potato beetle, tomato fruitworm, tomato hornworm, tomato pinworm, tarnished plant bug, and stink bug.	4 - 6 tsp./gallon	7 days	PHI = 3 days	Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.
Lettuce (head & leaf); Endive; Spinach; Garden Beets (tops only);	flea beetle, harlequin bug, leafhoppers, lygus bug, spittlebug, armyworm, corn earworm, fall armyworm, imported cabbageworm, tarnished plant bug, and stink bug.	4 - 6 tsp./gallon	7 days	PHI = 14 to 15 days	Spray thoroughly to wet upper and lower leaf surfaces, and stems. Do not repeat applications more than once a week.

6/11

Continued....

Vegetables, Berries, Citrus Fruit, Stone Fruits, Pome Fruits & Nuts	Insects	Maximum Application Rate, Tsp. of Product/ 1 Gal. of Water	Retreatment Interval, Days	Preharvest Interval, Days	Comments
Mustard Greens	flea beetle, harlequin bug, leafhoppers, lygus bug, spittlebug, armyworm, corn earworm, fall armyworm, imported cabbageworm, tarnished plant bug, and stink bug.	4 - 6 tsp./gallon	7 days	PHI = 13 to 14 days	Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.
Grape	European fruit lecanium, grape leafroller, grape leafhopper, Western grapeleaf skeletonizer, Western yellow-striped armyworm, cutworm, grape berry moth, and redbanded leafroller.	4 - 8 tsp./gallon	7 days	PHI = 7 days	Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.
Cherry; Peach; Plum; Apricot; Nectarine;	apple aphid, apple rust mite, apple mealybug, apple maggot, forbes scale, green fruitworm, black cherry aphid, brown soft scale, cherry fruitworm, European earwig, Japanese beetle, lecanium scales, pear psylla, pear leaf blister mite, pear rust mite, periodical cicada, plum curculio, redbanded leafroller, San Jose scale, tarnished plant bug, woolly apple aphid, codling moth, fruitree leafroller, lesser peachtree borer, peach twig borer, tussock moth, mealy plum aphid, prune leafhopper, and variegated leafroller.	4 tsp./gallon	7 days	PHI = 3 days	For trees taller than 10 feet, consider hiring a licensed professional. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.

Handwritten mark

Continued....

Vegetables, Berries, Citrus Fruit, Stone Fruits, Pome Fruits & Nuts	Insects	Maximum Application Rate, Tsp. of Product/ 1 Gal. of Water	Retreatment Interval, Days	Preharvest Interval, Days	Comments
Apple; Pear;	apple aphid, apple rust mite, apple mealybug, apple maggot, forbes scale, green fruitworm, black cherry aphid, brown soft scale, cherry fruitworm, European earwig, Japanese beetle, lecanium scales, pear psylla, pear leaf blister mite, pear rust mite, periodical cicada, plum curculio, redbanded leafroller, San Jose scale, tarnished plant bug, woolly apple aphid, codling moth, fruitree leafroller, lesser peachtree borer, peach twig borer, tussock moth, mealy plum aphid, prune leafhopper, and variegated leafroller.	4 tsp./gallon	7 days	PHI = 3 days	For trees taller than 10 feet, consider hiring a licensed professional. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week. To avoid undesired apple thinning, delay use until at least 30 days after full bloom.
Blueberry; Cranberry	cutworm, blueberry maggot, cherry fruitworm, cranberry fruitworm, cranberry fireworm, Japanese beetle, leafhoppers, European fruit lecanium, flea beetle, meadow spittlebug, strawberry leafroller, and strawberry weevil.	6 tsp./gallon	7 days	PHI = 7 days	Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.
Strawberry	cutworm, blueberry maggot, cherry fruitworm, cranberry fruitworm, cranberry fireworm, Japanese beetle, leafhoppers, European fruit lecanium, flea beetle, meadow spittlebug, strawberry leafroller, and strawberry weevil.	6 tsp./gallon	7 days	PHI = 7 days	Spray thoroughly to wet upper and lower leaf surfaces and stems. Do not repeat applications more than once a week.

11/8

Continued....

Vegetables, Berries, Citrus Fruit, Stone Fruits, Pome Fruits & Nuts	Insects	Maximum Application Rate, Tsp. of Product/ 1 Gal. of Water	Retreatment Interval, Days	Preharvest Interval, Days	Comments
Citrus (grapefruit, lemon, lime, oranges, & tangerine)	avocado leafroller, California orangedog, citrus cutworm, citrus root weevil, fruitree leafroller, and orange tortrix.	4 tsp./gallon	7 days	PHI = 5 days	Do not apply less than 10 gallons dilute spray per mature tree. For trees taller than 10 feet, consider hiring a licensed professional. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.
Filbert; Pecan; Walnut;	filbert aphid, filbert leafroller, filbertworm, pecan leaf phylloxera, pecan nut casebearer, pecan spittlebug, pecan weevil, walnut caterpillar, codling moth, calico scale, and European fruit lecanium.	4 tsp./gallon	7 days	PHI = 14 days	For trees taller than 10 feet, consider hiring a licensed professional. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.

11/1

10/11

ORNAMENTALS:

Roses, Flowers, Shrubs and Shade Trees - Do not use on Boston Ivy. Mix 4 teaspoons of product per gallon of water to control ants, armyworm, bagworm, boxelder bugs, Japanese beetle, blister beetle, Elm leaf beetle, flea beetle, willow leaf beetle, leafhoppers, tent caterpillar, June beetle, gypsy moth, leafrollers, birch leafminer, mealybugs, exposed thrips, stink bugs, oak leafminer, periodical cicada, and ticks. Apply dilute sprays to obtain thorough coverage of both upper and lower leaf surfaces. To control scale insects, treat trunks, stems and twigs in addition to plant foliage. The use of spreader-sticker may improve residual control. For trees taller than 10 feet, consider hiring a licensed professional. Spray thoroughly to wet upper and lower leaf surfaces, stems and branches. Do not repeat applications more than once a week.

Lawns and Ornamental Turf - For best results, mow lawn and water before spraying. Mix 1 pint of product in 15 gallons of water and spray over 1,250 square feet to control ants, armyworm, bluegrass billbug, centipedes, chiggers, cutworm, earwigs, fall armyworm, fleas, grasshoppers, June beetle, leafhoppers, millipedes, mosquitoes, sod webworm (lawn moth), sowbug, springtails, ticks, chinch bugs.

LIMITED WARRANTY AND DISCLAIMER.

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

The terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.

10/11

APPENDIX:

Advertising claims which may be presented on the container label:

- Kills Over 100 Insects on Listed Vegetables, Fruits, Ornamentals and Lawns.