

ACCEPTED
MAY 4, 1973
U.S. PATENT OFFICE

33155-536

NET CONTENTS
ONE QUART
U.S. MEASURE

ACME®



DO NOT USE IN THE HOME.

Lawn and Garden Insect Control

Spray Concentrate. Contains Diazinon Insecticide.
*Diazinon, a registered trademark of Ciba-Geigy Corp.

ACTIVE INGREDIENTS

O, O-diethyl O-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate 12.5%

Aromatic Petroleum Derivative Solvents 72.6%

INERT INGREDIENTS: 14.9%

TOTAL 100.0%

DO NOT REUSE CONTAINER DESTROY WHEN EMPTY

CAUTION: KEEP OUT OF REACH OF CHILDREN

See side panel for additional cautions.

EPA Reg. No. 577-536AA

CAUTION: COMBUSTIBLE MIXTURE

Caution: Keep out of reach of children.

Harmful if swallowed. May be absorbed through skin. Do not breathe spray mist. Avoid contact with skin, eyes, and clothing. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs.

Do not use on household pets or humans. Do not permit children or pets to go onto the sprayed grass until the spray has completely dried. Keep out of reach of domestic animals.

Do not use, pour, spill or store near open flame.

Do not contaminate ornamental fishponds.

Fish and Wildlife Caution: This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Sod Webworm Spray Lawn and Garden Insect Control

Kills listed outdoor insects which destroy lawns, roses, flowers, trees, and shrubs or annoy pets and people. Contains "Diazinon" insecticide which provides positive control of many "hard-to-kill" sucking and chewing insects. Sod Webworm Spray works fast and remains effective for an extended period of time.

How to Mix and Apply

Step #1: Determine size of spraying job. For lawns and vegetables measure the area to be sprayed. For all other applications apply as a thorough cover spray. Make applications when insects first appear, or in advance of an expected infestation. Repeat applications as needed or as noted in the rate chart.

Step #2: Use clean sprayer. Carefully measure and mix the amount of Sod Webworm Spray and water as suggested in the chart below.

Step #3: Spray as directed in rate chart. Use coarse drenching sprays for lawns, soil, and other surface applications. Use medium-fine mist sprays for trees and plants. Thorough coverage is important. Flush sprayer with clean water after each use.

Problem Area

Lawns: Bluegrass, Mignon, Fescue, St. Augustine, Dichondra, Bent, Bermuda

Roses and Flowers: Rose, Azalea, Chrysanthemum, Gladiol, Carnations

Trees and Shrubs: Oak, Maple, Locust, Holly, Boxwood, Pine, Spruce, Willow

Fruits: Apple, Peach, Orange, Plums, Grape, Cherry, Strawberry

Vegetables: Tomato, Carrot, Potatoes, Bean, Cucumber, Onion, Cantaloupe, Pepper, Cabbage

Cabbage, Carrot, Corn, Lettuce, Peas, Pole Bean, Radishes, Red Beet, Snap, Bean, Tomato, Turnip

Nuisance Pests in Outside Areas:

RATE CHART FOR APPLYING SOD WEBWORM SPRAY

Problem Area	Pests	Rate and Area Covered	Notes
Lawns	Chinch bug, Sod webworm, Bermudagrass, Ant, Tick, Earwig, Lawn tick, Millipede	4 tbsps. (2 fl. oz.) in 3 gal. of water covers 125 sq. ft. 16 fl. oz. in 12 gal. of water covers 1,000 sq. ft.	Thoroughly wet-down grass. Follow with re-applying. For Lawn Billbug, water down thoroughly after treatment. *In the north, if Chinch Bugs are the only insect problem, a lower rate of application can be used. 4 tbsps. in 3 gals. of water covers 250 sq. ft.
	Rhizoglyphus, Ground crawler	4 tbsps. (2 fl. oz.) in 3 gal. of water covers 125 sq. ft. 16 fl. oz. in 12 gal. of water covers 1,000 sq. ft.	Apply when Grub Crawler has emerged.
Roses and Flowers	Aphid, Mite, Groundhopper, Flea	4 tbsps. (2 fl. oz.) in 3 gal. of water. Apply as a thorough foliar spray.	Spray on underside of leaves. Wet plants to the drooping point.
	Aphid, Mite, Scale crawler, Leaf miner, Bagworm	4 tbsps. (2 fl. oz.) in 3 gal. of water. Apply as a thorough foliar spray.	Wet leaves and branches to the drooping point. Try to penetrate dense foliage.
Fruits	Codling moth, Mite, Aphids, Leaf roller, Scale crawler	4 tbsps. (2 fl. oz.) in 3 gal. of water. Apply as a full cover spray.	Start spraying after blossoms and repeat at 7 to 14 day intervals if needed. Do not spray fruit within 21 days of picking.
	Bear beetle, Aphid, Leaf miner, Cabbage worm, Cucumber beetle	4 tbsps. (2 fl. oz.) in 3 gal. of water covers 1,000 sq. ft.	Generally, insecticide should be applied before or after picking. Do not spray within one day of picking. Tomato: Do not spray before or after picking. Do not pick any other crops to be eaten for 14 days.
Vegetables	Cabbage, Carrot, Corn, Lettuce, Peas, Pole Bean, Radishes, Red Beet, Snap, Bean, Tomato, Turnip	Apply 4 tbsps. (2 fl. oz.) per 1,000 sq. ft. as a drenching spray. Do not spray until plants are well established.	Make application at least 14 days before harvest. Thoroughly wet plants to the top 1/2 inch of soil.
	Nuisance Pests in Outside Areas	Earwig, Ant, Cockroach, Boxelder bug, Chinch bug, Spider, Tick, Ground Squirrel	Thoroughly spray window frames, doors, sills, screens, porch and patio walls and ceiling, foundations, walks, garbage cans, and into cracks, and other places where insects may hide. Do not treat plants with this dosage rate.

See chart for complete insect list. Use dosages shown unless otherwise indicated. Do not spray on plants that are stressed or in bloom. Do not spray on plants that are in fruit. Do not spray on plants that are in bloom. Do not spray on plants that are in fruit.

LD 364V2 3 73 U.S. Patent No. 2,754,245
Sold by
THE SHERWIN-WILLIAMS CO. • Cleveland, Ohio 44101
Made in U.S.A.