

Box # 33955 - 538

FM - 10

181

ACME®

89 11

**Wasp & hornet
jet spray**

squirts a stream 10 or
more feet



KILLS: Hornets, Wasps, Yellow Jackets

ACTIVE INGREDIENTS

1,1-Dichloro-2,2-bis(4-chlorophenyl)ethane	0.1600
2,2,4,4-Tetrahydro-3H-pyridin-3-one	0.0200
Related Compounds	0.0000
Aromatic Petroleum Hydrocarbons	0.0000
Petroleum Distillate	26.3750
INERT INGREDIENTS	73.2950
TOTAL	100.0000

© 1987 G. P. B. I. / Gordon Corporation
U.S. Patent No. 3,466,007

KEEP OUT OF REACH OF CHILDREN

CAUTION

**HARMFUL OR FATAL IF SWALLOWED
CONTENTS UNDER PRESSURE**

See back panel for additional
precautionary statements

NET WEIGHT 13 OUNCES



**STOP! READ THE ENTIRE LABEL FIRST
OBSERVE ALL PRECAUTIONS AND
FOLLOW DIRECTIONS CAREFULLY**

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals

CAUTION Avoid inhalation of spray mist.
Avoid contact with skin & eyes.
Keep out of reach of children.
Do not contaminate food or feedstuffs.

Physical & Chemical Hazards

FLAMMABLE Contents under pressure. Do not use near
fire, sparks or flame. Do not puncture or otherwise
damage. Exposure to temperatures above 130°F may cause
bursting.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner
which starts with its labeling.

STORAGE & DISPOSAL

Keep out of the reach of children. **WHEN CONTAINER IS
EMPTY, WRAP AND PUT IN TRASH COLLECTION. DO
NOT ATTEMPT TO REUSE.**

INSTRUCTIONS (Cont)

Acme Wasp & Hornet Jet Spray may be used outdoors in attics,
warehouses and other storage areas where foods and feedstuffs
are not exposed.


FOR OUTDOOR USE Application should be made in late eve-
ning when hornets, wasps and yellow jackets are at rest. Hold
can upright, remove cap, point valve opening away from face, and
with wind at back aim at nests in trees, bushes, under eaves,
chimneys, cracks of crevices, etc.

FOR INDOOR USE Point valve away from face. Aim directly
at nests of hornets, wasps and yellow jackets under attics, etc.
Do not spray in areas where foods and feedstuffs are exposed
or handled.

GENERAL The readily saturate nest with spray, or treat with as
many insects as possible. To prevent escape from the spray mist,
and to this do not stand directly under nest. Best results can be
obtained if nest is sprayed in late evening when most insects are
at rest. If reinfestation occurs, repeat treatment.

EPA REG. NO. 11095-618
EPA EST. NO. 11-15 IN 1

689 581



07 02 1981

L. 33955-538