

**CERTAIN WHITE GRUBS** (larvae of European Chafer & Japanese Beetle): Apply at the rate of 5 to 10 lbs. per 500 sq. ft. as directed above. For best results, treat when grubs are young and actively feeding near the soil surface, usually during late July and August. Consult your local Agricultural Extension Service specialist for the best time to treat in your area.

**HYPERODES (turfgrass) WEEVILS:** Apply at the rate of 10 lbs. per 1,000 sq. ft. as directed above. Treat suspected problem areas in mid-April and again in mid-May. Consult your local Agricultural Extension Service specialist for the best time to treat in your area.

**SUGGESTED SPREADER SETTINGS FOR 5 LBS. / 1,000 SQ. FT.:**

These are approximate. Each individual spreader may require some adjustment for the proper dosage rate.

SCOTT 38-5	5	ASGROW	5
SEARS	5 1/2	LAWN BEAUTY	3
WARDS	4	JACKSON	5
VAUGHANS	4		

Try to maintain uniform walking speed. Shut off cart while turning.

**NOTICE:** Seller warrants that the product conforms to its chemical description and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions not reasonably foreseeable to Seller, and buyer assumes the risk on any such use.

EPA REG. NO. 33955-54  
EPA EST. NO. 2217-KS-1 635-1076



**ACME®**

**dursban**

**Granular Insecticide**

controls chinch bugs, soil webworms (lawn moths), ants, brown dog ticks, armyworms, chiggers, cutworms, crickets, earwigs, sowbugs (pillbugs), clover mites, fleas, grasshoppers, certain white grubs and hyperodes (turfgrass) weevils infesting home lawns and other ornamental and recreational turf areas.

**ACTIVE INGREDIENT**

Chlorpyrifos (0,0-diethyl (3,5,6-trichloro-2-pyridyl) phosphorothioate 0.5%

**INERT INGREDIENTS**

99.5%  
TOTAL 100.0%

\*DURSPAN is trademark of the Dow Chemical Company, U.S. Patent No. 3,244,565

**Keep Out of Reach of Children**

**CAUTION**

See back panel for additional precautionary statement.

**NET WEIGHT TEN POUNDS**

**ACCEPTED**

OCT 17 1980

U.S. Federal Insecticide, Fungicide, and Rodenticide Act

83 725 341

**STOP! READ THE ENTIRE LABEL AND ALL CAUTIONS AND DIRECTIONS FULLY.**

**PRECAUTIONS**

**HAZARDS TO HUMAN AND DOMESTIC ANIMALS:**

**CAUTION--** This product is toxic if swallowed. Do not take internally. Avoid contact with children. Avoid contact with clothing. Flush contaminated water and get medical attention if symptoms develop or persist. Wash thoroughly. Wash contaminated clothing. Keep children and pets away from treated areas. Material is washed into water. **Note to Physician:** Chlorpyrifos is a cholinesterase inhibitor. Treat symptoms with atropine as an antidote.

**ENVIRONMENTAL HAZARDS:**

This product is toxic to birds and wildlife. Keep out of areas where wildlife apply where runoff is likely to contaminate water by runoff. Dispose of wastes in accordance with instructions specified on this label.

**DIRECTIONS FOR USE**

**GENERAL CLASSIFICATION:** It is a violation of federal law to use this product in a manner inconsistent with the directions on the label.

**DISPOSAL:**

Do not reuse empty containers. Dispose of in trash collection.

**USES:**

- Chinch bugs
- Soil webworms (lawn moths)
- Ants
- Brown Dog ticks
- Armyworms
- Chiggers
- Cutworms

**NOTE:** Do not apply to dry grass. Apply Acme Dursban uniformly at the rate of 5 to 10 lbs. per 500 sq. ft. to non-irrigated areas. Apply by hand or power operated spreader designed for granular materials. Treat when insects are young. To help obtain uniform coverage, apply the required amount in two applications, at a right angle to each other. Thoroughly wash the insecticide into the soil. Remember, keep children and pets away from treated areas until the material is dry.

**ACME®**

# dursban

## Granular Insecticide

controls chinch bugs, soil webworms (lawn moths), ants, brown dog ticks, armyworms, chiggers, cutworms, crickets, earwigs, sowbugs (pillbugs), clover mites, fleas, grasshoppers, certain white grubs and hyperodes (turfgrass) weevils infesting home lawns and other ornamental and recreational turf areas.

### ACTIVE INGREDIENT

Chlorpyrifos (0,0-diethyl 1-(3,5,6-trichloro-2-pyridyl) phosphorothioate) 0.5%

INERT INGREDIENTS 99.5%  
TOTAL 100.0%

©1985 ACME, a trademark of the D.W. Chemical Company, U.S. Patent No. 3,244,183

Keep Out of Reach of Children

## CAUTION

See back panel for additional precautionary statement.

NET WEIGHT TEN POUNDS

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL CAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

### PRECAUTIONARY STATEMENTS

#### HAZARDS TO HUMANS & DOMESTIC ANIMALS:

**CAUTION** This product may be harmful if swallowed. Do not take internally. Keep out of reach of children. Avoid contact with eyes, skin, and clothing. Flush contaminated eyes with plenty of water and get medical attention if irritation develops or persists. Wash thoroughly after handling. Wash contaminated clothing before reuse. Keep children and pets off treated areas until the material is washed into the soil and grass is dry. **Note to Physician:** Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

#### ENVIRONMENTAL HAZARDS:

This product is toxic to fish, birds, and other wildlife. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

### DIRECTIONS FOR USE

#### GENERAL CLASSIFICATION

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

### DISPOSAL

Do not reuse empty container. Wrap and put in trash collection.

#### USES

Chinch bugs	Crickets
Soil webworms (lawn moths)	Earwigs
Ants	Sowbugs (Pillbugs)
Brown Dog ticks	Clover Mites
Armyworm	Fleas
Chigger	Grasshoppers
Cutworms	

**NOTE:** Do not apply to Dichondra

Apply Acme Dursban uniformly at the rate of 5 lbs. per 1,000 sq. ft. to non-turf areas using any hand or power operated gravity or rotary-type spreader designed for application of granular material. Treat when insects appear; retreat as necessary. To help obtain uniform distribution, apply one-half the treated amount in one direction; then apply the balance at a right angle. **Immediately after application,** thoroughly water the treated turf to wash the insecticide into the zone of insect infestation. Remember, keep children and pets off treated areas until the material is washed into the soil and grass is dry.

ACCEPTED

OCT 17 1980

U.S. Federal Insecticide, Fungicide, and Rodenticide Act  
33-735-341

47

48