



**STOP! READ THE ENTIRE LABEL  
FIRST OBSERVE ALL CAUTIONS AND  
FOLLOW DIRECTIONS CAREFULLY**

**PRECAUTIONARY STATEMENTS**

**Hazards to Humans & Domestic Animals**

**CAUTION** - Irritant. May cause skin irritation. Avoid contact with eyes. Wash thoroughly after use. Do not eat, drink, or smoke while using this product. Do not use near children or pets.

**Statement of Practical Treatment**

**If Swallowed** - Do not induce vomiting. Drink plenty of water. If symptoms persist, seek medical attention.

**If Inhaled** - Remove victim to fresh air. Administer first aid as directed on label.

**If In Eyes** - Flush eyes with plenty of water for at least 15 minutes. Get medical attention immediately.

**If On Skin** - Wash contaminated skin with soap and water.

**Environmental Hazards**

This product is toxic to fish, birds, and other wildlife. Birds feeding on treated areas may be killed. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from target area. Apply this product only as specified on this label.



**Bees** - This pesticide is highly toxic to bees exposed to direct treatment or residues remaining on the treated area. Do not apply when bees are actively visiting the crop cover, top or weeds blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

**DIRECTIONS FOR USE**

Use in accordance with federal law to use this product in a manner consistent with its labeling.

**DISPOSAL** - Do not reuse empty container. Wrap and put in trash collection.

EPA REG. NO. 33955-545  
EPA EST. NO. 2217-KS-1



a foliage insecticide for vegetables,  
fruit and flowers, and, a preplant soil  
insecticide for vegetables.

**ACTIVE INGREDIENT**  
Diazinon (0.02% diazinon, 0.02% methyl 4-phosphorothioate) 4.0  
**INERT INGREDIENTS** 96.0  
Total 100.0

**Keep Out of Reach of Children  
CAUTION**

**NET WEIGHT ELEVEN OUNCES**

Use a hand operated duster or shake from this convenient shaker top canister.

**FOLIAGE APPLICATIONS**  
Use on the following plants and vegetables:  
Beans, Broccoli, Cabbage, Cauliflower, Carrots, Radishes, Onions, Cucumbers, Summer Squash, Winter Squash, Melons, Apples, Pears, Strawberries, Roses, Flowers, Shrubs, Ornamentals.

PLANT	LAST APPLICATION BEFORE HARVEST	PROTECTION FROM
BEANS	7 days	
PEAS	14 days before harvest	
POTATOES	25 days	Aphids & Leafhoppers
TOMATOES	7 days	
BEETS	14 days	
BROCCOLI	5 days	
CABBAGE	7 days	
CAULIFLOWER	5 days	Aphids
CARROTS	10 days	
RADISHES	10 days	
ONIONS	10 days	
CUCUMBERS	7 days	Aphids, Thrips
SUMMER SQUASH	7 days	Miners & Mites
WINTER SQUASH	7 days	
MELONS	3 days	
APPLES	14 days	
PEARS	14 days	Aphids & Mites
STRAWBERRIES	5 days	Aphids, Cyclamen & Two Spotted Mites
ROSES	not applicable	Aphids, Mites, Leafhoppers, Mealy bugs, White fly, Scale, crawlers & Lace bugs
FLOWERS		
SHRUBS		
ORNAMENTALS		

**USE PRECAUTIONS**  
Do not use on certain ferns, poinsettias, hibiscus, and gardenias since injury may occur. Not effective against leaf or spider mites on evergreens. Do not apply when bees are foraging in the garden.

**SOIL APPLICATION** - Use on planting the following vegetables:  
Beans, Broccoli, Cabbage, Cauliflower, Carrots, Radishes, Onions, Cucumbers, Summer Squash, Winter Squash, Melons, Apples, Pears, Strawberries, Roses, Flowers, Shrubs, Ornamentals.

To control Cutworms, Seed corn Maggots, Wire worms, etc., apply 1/2 ounce per 100 square feet of soil at planting time. For control of other soil insects, apply 1/2 ounce per 100 square feet of soil at planting time.

**SPECIAL USES**  
On onion maggot, apply 1/2 ounce per 100 square feet of soil at planting time. On radishes, apply 1/2 ounce per 100 square feet of soil at planting time.

ACCEPTED

DEC 0 1981