

ACCEPTED
 JUN 16 1982
 U.S. Environmental Protection Agency
 Federal Insecticide, Fungicide, and Rodenticide Act
 Registration for the pesticide
 Chlorpyrifos
 EPA Reg. No. 33955-547

PRECAUTIONARY STATEMENTS
Hazards to Humans & Domestic Animals
WARNING—This product may be fatal if swallowed. Do not take internally. It can be absorbed through skin and may be innocuous to both skin and eyes. Do not get on skin or into eyes. Wash thoroughly after handling. Wash contaminated clothing before reuse. Do not breathe vapors or spray mist. Do not allow spray to get on food crops. Keep away from food, feedstuffs and domestic water supplies. Keep container closed when not in use and handle concentrate only in a ventilated area.
Statement of Practical Treatment
If Swallowed—Do not induce vomiting. Call a physician immediately. **If On Skin**—In case of contact, remove contaminated clothing and immediately wash skin with soap and water. **If In Eyes**—Flush eyes with plenty of water for 15 minutes and call a physician.
Note to Physician—Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine only by injection is an antidote.
Environmental Hazards
 This product is toxic to fish, birds and other wildlife. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from target area. Do not contaminate water by cleaning of equipment or disposal of wastes.
Physical or Chemical Hazards
COMBUSTIBLE! Do not use or store near heat or open flame.

DIRECTIONS FOR USE
 It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
 EPA REG. NO. 33955-547
 EPA EST. NO. 2217-KS-1
 (Cont. on right panel)
 927-582



Controls: Chinch Bugs, Sod Webworms, Cutworms, Grasshoppers, Mealybugs, Leafhoppers and many other insects in home lawns and ornamental plants.

ACTIVE INGREDIENTS

Chlorpyrifos (0.025% ethyl ester)	6.62%
(3,5,6-trichloro-2-pyridyl) phosphorothioate	88.86%
Kyline Range Aromatic Petroleum Sulfonates	4.52%
INERT INGREDIENTS	
This Product Contains	TOTAL 100.00%
0.50 lb. Chlorpyrifos per gallon	

DURSBAN® is a registered trademark of The Dow Chemical Company, U.S. Patent No. 3,244,588

Keep Out of Reach of Children
WARNING
 See side panels for additional precautions and statement of practical treatment.

NET CONTENTS ONE PINT

DIRECTIONS FOR USE (Cont. from left panel)
 It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE & DISPOSAL
 Do not contaminate water, food or feed by storage or disposal. Do not reuse empty container. Wrap and put in trash collection.

Home Lawns
 For best results, lawn should be moist at time of treatment. Dilute Acme Dursban in enough water to obtain complete and uniform coverage. Apply as a coarse low pressure spray using suitable application equipment. Thoroughly water lawn after treatment to wash the insecticide into the grass, except as otherwise noted. To control ants, armyworms, Brown dog ticks, chiggers, chinch bugs, clover mite crickets, cutworms, earwigs, fleas, grasshoppers and sod webworms, use 6 fl. oz. or 12 lbs. and spray on 1,000 sq. ft. when pests first appear. Retreat when needed. For sod webworms, delay watering or mowing for 12 to 24 hours after treatment. To control turfgrass (Hyperdoel) weevil, use 12 fl. oz. or 24 lbs. and spray suspected problem area in mid April and again in mid May or as recommended by the local Agricultural Extension Service specialist in your area.

Ornamentals
NOTE: Do not use on azaleas, camellias, poinsettias, rose bushes or variegated ivy because injury to these plants may occur. Treat flowers, shrubs, vines, shade and flowering trees and evergreens found to be infested. Dilute with water as instructed and apply using a suitable hand or power sprayer in a manner to provide complete and uniform coverage. For best results, apply a wetting spray to both upper and lower leaf surfaces and infested limb and trunk areas. Attempt to penetrate dense foliage but avoid over spraying to the point of excessive run off. To control aphids, bagworms, Eastern tent caterpillar, Fall webworm, grasshoppers, mites, Orangestriped oakworm, spittlebugs, whiteflies, Yellownecked caterpillar, use 2 fl. oz. or 4 lbs. in 3 gals. water. To control cutworms, leafhoppers, mealybugs, Mamee webworm, Redhumped caterpillar, thrips (leopard), use 3 fl. oz. or 6 lbs. in 3 gals. water. To control Ash borer, Lilac borer, scale crawlers, Oak lecanium, pine needle leaf white, peach use 8 fl. oz. or 16 lbs. in 3 gals. of water.
NOTICE: Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, Seller makes no claims other than those stated on this label.

COMPANY
NUMBER 34704-

PRODUCT
SERIAL NO. 10

LABEL
APPROVAL DATE 7-28-82