

ACME® DURSBAN® INSECTICIDE FOR LAWNS & ORNAMENTALS

Controls Chinch Bugs, Sod Webworms, Cutworms,
Grasshoppers, Mealybugs, Leafhoppers and other
insects in home lawns and ornamental plants.

ACTIVE INGREDIENT:

Chlorpyrifos: 0,0-diethyl 0-(3,5,6-trichloro-2-pyridyl) phosphorothioate..... 6.62%

INERT INGREDIENTS*..... 93.38%

TOTAL 100.00%

This Product Contains:

0.50 lb. Chlorpyrifos insecticide per gallon.


DURSBAN® is a trademark of DowElanco for chlorpyrifos insecticide.

*CONTAINS XYLENE RANGE AROMATIC PETROLEUM SOLVENT

KEEP OUT OF REACH OF CHILDREN

WARNING

See side panels for Precautionary Statements
and Statement of Practical Treatment.



NET CONTENTS ONE QUART

9271 AP
EPA REG NO. 33955-547
EPA EST NO. 2217-KS-1

Manufactured by ACME DIVISION
PBI/GORDON CORPORATION
KANSAS CITY, MISSOURI 64101

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS
AND FOLLOW DIRECTIONS CAREFULLY.

2/4

PRECAUTIONARY STATEMENTS

Hazards to Humans & Domestic Animals:

WARNING: This product may be fatal if swallowed. Do not take internally. It can be absorbed through skin and may be injurious to both skin and eyes. Do not get on skin or into eyes. Wash thoroughly after handling. Wash contaminated clothing before reuse. Do not breathe vapors or spray mist. Keep away from food, feedstuffs and domestic water supplies. Keep container closed when not in use and handle concentrate only in a ventilated area.

KEEP CHILDREN AND PETS OFF OF TREATED AREAS UNTIL SPRAY IS WASHED INTO THE SOIL.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician or Poison Control Center. Do not induce vomiting because of possible aspiration hazard.

IF ON SKIN: In case of contact, remove contaminated clothing and immediately wash skin with plenty of soap and water.

IF IN EYES: Flush eyes with plenty of water and call a physician.

IF INHALED: Remove to fresh air if symptoms of cholinesterase inhibition appear and get medical attention.

NOTE TO PHYSICIAN: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine only by injection is an antidote.

Environmental Hazards: This product is toxic to fish, birds, and other wildlife. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from target area. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

Physical or Chemical Hazards: COMBUSTIBLE! Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes. For use on plants intended for aesthetic purposes, ornamental gardens or parks, or on golf courses or lawns and grounds.

STORAGE & DISPOSAL

STORAGE: Store in original bottle in a locked storage area inaccessible to children or pets.

PESTICIDE DISPOSAL: Securely wrap original container in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse empty container. Discard in trash.

HOME LAWNS, PICNIC SITES, and RECREATIONAL AREAS

SITES: Use Acme® Dursban® Insecticide for Lawns and Ornamentals for the control of pests listed in Table 1 for home lawns, picnic sites and recreational areas.

SPRAY EQUIPMENT AND SPRAY VOLUME: Apply Acme® Dursban® Insecticide for Lawns and Ornamentals in water with hydraulic power sprayers, knapsack sprayers, hand-operated sprayers, or other suitable equipment. Spray volumes ranging from 1 to 5 gallons of product per 1000 square feet will provide sufficient coverage. Apply as a coarse low pressure spray pattern.

SPRAY PREPARATION: See Table 1 for the preparation of spray solutions.

APPLICATION SCHEDULE: Treat problem areas when pests first appear. If retreatment is needed, observe a 7 day application interval.

MANAGEMENT CARE: For best results, the lawn should be moist at the time of treatment. Except for sod webworm, the lawn should be irrigated, sprinkled, or watered within 12 hours of treatment. For sod webworm, delay mowing and watering for 12 to 24 hours after treatment.

PRECAUTIONS: KEEP CHILDREN AND PETS OFF OF TREATED AREAS UNTIL SPRAY IS WASHED INTO THE SOIL.

TABLE 1. DOSAGE RATES OF ACME DURSBAN IN HOME LAWNS, PICNIC SITES, and RECREATIONAL AREAS.

Site	Pests	Amount of Acme Dursban		Directions
		Fluid Ounces per 1000 square feet	Tablespoons per 1000 square feet	
Home lawns, Picnic Sites, and Recreational areas.	Ants, Armyworms, Brown Dog Ticks, Chiggers, Chinch bugs, Clover Mites, Crickets, Cutworms, Earwigs, Fleas, Grasshoppers	6	12	Spray volumes ranging from 1 to 5 gallons per 1000 square feet should provide complete coverage. For best results, lawn should be moist at the time of application, and the lawn should be irrigated, sprinkled or watered after application. Retreat as needed.
	Sod Webworms	6	12	Spray volumes ranging from 1 to 5 gallons per 1000 square feet should provide complete coverage. For best results, lawn should be moist at time of application. Delay mowing and watering for 12 to 24 hours after treatment. Retreat as needed.
	Turfgrass Weevil (Hyperodes)	12	24	Spray problem areas in mid-April and mid-May, or as recommended by your State Agricultural Extension Service.

ORNAMENTALS, OUTDOORS

SITES: Use Acme® Dursban® Insecticide for Lawns and Ornamentals for the control of the insect pests listed in Table 2 for flowers, shrubs, shade trees, flowering trees, evergreens, vines, and groundcovers.

SPRAY EQUIPMENT AND SPRAY VOLUME: Apply Acme® Dursban® Insecticide for Lawns and Ornamentals in water with hydraulic/power sprayers, knapsack sprayers, hand-operated sprayers, or other suitable equipment. Apply a coarse spray to wet thoroughly the leaves, branches, limbs, and trunks. The wetting of infested limbs and trunks and the wetting of the upper and lower leaf surfaces will provide the best results. Avoid spray runoff.

SPRAY PREPARATION: See Table 2 for the preparation of three gallons of spray solution.

APPLICATION SCHEDULE: Treat ornamentals when pests first appear. If needed, repeat treatments at 7 day intervals.

