

4-13-81

**PRECAUTIONARY STATEMENTS**

**Hazards to Humans & Domestic Animals:**

**WARNING**—This product may be fatal if swallowed. Do not take internally. It can be absorbed through skin and may be injurious to both skin and eyes. Do not get on skin or into eyes. Wash thoroughly after handling. Wash contaminated clothing before reuse. Do not breathe vapors or spray mist. Do not allow spray to get on food crops. Keep away from food, feedstuffs and domestic water supplies. Keep container closed when not in use and handle concentrate only in a ventilated area.

**Statement of Practical Treatment:**

**If Swallowed:** Do not induce vomiting. Call a physician immediately. **If On Skin:** In case of contact, remove contaminated clothing and immediately wash skin with soap and water. **If In Eyes:** Flush eyes with plenty of water for 15 minutes and call a physician.

**Note to Physician:** Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine only by injection is an antidote.

**Environmental Hazards:**

This product is toxic to fish, birds and other wildlife. Keep out of lakes, streams and ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from target area. Do not contaminate water by cleaning of equipment or disposal of wastes.

**Physical or Chemical Hazards:**

**COMBUSTIBLE!** Do not use or store near heat or open flame.

**STORAGE & DISPOSAL**

Do not contaminate water, food or feed by storage or disposal. Do not reuse empty container. Wrap and put in trash collection.

EPA REG NO 33955-547  
EPA EST. NO. 2217-RS 1

927/381



**ACME**

17-14

**DURSBAN**  
insecticide  
for lawns & ornamentals

**Controls:** Chinch Bugs, Sod Webworms, Cutworms, Grasshoppers, Mealybugs, Leafhoppers and many other insects in home lawns and ornamental plants.

**ACTIVE INGREDIENTS:**

Chlorpyrifos (0,0-diethyl O-  
(3,5,6-trichloro-2-pyridyl) phosphorothioate 6%  
Xylene-Range Aromatics, Petroleum Solvent 88%

**INERT INGREDIENTS**

4.52%

**This Product Contains:**

0.50 lb Chlorpyrifos per gallon

**TOTAL 100.00%**

\*DURSBAN® is a registered trademark of The Dow Chemical Company U.S. Patent No. 3,244,586

**Keep Out of Reach of Children**  
**WARNING**

See side panels for additional precautions and statement of practical treatment.

**NET CONTENTS ONE PINT**

**ACCEPTED**  
**MAY 21 1981**  
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for the pesticide registered under EPA Reg. No. 33955-547

**DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

**Home Lawns:**

For best results, lawn should be moist at time of treatment. Dilute Acme Dursban in enough water to obtain complete and uniform coverage. Apply as a coarse, low-pressure spray, using suitable application equipment. Thoroughly water lawn after treatment to wash the insecticide into the grass, except as otherwise noted.

To control ants, armyworms, brown dog ticks, chiggers, chinch bugs, clover mite, crickets, cutworms, earwigs, fleas, grasshoppers and sod webworms, use 6 fl oz or 12 tbs. and spray on 1,000 sq ft when pests first appear. Retreat when needed. For sod webworms, delay watering or mowing for 12 to 24 hours after treatment. To control turfgrass (Hyperodes) weevil, use 12 fl oz or 24 tbs. and spray suspected problem area in mid-April and again in mid-May, or as recommended by the local Agricultural Extension Service specialist in your area.

**Ornamentals:**

Treat flowers, shrubs, vines, shade and flowering trees and evergreens found to be infested. Dilute with water as instructed and apply using a suitable hand or power sprayer in a manner to provide complete and uniform coverage. For best results, apply a wetting spray to both upper and lower leaf surfaces and infested limb and trunk areas. Attempt to penetrate dense foliage but avoid over spraying to the point of excessive run-off.

To control aphids, bagworms, Eastern tent caterpillar, Fall webworm, grasshoppers, mealybugs, mites, Orange-striped oakworm, spittlebugs, whiteflies, Yellow-necked caterpillar, use 2 fl oz or 4 tbs in 3 gals water. To control cutworms, leafhoppers, mealybugs, Mameose webworm, Red-tipped caterpillar, Stages (exposed), use 3 fl oz or 6 tbs in 3 gals water. To control Ash borer, Lilac borer, scale crawlers: Oak lecanium, pine needle, tea white peach, use 8 fl oz or 16 tbs in 3 gals of water.

**NOTICE:** Seller warrants that this product conforms to the ingredient statement on the label. Since conditions of use, such as weather, compatibility with other chemicals, and condition of application equipment will vary, Seller makes no claims other than those stated on this label.

19