

34052-9

8/20/1985

182

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

Disinfectant-Sanitizer - Use with a cloth, mop or mechanical spray device. When applied with a mechanical spray device, surface must be sprayed until thoroughly wetted. Treated surfaces must remain wet for 10 minutes. Fresh solution should be prepared daily or when the use solution becomes visibly dirty.

Fungicide-Virucide-Deodorizer with Organic Soil Tolerant - Disinfectant in Hospitals, Nurseries, Restrooms and Other Health Care Institutions. For disinfecting floors, walls, countertops, bathing areas, lavatories, bedframes, tables, chairs, garbage pails and other hard non-porous surfaces.

For Hospital, Institutional, Industrial, School, Dairy and Other Farm Use - Add 1 1/2 oz. ENFORCER to 5 gallons water. Apply to previously cleaned hard surfaces. At this use level, ENFORCER is effective against Pseudomonas aeruginosa, Staphylococcus aureus and Salmonella choleraesuis in the presence of 5% blood serum which established by the AOAC Use-Dilution Test.

Virucidal Performance - At a 1 1/2 oz./5 gallon use level, ENFORCER was evaluated in the presence of 5% serum and found to be effective against the following viruses: Herpes simplex type 1, 2, Vaccinia, and Influenza A₂ (Hong Kong) on inanimate hard non-porous environmental surfaces.

Fungicidal Performance - At 1/2 oz. to 6 gallons of water use level, ENFORCER is an effective fungicide against Trichophyton meningophyces (the athlete's foot fungus) when used on surfaces in areas such as locker rooms, dressing rooms, shower and bath areas and exercise facilities, utilizing the AOAC Fungicidal Test.

Disinfection of Barber Tools - Preheated barber tools (such as combs, brushes, razors, and scissors) can be disinfected by immersing in 0.2% aqueous solution of ENFORCER.

Disinfection of Poultry Equipment, Animal Quarters and Kennels - Poultry brooders, watering troughs, feeding equipment and other animal quarters (such as stalls and kennel area) can be disinfected after thorough cleaning by applying a solution of 1 oz. ENFORCER to 6 gallons of water. Small contents should be immersed in this solution. Prior to disinfection, all poultry, other animals and their feeds must be removed from the premises. This includes emptying all troughs, racks and other feeding and watering appliances. Remove all litter and droppings from floors, walls and other surfaces occupied or traversed by poultry or other animals.

After disinfection, ventilate buildings, coops and other closed spaces. Do not house poultry, or other animals or employ equipment until treatment has been absorbed, set or dried. All treated equipment that will contact feed or drinking water must be rinsed with potable water before reuse.

Disinfection in Non-Medical Institutions, Industry and Schools - Add 1 oz. Enforcer to 6 gallons of water for disinfecting floors, walls, countertops, bathing areas, lavatories, tables, chairs, garbage pails, and other hard, non-porous surfaces.

ACCEPTED
34052-9
AUG 20 1985

ENFORCER

Under the Federal Insecticide, Fungicide, and Rodenticide Act, EPA has registered this product for use as a Disinfectant-Sanitizer with Organic Soil Tolerant for Hospital, Institutional, Industrial, School, Dairy and Other Farm Use.

Active Ingredients

- Octyldecyl dimethyl ammonium chloride 6.0 %
- Didecyl dimethyl ammonium chloride 3.0 %
- Dioctyl dimethyl ammonium chloride 3.0 %
- Alkyl (C₁₂, 50%; C₁₇, 40%; C₁₆, 10%) dimethyl benzyl ammonium chloride 8.0 .06
- Inert Ingredients 80.0 %
- 100.0 %

KEEP OUT OF REACH OF CHILDREN

DANGER

STATEMENT OF PRACTICAL TREATMENT

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse.

If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

SEE LEFT PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EPA Registration No. 34052 - 9

EPA Establishment No. 34052 - CA - 1

Net Contents

MANUFACTURED BY:

H. Wilson Mfg. Co., P.O. Box 481, Jefferson, CA 95849

For Sale For Use And Storage By Maintenance Personnel Only

Sanitizing of Food Processing Equipment and Other Hard Surfaces in Food Contact Locations. For sanitizing food processing equipment, dairy equipment, food utensils, dishes, silverware, glasses, sink tops, countertops, refrigerated storage and display equipment and other hard non-porous surfaces. No Potable water rinse is required.

Wash and rinse all articles thoroughly, then apply a solution of 1 oz. ENFORCER in 10 gallons of water (150ppm active). Surfaces should remain wet for at least one minute followed by adequate draining and air drying. Fresh solution should be prepared daily or when use solution becomes visibly dirty. For mechanical application, use solution may not be reused for sanitizing applications.

Apply to sink tops, countertops, refrigerated storage and display equipment and other stationary hard surfaces by cloth or brush or mechanical spray device. No Potable water rinse is required.

Dishes, silverware, glasses, cooking utensils and other similar size food processing equipment can be sanitized by immersion in a 0.2% aqueous solution of ENFORCER. No Potable water rinse is required.

At 1 oz./10 gallons ENFORCER fulfills the criteria of Appendix F of the Grade "A" Pasteurized Milk Ordinance 1978 Recommendations of the U.S. Public Health Service in waters up to 800 ppm of hardness calculated as CaCO₃ when evaluated by the AOAC Germicidal and Detergent Sanitizer Method against Escherichia coli and Staphylococcus aureus. The washers, tanks and stalls of dairy cows can be sanitized by washing with a solution of 1 oz. ENFORCER in 10 gallons of warm water. No Potable water rinses are required. Use a fresh towel for each cow. Avoid contamination of sanitizing solution by dirt and soil. Do not dip used towel back into sanitizing solution. When solution becomes visibly dirty, discard and provide fresh solution.

Precautionary Statements

Hazardous to Humans and Domestic Animals

DANGER

Keep Out of Reach of Children. Corrosive. Causes eye and skin irritation. Do not get in eyes, on skin, or on clothing. Wear goggles or face shield and rubber gloves when handling. Harmful if swallowed. Do not breathe spray mist. Avoid contamination of food.

Storage and Disposal


- Do not contaminate water, food, or feed by storage disposal. - Do not store on side. - Avoid creasing or impacting of side walls.

PESTICIDE DISPOSAL

Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray or material of this type in violation of Federal Law, if these wastes are disposed of by use according to label instructions, contacts your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL

Triple rinse (or equivalent). Then offer for recycling (reconditioning or puncturing and crushing by a recognized recycling operation, or if allowed by state and local authorities by burning, if burned, stay out of smoke). Metal Container: Triple rinse (or equivalent), then offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. (If container is 1 gallon or less, use the container disposal statement.) Do not reuse empty container (bottle, can, bucket, wrap container and put it to rest).



SANITIZING HATCHING EGGS

Hatching eggs can be sanitized by using a solution of 1 oz. ENFORCER in 10 gallons of water. No Potable water rinse is required. Avoid contamination of sanitizing solution by dirt and soil. Surfaces should remain wet for at least one minute followed by adequate draining and air drying. Fresh solution should be prepared daily or when use solution becomes visibly dirty.