

[34224-5]

Chemrite

CR-174 VAPORITE

INSECTICIDE VAPORIZING SPRAY
CONCENTRATED LIQUID

FOR COMMERCIAL AND INDUSTRIAL USE

IN FOGGING EQUIPMENT AND HAND SPRAYERS

To Kill Accessible Stages of Insects* Infesting Dried Fruit Products In Storage or Being Processed. Close room or warehouse and shut off all ventilating systems. Dispense this product as a fine mist in the air above trays or shelves of dried fruit. Use a spraying device similar to the Microsol machine or use fixed ceiling sprayers that produce a very fine spray. Do not apply to fruit directly. If overhead sprayers are used, be sure that the spray is discharged from a position of at least 6 feet over the top trays of fruit. Use a dosage of 1 gallon of this spray per 50,000 cubic feet of room space. Repeat applications at this dosage in any given month. More than ten applications may also be made if this spray is diluted with more petroleum distillate so that the pyrethrins and synergists in the final spray are reduced proportionately. Do not remain in treated areas. Ventilate the area after treatment is completed. Maximum dilution is 1 part 5192 plus 3 parts oil.

*Indian Meal Moth, Dried Fruit Beetle, Confused Flour Beetle, Saw-Toothed Grain Beetle and Roaches.

To Kill The Adult and Accessible Forms of Insect Pests* of Stored Food (Stored in Multiwall Paper Bags or in Cloth Bags) in Warehouses, Storage Rooms and Similar Locations. Close room or warehouse and shut off all ventilating systems. Use a spraying device similar to the Microsol machine, or use fixed ceiling sprayers that produce a very fine spray. Use a dosage of one fluid ounce of this spray per 1000 cubic feet of room space. Do not remain in treated areas. Ventilate the area after treatment is completed. Repeat applications as necessary for insect control.

*Rice Weevil, Yellow Meal Worm, Cadelle, Confused Flour Beetle, Saw-toothed Grain Beetle, Angoumois Grain Moth, Mediterranean Flour Moth and Indian Meal Moth.

To Kill Flies, Mosquitoes and Gnats. Close room and discharge spray mist as described above. Use a dosage of 1 fluid ounce per 1000 cubic feet of room space. Leave room closed for at least ten minutes. Sweep up and destroy fallen insects. Repeat, as necessary for control. Do not remain in treated areas. Ventilate the area after treatment is completed.

IN PRESSURIZED TANK SPRAYERS FOR WETTING CONTACT SPRAYS

As a Surface Spray to Kill the Accessible Stages of Granary Insects: Rice Weevils, Confused Flour Beetles, Rust Red Flour Beetles, Saw-Toothed Grain Beetles, Cadelles, Meal Moth Larvae and Adults, Cigarette Beetles, Drug Store Beetles and Granary Weevils. Apply as a coarse, wet spray to all areas infested by these insects. Use a dosage of 1 gallon per 1000 square feet. Pay particular attention to cracks and crevices in floor, walls and ceilings. Spray

ACTIVE INGREDIENTS: 100%
Pyrethrins .30%
*Piperonyl butoxide, technical .60%
-N-octyl bicycloheptene dicarboximide 1.00%
Petroleum distillate 98.10%

*Equivalent to .48% (butylcarbityl) (6-propylpiperonyl) ether and .12% related compounds.
-MGK-264, Insecticide Synergist.

EPA REG. NO. 34224-5


CAUTION

KEEP OUT OF THE REACH OF CHILDREN

SEE OTHER PRECAUTIONS ON THE BACK/SIDE PANEL.

NET CONTENTS

Manufactured by
CHEMRITE CORPORATION
12600 SOUTH DAPHNE AVENUE
HAWTHORNE, CALIFORNIA 90250



To Kill Accessible Stages of Insects* Infesting Dried Fruit Products In Storage or Being Processed. Close room or warehouse and shut off all ventilating systems. Dispense this product as a fine mist in the air above trays or shelves of dried fruit. Use a spraying device similar to the Microsol machine or use fixed ceiling sprayers that produce a very fine spray. Do not apply to fruit directly. If overhead sprayers are used, be sure that the spray is discharged from a position of at least 6 feet over the top trays of fruit. Use a dosage of 1 gallon of this spray per 50,000 cubic feet of room space. Repeat applications at this dosage in any given month. More than ten applications may also be made if this spray is diluted with more petroleum distillate so that the pyrethrins and synergists in the final spray are reduced proportionately. Do not remain in treated areas. Ventilate the area after treatment is completed. Maximum dilution is 1 part 5192 plus 3 parts oil.

To Kill The Adult and Accessible Forms of Insect Pests* of Stored Food (Stored in Multiwall Paper Bags or in Cloth Bags) in Warehouses, Storage Rooms and Similar Locations. Close room or warehouse and shut off all ventilating systems. Use a spraying device similar to the Microsol machine, or use fixed ceiling sprayers that produce a very fine spray. Use a dosage of one fluid ounce of this spray per 1000 cubic feet of room space. Do not remain in treated areas. Ventilate the area after treatment is completed. Repeat applications as necessary for insect control.

CAUTION

KEEP OUT OF REACH OF CHILDREN
Harmful if swallowed. Do not apply directly to food. Cover or remove all food and food processing equipment during application. Do not apply space sprays while food processing is underway. Food processing operations do not have to be stopped while applying a wet spray with care and in accordance with the directions and cautions above. After spraying in meat packing plants, bakeries and other food processing plants, wash with potable water all equipment benches shelving etc. where exposed food will be handled. In case washing is impractical, do not allow food to come in contact with treated surfaces for at least 48 hours.

Remove pets and cover fish bowls before spraying. This product is toxic to fish. Keep out of lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of wastes.

Do not re-use empty containers. Wash with water and discard in a safe place. Do not reuse empty containers. Return to manufacturer or distributor for recycling information. Wash and store in a safe place away from water supplies.

Keep container closed. Do not pour, store or spill near heat or open flame.

Note: This spray when applied per the directions and cautions above, conforms with Food Additive Regulations 121.1074, 121.1075, and 121.1205 and the food additive tolerances established thereunder.

MADE FOR INDUSTRIAL & INSTITUTIONAL USE ONLY