

Chemrite

CR-350 BIO-THANARI

DISINFECTANT SANITIZER
DEODORIZER

ACTIVE INGREDIENTS:

n-Alkyl (60% C₁₄, 30% C₁₆, 5% C₁₂, 5% C₁₈)
dimethyl benzyl ammonium chlorides..... 5%
n-Alkyl (68% C₁₂, 32% C₁₄) dimethyl
ethylbenzyl ammonium chlorides 5%

INERT INGREDIENTS:..... 90%
100%

CR-350 is a concentrate and should be diluted before using.

DANGER

KEEP OUT OF REACH OF CHILDREN. "E"
(See other precautionary and first aid statements on left side panel.)

Manufactured By
CHEMRITE CORPORATION
12600 SOUTH DAPHNE AVENUE
HAWTHORNE, CALIFORNIA 90250

CAUTION: Keep out of reach of children

Cold Disinfection

For disinfection of previously cleaned surgical instruments, barber tools and dental equipment, simply submerge instruments into solution containing 1½ ounce CR-350 per gallon of water for 10 minutes.

FOR ROOM FOGGING

DIRECTIONS:

To minimize the danger of cross-contamination from environmental surfaces apply solution of CR-350 as a fog as described below, before applying standard routine terminal cleaning and disinfecting practices. 1. Remove all human, animal, and plant life from room. 2. Open closet doors and drawers. 3. Set up 34" revolving platform in center of room. 4. Mount Jet FOG-MASTER TRI-JET MODEL 6208 or other fogging device delivering equivalent spray at 3 RPM. 5. Fill sprayer reservoir with 1.5 oz. of CR-350 to 1 gallon of water to produce 1150 ppm active ingredients. 6. Set sprayer mechanism to deliver 1 gallon of solution. 7. Fog for 15 minutes for an average 2100 cu. ft. room. For different room sizes, vary spray time proportionately to ensure complete wetting of exposed surfaces. 8. Wait 2 hours before entering room after treatment. 9. Rinse fogging equipment thoroughly with clear water following use.

DANGER

Keep Out of Reach of Children. Corrosive. Causes severe eye and skin damage. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling concentrate. Harmful if swallowed. Avoid contamination of food.

FIRST AID

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash all contaminated clothing before reuse. If swallowed, drink milk, egg whites, gelatin solution, or if these are not available, drink large quantities of water. Call a physician.

NOTE TO PHYSICIAN:

Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

Rinse empty container thoroughly with water and discard.

NET CONTENTS: _____

MADE FOR INDUSTRIAL & INSTITUTIONAL USE ONLY

rite

ACCEPTED
APR 10 1975
 UNDER THE FEDERAL INSECTICIDE
 FUNGICIDE AND RODENTICIDE ACT
 FOR ECONOMIC POISON REGISTERED
 UNDER NO. 34224-8. SUBJECT
 TO ATTACHED COMMENTS

CR-350 BIO-THANARITE

DISINFECTANT SANITIZER
 DEODORIZER

ACTIVE INGREDIENTS:

n-Alkyl (60% C₁₄, 30% C₁₆, 5% C₁₂, 5% C₁₈)
 dimethyl ammonium chlorides..... 5%
 n-Alkyl (68% C₁₂, 32% C₁₄) dimethyl
 ethylbenzyl ammonium chlorides 5%

INERT INGREDIENTS:..... 90%
 100%

CR-350 is a concentrate and should be diluted before using.

DANGER

KEEP OUT OF REACH OF CHILDREN.
 (See other precautionary and first aid statements on left side panel.)

"EPA Reg. No. 34224-8"

Manufactured By
CHEMRITE CORPORATION
 12600 SOUTH DAPHNE AVENUE
 HAWTHORNE, CALIFORNIA 90250

USE DIRECTIONS:

Household

For sanitizing and disinfecting floors, walls and inanimate hard surfaces. For Schools, Homes, Locker Rooms, Garbage Pails, Sink Tops, Corridors, Classrooms, Offices and shower Stalls. Apply solution with mop or cloth. Sanitize with 1 ounce **CR-350** to 4 gallons of water. Disinfect with 3 ounces (Name) to 5 gallons of water. Kills Staphylococcus aureus and Salmonella choleraesuis. For heavily soiled or contaminated areas, a pre-cleaning step is recommended.

Hospitals

For floors, walls and other hard surfaces. In Nursing Homes, Institutions, as well as Sink Tops, Garbage Pails, Telephones and Restrooms, use **CR-350** at 3½ ounces per 5 gallons of water. Apply with cloth, mop, or sponge. At this level **CR-350** will be effective against Pseudomonas aeruginosa PRD-10. For heavily soiled or contaminated areas, a pre-cleaning step is recommended.

Food Processing Equipment

For sanitization of previously cleaned food processing equipment and food utensils dilute 1 ounce **CR-350** per 4 gallons of water to provide 200 ppm of active quaternaries. At this level, **No Potable Water Rinse** is required.

Dairies

To sanitize dairy equipment such as tanks, lines, pails, and milk cans, first clean and rinse the equipment thoroughly. Then apply sanitizing solution containing 1 ounce **CR-350** to 4 gallons of water (200 ppm). At this level, **No Potable Water Rinse** is required. Follow recommendations of local Health Board.

Restaurant and Bar Rinse

Dishes, glassware, silverware, cooking utensils, wash with soap or synthetic detergent; rinse thoroughly and immerse in a sanitizing solution containing 1 ounce **CR-350** to 4 gallons of water. **No Terminal Potable Water Rinse Required.**

This product when used on environmental, inanimate, hard surfaces at 3½ ounces per 5 gallons of water is effective against Influenza A₂, Herpes Simplex, Adenovirus type 2, and Vaccinia Viruses.

CAUTION: Keep out of reach of children