

INSE
SANITIZER DEODORIZER

For Hospital, Institutional, Industrial,
 Other Farm and Home Use.

ENOL COEFFICIENTS

CC NO. 6538	79.2
NO. 6539	78.1
C ₁₆ , 10%)	10%
Chloride	2%
	88%
	100.0%

DANGER:
 OF CHILDREN. SEE FIRST
 CAUTIONS ON RIGHT PANEL.

Sold by
CHEMICAL CO.
 LE PHILA., PA. 19124

ACCEPTED
 JUN 9 1976
 Under the Federal Insecticide,
 Fungicide and Rodenticide Act,
 as amended, and the pesticide
 EPA Reg. 40 CFR 152.101
 registration no. 34282-6

USE DIRECTIONS FOR DISINFECTION

Disinfection in Hospitals, Nursing Homes and Other Health Care Institutions.

For disinfecting floors, walls, countertops, bathing areas, lavatories, bedframes, tables, chairs, garbage pails and other hard surfaces.

Add 4 1/4 oz. RINSE to 4 gallons water. Apply to previously cleaned hard surface with mop or cloth.

At this use-level, RINSE is effective against *Pseudomonas aeruginosa*.

Disinfectant in Institutions (i.e. Educational, Penal, Military), Industry, Schools and Homes

For disinfecting floors, walls, bedframes, countertops, tables, chairs, garbage pails, bathroom fixtures and other hard surfaces.

Add 3 oz. of RINSE to 5.5 gallons of water. Apply to previously cleaned hard surface with mop or cloth.

At 3oz./5.5 gallon use-level, RINSE is effective against *Staphylococcus aureus*, and *Salmonella choleraesuis*.

Disinfection of Barber Tools

Barber tools (such as combs, brushes, razors, and scissors) can be disinfected by immersing in a 1/2-oz./gallon solution of RINSE.

Disinfection of Poultry Equipment, Animal Quarters and Kennels.

Poultry brooders, watering founts, feeding equipment and other animal quarters (such as stalls and kennel areas) can be disinfected after thorough cleaning by applying a solution of 4 1/4 oz. RINSE to 4 gallons of water with a mop, cloth or brush. Small utensils should be immersed in this solution.

Prior to disinfection, all poultry, other animals and their feeds must be removed from the premises. This includes emptying all troughs, racks and other feeding and watering appliances. Remove all litter and droppings from floors, walls and other surfaces occupied or traversed by poultry or other animals.

After disinfection, ventilate buildings, coops and other closed spaces. Do not house poultry, or other animals, or employ equipment until treatment has been absorbed, set or dried.

All treated equipment that will contact feed or drinking water must be rinsed with potable water before reuse.

Rinse empty container thoroughly with water and discard it.

USE DIRECTIONS FOR SANITIZING

Sanitizing of Food Processing Equipment and other Hard Surfaces in Food Contact Locations.

For sanitizing food processing equipment, dairy equipment, food utensils, dishes, silverware, glasses, sink tops, countertops, refrigerated storage and display equipment and other hard surfaces.

Wash and rinse all articles thoroughly, then apply a solution of 1 oz. RINSE in 4 gallons of water.

Apply to sink tops, countertops, refrigerated storage and display equipment and other stationary hard surfaces by cloth or brush.

Dishes, silverware, glasses, cooking utensils and other similar size food processing equipment can be sanitized by immersion in a 1 oz./4 gallon dilution of RINSE.

No Potable water rinse is required.

At 1 oz./4 gallons RINSE has a hard water tolerance of 500PPM of hardness calculated as Ca CO₃ when evaluated by the AOAC Germicidal and Detergent Sanitizer Method against *Escherichia coli* and *Staphylococcus aureus*.

The udders, flanks, and teats of dairy cows can be sanitized by washing with a solution of 1 oz. RINSE in 4 gallons of warm water.

Use a fresh towel for each cow. Avoid contamination of sanitizing solution by dirt and soil. Do not dip used towel back into sanitizing solution. When solution becomes visibly dirty, discard and provide fresh solution.

FIRST AID:

In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution; or if these are not available, drink large quantities of water. Avoid alcohol. Call a physician immediately.

NOTE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression, and convulsion may be needed.

DANGER:

Keep Out of Reach of Children. Corrosive. Causes eye damage and skin irritation. Do not get in eyes, on skin, or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food.