DIRECTIONS

HOTELS-RESTAURANTS-TAVERNS SANITIZING AND DEODORIZING

- 1. Scrape and prewash all utensils and glassware
- 2 Thoroughly wash in solution of 1 ounce of Sod um Hypochiorite to 2 gallons water
- 3. Rinse in clear potable water
- 4 Sanitize in solution of 1 ounce Sodium Hypochlorite to 712 gallons water 130 PPM available chlorine. Immerse all utensils for a minimum of 2 minutes or contact time specified by governing sanitary code.
- 5 Place sanitized utensils on rack or drainboard to air dry

ENVIRONMENTAL PRECAUTIONS

This product is toxic to fish. Treated pool effluent should not be discharged where it will drain into lakes, streams, ponds, or public water. Do not contaminate water by cleaning equipment or disposal of wastes. Apply only as specified on label

WAF:NING: Keep out of reach of children Harmful it swallowed Strong Oxidant. May produce chemical burns. Do not get on skin or in eyes. Do not use ammonia or other bowl cleaners with this product. To do so will release hazardous gases

FIRST AID: In case of skin and eye contact, flush with plenty of water. If skin irritation persists, get medical attention. For eyes get prompt medical attention Internal—Give milk, water, or egg whites. Call Physician Immediately.

SODIUM HYPOCHLORITE SOLUTION

Active Ingredient: Sodium Hypochlorite 12.5% Inert Ingredient: 87.5%

Degrades with age. Use a test kit and increase dosage as necessary to obtain required level of available chlorine. See Directions on Right Panel.

DISINFECTANT BACTERICIDE DEODORANT FOR

FOOD PROCESSING PLANTS, DAIRY PLANTS, FARMS, HOTELS, RESTAURANTS, TAVERNS, SWIMMING POOLS.

WARNING

KEEP OUT OF REACH OF CHILDREN
SEE FIRST AID AND OTHER PRECAUTIONS ON LEFT PANEL

Produced By
The Chloramone Corporation
Delaware City, Delaware 19706

For ne each

Add 10 alent to with a tinue a 10,000 availab residua as dete at 7.2 to an ade

Sodiun Store i

Use Tes necess residua

Test Kil dealer.

ONDER THE PEDERAL INSPITICADE FUNCTION OF AND RODING THE ACT TO BE FOROMIC POISON RECISION.

RNS NG glassware ounce of

vater.

ium Hypo-Mavailable inimum of by govern-

drainboard

INS
of effluent
I drain into
er. Do not

nent or dis**d** on label.

ren. Harmkidant. May Do not get ot use amaners with vill release

ntact, flush in irritation ention. For l attention. er. or egg mediately.

SODIUM HYPOCHLORITE SOLUTION

Active Ingredient: Sodium Hypochlorite 12.5% Inert Ingredient: 87.5%

Degrades with age. Use a test kit and increase dosage as necessary to obtain required level of available chlorine. See Directions on Right Panel.

DISINFECTANT BACTERICIDE DEODORANT FOR

FOOD PROCESSING PLANTS, DAIRY PLANTS, FARMS, HOTELS, RESTAURANTS, TAVERNS, SWIMMING POOLS.

WARNING

KEEP OUT OF REACH OF CHILDREN
SEE FIRST AID AND OTHER PRECAUTIONS ON LEFT PANEL

Produced By
The Chloramone Corporation
Delaware City, Delaware 19706

DIRECTIONS

For newly filled pools or at start of each season:

Add 10 oz. 10,000 gallons of water (equivalent to 1.0 ppm available chlorine). Test with a reliable chlorine test kit and continue adding in smaller amounts (1 oz. 10,000 gallons equivalent to 0.1 ppm available chlorine) if necessary until a residual of 0.6 to 1.0 ppm is established as determined by the test kit. Keep the pH at 7.2 to 7.6 at all times as determined by an adequate pH test kit.

Sodium Hypochlorite degrades with age. Store in cool, dark place.

Use Test Kit often and increase dosage as necessary to maintain required level of residual chlorine.

Tost l'ata available from your pool supply dealer.