


**PRODUCT INFORMATION**

PIRIMOR 50 W is a quick-acting selective insecticide for the control of aphids on potatoes. When applied to foliage, PIRIMOR 50 W controls aphids through contact action, shows fumigant action against aphids and may give a high

degree of control within a few hours. It has also been shown to be effective against aphids tolerant to organophosphorus insecticides. PIRIMOR 50 W is relatively inactive against predator and non-target beneficial insects.



**Aphid Control Spray**

**Pirimor** 50 W

**Pirimor**

**DIRECTIONS FOR USE**

PIRIMOR 50 W will control green peach aphid, potato aphid and buckthorn aphid. By ground sprayer, apply 4 to 8 oz. PIRIMOR 50 W per acre in 30 to 150 gallons of water. **SPRAY TO WET ALL FOLIAGE, MAKING SURE LOWER LEAVES ARE WETTED.** Spray in a 7 to 10-day program or as needed. The higher dose rate may be required under conditions of heavy infestation. For the

most effective and economic control, begin to apply PIRIMOR 50 W as soon as aphids appear. **Compatibility:** Although no phytotoxicity has been observed when PIRIMOR 50 W was combined with commonly used pesticides, additional compatibility information must be obtained before a recommendation for such use can be made.

**NOTICE TO BUYER:** Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in accordance with directions under normal conditions of use. This warranty does not extend to the use of this product contrary to label instructions, or under abnormal use conditions, or under conditions not reasonably foreseeable to Seller, and Buyer assumes the risk of any such use. Seller disclaims all other warranties, expressed or implied, including any warranty of FITNESS OR MERCHANTABILITY. Seller shall not be liable for consequential, special or indirect damages resulting from the use or handling of this product and Seller's sole liability and Buyer's exclusive remedy shall be limited to the refund of the purchase price.

Distributed by:



**ICI United States Inc.**  
Agricultural Chemicals Division  
Wilmington, Delaware 19897

Contents  
manufactured in  
Great Britain

EPA Registration No. 10182-7

Active Ingredient: 2-(dimethylamino)-5,6-dimethyl-4-pyrimidinyl dimethylcarbamate . . . . .	<b>50%</b>
Inert ingredients: . . . . .	<b>50%</b>
Total . . . . .	<b>100%</b>

**NET CONTENTS: 2½ LBS.**

U.S. Patent Nos. 3493574 and 3577543

PIRIMOR is a registered trademark of Imperial Chemical Industries Limited.

EPA Est. No. 09779-NC-01

**WARNING**

**KEEP OUT OF REACH OF CHILDREN**

MAY BE FATAL IF SWALLOWED. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONS AND DIRECTIONS AND IN CONFORMITY WITH FEDERAL AND STATE REGULATIONS.

Avoid unnecessary exposure to spray mist and wash thoroughly after handling.

**FIRST AID:** If illness occurs following exposure to PIRIMOR 50 W, call a physician.

If swallowed, induce vomiting. Repeat until vomit is clear. Have patient lie down and keep quiet. Call a physician.

**NOTE TO PHYSICIAN:** Prolonged exposure may cause cholinesterase inhibition. Atropine is antidotal.

Avoid contamination of food, feed, water supplies, lakes, streams and ponds.

Do not reuse container. Destroy when empty.



**ICI United States Inc.**  
Agricultural Chemicals Division  
Wilmington, Delaware 19897



# 10-35-5

# SYSTEMIC-T

## fertilizer plus insecticide

FOR USE ON BEANS, POTATOES, CORN AND SUGAR BEETS TO CONTROL THE INSECTS SHOWN UNDER DIRECTIONS.

Contains 0.2% THIMET®  
Soil Insecticide

### Statement of Ingredients


**Active Insecticide Ingredients:**  
 Phorate (O, O-diethyl S-( ethylthio) methyl) phosphorodithioate ..... 0.20%  
 Insecticidally Inert Ingredients ..... 99.80%

### Fertilizer Ingredients

Total Nitrogen (Nitrogen (N) ..... 10%  
 Available Phos. Acid (P<sub>2</sub>O<sub>5</sub>) ..... 35%  
 Water Sol. Potash (K<sub>2</sub>O) ..... 5%

## DANGER!

### Keep Out of Reach of Children



Not for use or storage in or around the home. Poisoning may occur as a result of exposure. May be fatal if swallowed or inhaled. Rapidly absorbed through the skin. Do not breathe dust. Do not get in eyes or on skin. Wash all contaminated clothing with soap and hot water before reuse. Wash thoroughly after handling, and before eating or smoking. Do not contaminate feed or food. Do not reuse empty container. Burn empty bags and stay away from smoke or fumes. This product is toxic to fish and wildlife. Keep out of lakes, streams or ponds. Birds feeding on treated areas may be killed. Apply this product only as specified on this label. Antidote:— Atropine.

### Directions

Thimet is an effective soil and systemic insecticide for early season insect control, which has been combined with fertilizer for efficient and uniform application. Application should be made at planting time, in the row to the side of the seed. Do not apply directly to the seed.

### Recommendations for Use on Crops

Crop	Insects	Rate-lbs./acre	Application
<u>POTATOES</u>	Aphids, leafhoppers, leafminers, psyllids, wireworms, flea beetle larve and reduction of flea beetle adults	1000 to 1500	Granules may be banded on each side of the row at planting time. Use 500 lbs. in sandy or light soils, 750 lbs. in heavy silt, clay loam or muck soils (38" row spacing). Do not apply more than once (1) per season.
<u>CORN</u> At planting time	Corn rootworms	252 to 500	At planting time, in the furrow 1/2" to 1" above seeds on each side of the seed row. Do not place in direct contact with seed. Use 500 lb. rate in areas having high rootworm populations.
<u>SUGAR BEETS</u>	Sugar beet root maggot, leafhoppers, mites, aphids, pea leafminers	500	Apply in 7" band over the row at planting time. Do not place in contact with seed. Make no application later than 30 days before harvest. Do not feed beet tops or silage to dairy cattle.
<u>BEANS</u>	Leafhoppers, aphids, lygus bugs, thrips, mites and seedcorn maggots	500 to 740	Distribute the granules evenly in the row to the side of the seed at planting time. Apply 30 to 40 lbs. per 1000 feet of row. Do not place in direct contact with the seed. Do not feed foliage within 60 days of treatment.

®Trademark of American Cyanamid Company

U.S.D.A. REG. No. 10501-3



Produced by  
**FARM BUREAU CO-OPERATIVE**  
 Antigo, Wisconsin 54409

Handwritten notes:  
 10501-3

## ENVIRONMENTAL HAZARDS

**PRECAUTIONS IN USING:** "This product is toxic to fish and wildlife," "Birds and other wildlife in treated areas may be killed," "Keep out of lakes, streams, or ponds," "Do not apply where runoff is likely to occur," "Do not apply when weather conditions favor drift from areas treated," "Do not contaminate water by cleaning of equipment, or disposal of wastes," "Apply this product only as specified on this label."

**BEE CAUTION:** This product is toxic to bees and should not be applied when bees are actively visiting the area.

### RESTRICTIONS

**USE PRECAUTIONS:** Do not apply to or allow to drift onto varieties of sorghum which are sensitive to phosphates. To do so may cause burning of the sorghum or milo. Do not use on other crops used for food or forage. Use only according to label directions. Application at rates above those shown may result in illegal crop residues. Do not treat food crops grown in the greenhouse.

### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

# TIFCHEM CUTWORM & CRICKET BAIT

Contains 5% **DYLOX**

#### ACTIVE INGREDIENT:

Dimethyl (2,2,2-trichloro-1-hydroxyethyl)  
Phosphonate† ..... 5.00%

INERT INGREDIENTS ..... 95.00%

Total ..... 100.00%

DYLOX is a Reg. TM of the Parent Company of  
Farbenfabriken Bayer GmbH, Leverkusen.

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

May be harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with skin. Do not breathe dust. Wash thoroughly with soap and warm water after handling. Wash contaminated clothing with soap and hot water before re-use. Do not contaminate feed or food. If misuse results in

Lima Beans

Cabbage  
Cauliflower  
Peppers

Collards  
& Lettuce  
Tomatoes

Crop  
Corn

Cowpeas  
(black  
crow  
southern)

Tobacco

Peas