



MALATHION GRAIN PROTECTOR

Reg # 34704-8

PM-14

PRECAUTIONARY STATEMENTS HARMFUL TO HUMANS AND DOMESTIC ANIMALS **CAUTION**

182

Harmful if Swallowed. Avoid contact with skin. Wash thoroughly after using. Avoid breathing of dust. Avoid contamination of feed and foodstuff. Keep out of reach of children.

STATEMENT OF PRACTICAL TREATMENT

If Swallowed—Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching the back of throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.
If Inhaled—Remove victim to fresh air. Apply artificial respiration if indicated.
If On Skin—Remove contaminated clothing and wash affected areas with soap and water.
If In Eyes—Flush eyes with plenty of water. Call a physician if irritation persists.
Note to Physician: This product may cause cholinesterase inhibition. Atropine is antidotal. 2 PAM may be effective as an adjunct to atropine.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish, aquatic invertebrates, and aquatic life stages of amphibians. Do not apply directly to water or wetlands (i.e., swamps, bogs, marshes, and potholes). Drift and runoff may be hazardous to aquatic organisms in areas near the application site. Do not contaminate water when disposing of equipment washwaters.
This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

IMPORTANT

Use of this product, as directed, will be within the tolerance of 8 parts per million for MALATHION. Treated grain is not subject to downgrading and can be milled into flour or feed using normal production methods.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Do not store under conditions which might adversely affect the container or its ability to function properly. Such conditions include, but are not limited to, positioning of the container in storage, storage temperature, potential for crushing or damage due to stacking, and penetration of moisture.
STORAGE: Store in safe manner. Store in original container only. Store in cool, dry place. Reduce stacking height where local conditions, such as humidity or pallet overhang, can effect package strength. Personnel should use clothing and equipment consistent with good pesticide handling.
PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.
CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

ACTIVE INGREDIENT;

Malathion*	
*O,O-dimethyl phosphorodithioate of diethyl mercaptosuccinate.....	2.00%
INERT INGREDIENT.....	98.00%
TOTAL	100.00%

KEEP OUT OF REACH OF CHILDREN **CAUTION**

See side panel for additional precautionary statements.

PROTECTS AGAINST DAMAGE from Confused Flour Beetle, Rice Weevil, Granary Weevil, Saw-toothed Grain Beetle, Flat Grain Beetle, Red Flour Beetle, Rusty Grain Beetle, Lesser Grain Borer, Indian Meal Moth.

FOR USE ON STORED GRAINS:

Wheat, Corn, Oats, Rice, Barley, Grain Sorghum, Field and Garden Seeds.

NOTICE

Buyer assumes all risk of use, storage, or handling of this product when not in strict accordance with the directions given herewith.

NET WEIGHT 50 LBS.

EPA REG. NO. 34704-8

BEST AVAILABLE COPY

PLATTE CHEMICAL COMPANY, INC.
130 S. MAIN STREET
FREMONT, NEBRASKA 68025

ACCEPTED
JAN 23 1989
EPA Reg. No. 34704-8

292

SHELLED CORN AND SMALL GRAINS - The most convenient way to apply MALATHION GRAIN PROTECTOR is to treat it when it is in the truck or wagon prior to binning. Determine the amount in the truck or wagon in this manner:

Shelled Corn or Small Grains - multiply the length by width by depth (all in feet) and multiply this total by 0.8 to get the number of bushels.

Measure out the correct amount of MALATHION GRAIN PROTECTOR (30 lbs. per 1,000 bushels).

Spread the PROTECTOR evenly over the surface of the grain in the truck or wagon. Cut the PROTECTOR into the grain with a shovel. The PROTECTOR will become mixed with grain as it is dumped from the truck into the auger hopper. Further mixing will occur when the grain is augered or blown into the bin. Mixing is completed as the PROTECTOR and grain roll down the cone shaped pile in the bin.

Apply MALATHION GRAIN PROTECTOR to the grain at the rate of 30 pounds per 1,000 bushels as the grain is being loaded or turned into final storage.

Field and garden seeds can be protected against grain pests with a dosage of 30 pounds of MALATHION GRAIN PROTECTOR per 1,000 bushels of seed.

To protect stored grains from attack by Indian Meal Moth, apply MALATHION GRAIN PROTECTOR to the surface of clean or infested grain at the rate of 15 pounds per 1,000 square feet of grain surface area. Apply the dust evenly over the surface of the grain. Apply immediately after grain is loaded into storage and repeat if necessary.

BEST AVAILABLE COPY