

R23 # 34704-10 PM 12

11/12



RESTRICTED USE PESTICIDE

FOR RETAIL SALE TO AND USE ONLY
BY CERTIFIED APPLICATORS OR PER-
SONS UNDER THEIR DIRECT SUPER-
VISION AND ONLY FOR THOSE USES
COVERED BY THE CERTIFIED APPLI-
CATOR'S CERTIFICATION.

METHYL PARATHION 4E

Emulsifiable Liquid - An Agricultural Insecticide

ACTIVE INGREDIENTS:

*O,O-dimethyl O-p-nitrophenyl phosphorothioate	43.2%
Related Products	2.3%
Xylene Range Aromatic Solvent	47.8%

INERT INGREDIENTS:

TOTAL 100.0%

*Methyl Parathion Contains 4 lbs. Methyl Parathion per Gallon.

KEEP OUT OF REACH OF CHILDREN

**DANGER
PELIGRO**



POISON

PRECAUCION AL USARIO: Si usted no lee ingles, no use este producto hasta
que le etiqueta haya sido explicado ampliamente.



**CAN KILL YOU
IF SWALLOWED**

This product can kill you if swallowed
even in small amounts: spray mist or
dust may be fatal if swallowed.



**CAN KILL YOU
BY SKIN CONTACT**

This product can kill you if touched
by hands or spilled or splashed on
skin, in eyes, or on clothing (liquid
goes through clothes).



**CAN KILL YOU
IF BREATHED**

This product can kill you if vapors,
spray mist or dust are breathed.

See Side Panels For Antidote and Precautions

and oral warnings must include the following information:

DANGER

... or field description treated with methyl parathion or insert date of application.) Do not enter without appropriate protective clothing. In case of skin or eye exposure: Call a doctor (physician), clinic or hospital immediately. Explain that the person has been exposed to methyl parathion and describe his condition. For further information, see the treatment of FIRST AID TREATMENT portion of the pesticide label.

STORAGE AND DISPOSAL

PRECAUTIONS: Do not contaminate water, food, or feed by storage or disposal. Do not store under conditions which might adversely affect the container or its ability to function properly.

STORAGE: Do not store below temperature of 55°F. Store in safe manner. Store in original container only. Keep container tightly closed when not in use. Reduce stacking height where local conditions can affect package strength. Persons should use clothing and equipment listed under "PRECAUTIONARY STATEMENT" when handling open containers.

SPILLED MATERIAL: Block or dike to prevent spreading of spill. Cover with absorbent material such as lime, clay or sand. Collect material and place in a disposable container. Wash area with strong lye solution, absorb and place into a disposable container.

HAZARDOUS WASTE: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mix, or empty containers is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

DIRECTIONS

Be sure to read the precautionary statements before using!

This product is designed for application after dilution with water and for use by trained operators using airplane or power ground equipment. The hazards and precautions for handling the product in this container are equally applicable to it after dilution with water for spray application. Add the concentrate to the spray tank while filling with water, and mix thoroughly either by means of a tank agitator or pump by-pass. For best results, thoroughly cover all surfaces to be treated with spray. Rates of application given below should not be exceeded. Never apply later than indicated to assure residue levels at harvest are below tolerances.

Consult the State Agricultural Extension Service or Experiment Station for specific recommendations regarding application, storage and timing of sprays. For application by ground equipment, add the desired amount of concentrate to sufficient water to apply at least 3 gallons of water per acre. For application by aircraft, add the amount of concentrate desired per acre to 1 to 3 gallons of water consistent with crop growth and good coverage. Greater quantities of water may be required to give sufficient coverage of orchard trees.

NOTICE

Because Platte Chemical Co., Inc. has no control over storage, handling, and conditions of use, which are of critical importance, Platte Chemical Co., Inc. makes no representation or warranty, either express or implied, for results or residues greater than any tolerance which may be established by appropriate governmental agencies, due to misuse, improper handling or storage of this material. Nor does Platte Chemical Co., Inc. assume any responsibility for injury to persons, crops, animals, soil or property arising out of misuse, improper handling or storage of this material.

FORMULATED FOR

PLATTE CHEMICAL COMPANY, INC.

154 SO. MAIN STREET, FREMONT, NEBRASKA 68025

NOT FOR HOME USE

Complete Directions for Use

EPA REG. NO. 34704-10

Use only according to these label instructions

CLEAN CROP is a Registered T.M.

of United Agri Products, Inc.

ACCEPTED

JUL 26 1985

Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 34704-10

NOTICE

Because Platte Chemical Co., Inc. has no control over storage, handling, and conditions of use, which are of critical importance, Platte Chemical Co., Inc. makes no representation or warranty, either express or implied, for results or residues greater than any tolerance which may be established by appropriate governmental agencies due to misuse, improper handling or storage of this material. Nor does Platte Chemical Co., Inc. assume any responsibility for injury to persons, crops, animals, soil or property arising out of misuse, or improper handling or storage of this material.

ATTENTION

Do not use in any manner other than recommended on this label.

To avoid excessive residues of methyl parathion on food or forage crops, always observe the statements found under "Directions for Use," limiting the time before harvest when methyl parathion may be applied.

ANTIDOTE	8
FIRST AID	8
GENERAL DIRECTIONS	9-11
HAZARDS, SAFEGUARDS	5-9
INGREDIENTS	5
NOTICE	2
POISON SIGNS	7

CEREAL INSECTS

BARLEY	11
CORN	11
OATS	11
RICE	11
RYE	11
WHEAT	11

FIELD AND FORAGE CROPS INSECTS

ALFALFA	12
CLOVER	12
COTTON	12
GRASS (Forage)	12
HOPS	12
SOYBEANS	12
SUGAR BEETS	13
TOBACCO	13
VETCH	13

FRUIT INSECTS

APPLES	13
APRICOTS	14
CHERRIES	14
GOOSEBERRIES	14
GRAPES	14
PEACHES	14
PEARS	14
PLUMS	14
PRUNES	14
STRAWBERRIES	14

Contents 4

OIL SEED CROP INSECTS	
SAFFLOWER	14
SUNFLOWER	14

VEGETABLE INSECTS

ARTICHOKE	15
BEANS Dry and Green	16
BEETS (Red)	16
BROCCOLI	15
BRUSSELS SPROUTS	15
CABBAGE	15
CARROTS	18
CAULIFLOWER	15
CELERY	16
COLLARDS	16
CUCUMBERS	16
KALE	16
KOHLRABI	15
LETTUCE	17
MUSTARD	16
ONIONS	15
PEAS	16
PEPPERS	18
POTATOES	15
RUTABAGAS	15
SPINACH	17
SWEET POTATOES	15
TOMATOES	16
TURNIPS	17

FOREST, NONAGRICULTURAL INSECTS

ORNAMENTALS	17
PINE FOREST	17
FOREST, NONAGRI	17

Hazards and Ingredients 5

KEEP OUT OF REACH
OF CHILDREN

DANGER
POISON  **PELIGRO**

PRECAUCION AL USARIO: Si usted no lee ingles, no use este producto hasta que le etiqueta haya sido explicado ampliamente.

See antidote statments and other required warning statements through page 10.

ACTIVE INGREDIENTS:

*O,O-dimethyl O-p-nitrophenyl phosphorothioate	43.2%
Related Products	2.3%
Xylene Range Aromatic Solvent	47.8%
INERT INGREDIENTS:	8.7%

TOTAL 100.0%

*Methyl Parathion
Contains 4 lbs Methyl Parathion per Gallon

BEST AVAILABLE COPY

Hazards

DANGER—POISON—PRECAUTIONS **KEEP OUT OF REACH OF CHILDREN**

PELIGRO DANGER POISON



PRECAUCION AL USARIO: Este producto es muy venenoso y puede causar la muerte si se lo ingiere o si se inhala. Evite el contacto con la piel y los ojos.



CAN KILL YOU IF SWALLOWED

This product can kill you if swallowed even a small amount. Spit it out if it gets in your mouth. Do not swallow it.



CAN KILL YOU BY SKIN CONTACT

This product can kill you if it touches your skin or gets in your eyes or on your clothing. Wash it off immediately with soap and water.



CAN KILL YOU IF BREATHED

This product can kill you if you breathe it in. Do not breathe the vapors.

PRECAUTIONARY STATEMENTS **HAZARDS TO HUMANS & DOMESTIC ANIMALS**



DANGER

USE ONLY WHEN WEARING THE FOLLOWING PROTECTIVE EQUIPMENT AND CLOTHING:

- (1) Wear water-proof pants, coat, hat, rubber boots and gloves.
- (2) Wear safety goggles.
- (3) Wear a pesticide-resistant suit, hood and gloves.
- (4) Wear a respirator approved by the Mine Safety and Health Administration, the U.S. Department of Labor, and the National Institute for Occupational Safety and Health.
- (5) Wear a respirator approved by the Mine Safety and Health Administration, the U.S. Department of Labor, and the National Institute for Occupational Safety and Health.

Hazards

WORKER SAFETY RULES

Keep all unprotected persons and children away from treated area or where there is danger of drift.

TO PREVENT PERSONAL INJURY AND POSSIBLE FATALITIES keep all unprotected person and animals out of treated areas for 48 hours.

Do not rub eyes or mouth with hands. If you feel sick in any way, **STOP** work and get help right away. Call a doctor (physician), clinic or hospital immediately. Explain that the victim has been exposed to parathion and describe his condition. After first aid is given (see First Aid Treatment Section) and if a doctor cannot come, take victim to clinic or hospital.

IMPORTANT! Before removing gloves, wash them with soap and water. Always wash hands, face and arms with soap and water before smoking, eating or drinking.

AFTER WORK, take off all work clothes and shoes. Shower, using soap and water. Wash and clean clothes when leaving job. Do not wear contaminated clothing. Wash protective clothing and protective equipment with soap and water after each use. Respirator should be cleaned and later replaced according to instructions included with respirator.

POISON SIGNS (Symptoms)

Methyl Parathion is a very dangerous poison. It rapidly enters the body on contact with the skin, clothes and eyes. Clothing wet with this material must be removed immediately. Exposed persons must receive prompt medical treatment or they may die.

Some of the signs and symptoms of poisoning are: Headache, nausea, vomiting, cramps, weakness, blurred vision, pinpoint pupils, tightness in chest, labored breathing, nervousness, sweating, watering of eyes, drooling or frothing of mouth and nose, muscle spasms and coma.

BEST AVAILABLE COPY

FIRST AID TREATMENT

Call a doctor (physician) clinic or hospital immediately. Explain that the victim has been exposed to methyl parathion and describe his condition.

If breathing has stopped, start artificial respiration immediately and maintain until doctor sees victim.

If swallowed — Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

In case of contact, immediately flush eyes or skin with plenty of water while removing contaminated clothing and shoes. See doctor immediately.

NOTE TO PHYSICIAN

Antidote — administer atropine sulfate in large doses. TWO to FOUR mg. intravenously or intramuscularly as soon as cyanosis is overcome. Repeat at 5 to 10 minute intervals until signs of atropinization appear. 2-PAM chloride is also antidotal and may be administered in conjunction with atropine. DO NOT GIVE MORPHINE OR TRANQUILIZERS. Methyl parathion is a strong cholinesterase inhibitor affecting the central and peripheral nervous systems and producing cardiac and respiratory depression.

At first sign of pulmonary edema, the patient should be given supplemental oxygen and treated symptomatically. Continued absorption of the poison may occur and further doses may be required after some improvement. VERY CLOSE SUPERVISION OF THE PATIENT IS INDICATED FOR AT LEAST 48 HOURS.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply under weather conditions favoring drift from areas treated. Do not contaminate water by cleaning of equipment or disposal. Lake trout, Striped and crab may be killed at application rates recommended on this label. Do not apply where these are important resources. Apply this product only as specified on this label.

This product is highly toxic to humans exposed to direct treatment or residues on crops. Protective information may be obtained from your Agricultural Extension Service.

PHYSICAL & CHEMICAL HAZARDS

WARNING: COMBUSTIBLE

KEEP AWAY FROM HEAT AND OPEN FLAME

DO NOT USE IN UNDILUTED FORM

NOT FOR USE OR STORAGE IN OR AROUND THE HOME

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

RE-ENTRY

Do not reenter treated areas for 48 hours unless appropriate protective clothing is worn. Because certain states may require more restrictive reentry intervals for various applications with this product, consult your State Department of Agriculture for further information. Do not apply this product near to areas where as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. When oral warnings are given, warnings shall be given in a language customarily understood by workers.

Written or oral warnings must include the following information:

DANGER

(Insert area or field description) treated with methyl parathion on (insert date of application). Do not enter within (insert appropriate protective clothing) for 48 hours. In case of accidental exposure: Call doctor (physician), clinic or hospital immediately. Explain that the victim has been exposed to methyl parathion and describe his condition. For further information see the statement of FIRST AID TREATMENT portion of the product label.

BEST AVAILABLE COPY

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Do not store under conditions which might adversely affect the container or its ability to function properly.

STORAGE: Do not store below temperature of 0°F. Store in safe manner. Store in original container only. Keep container tightly closed when not in use. Reduce stacking height where local conditions can affect package strength. Personnel should use clothing and equipment listed under "PRECAUTIONARY STATEMENT" when handling open containers.

SPILLED MATERIAL: Block or dike to prevent further spill. Cover with absorbent material such as sawdust, straw, or sand. Scoop and sweep into a disposable container. Wash area with strong lye solution, absorb and place into a disposable container.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities.

DIRECTIONS

Be sure to read the precautionary statements before using! This product is designed for application after dilution with water and for use by trained operators using airplane or power ground equipment. The hazards and precautions for handling the product in this container are equally applicable to it after dilution with water for spray application. Add the concentrate to the spray tank while filling with water, and mix thoroughly either by means of a tank agitator or pump by-pass. For best results, thoroughly cover all surfaces to be treated with spray. Rates of application

given below should not be exceeded. Never apply later than indicated to assure residue levels at harvest are below tolerances established by the Food and Drug Administration.

Consult the State Agricultural Experiment Station or Experiment Station for specific recommendations regarding application, dosage and timing of sprays. For application by ground equipment, add the desired amount of concentrate to sufficient water to apply at least 3 gallons of water per acre. For application by aircraft, add the amount of concentrate desired per acre to 1 to 3 gallons of water consistent with crop growth and good coverage. Greater quantities of water may be required to give sufficient coverage of orchard trees.

CEREAL INSECTS

BARLEY, OATS, RYE AND WHEAT: For barley thrips, use 1/2 to 1 1/2 pints per acre. For aphids, black grass bugs, stink bugs and winter grain beetles, use 1/2 to 1 1/2 pints per acre. For climbing cutworms and grasshoppers, use 1 to 1 1/2 pints per acre. For armyworms and leathoppers, use 1/2 to 1 pint per acre. For chinch bugs, false chinch bugs and bay-leaved bugs, use 1 1/2 pints per acre. For western or brown wheat beetles and grasshoppers, use 1 pint per acre. Do not use more than 1/2 pint per acre within 15 days of harvest.

CORN: For control of armyworms, climbing cutworms, corn leaf aphids, corn rootworm adults, for armyworms and stink bugs, use 1/2 pint per acre. For grasshoppers, use 1 pint per acre. Do not apply within 12 days of harvest.

RICE: For rice stink bugs, use 1 to 1 1/2 pints per acre. For grasshoppers, use 1 pint per acre. Do not apply within 15 days of harvest or within 5 days after propane has been applied. Shrimp, crabs and crayfish may be killed. Do not apply where these are important resources.

BEST AVAILABLE COPY

FIELD AND FORAGE CROPS

ALFALFA AND CLOVER — For control of alfalfa weevil larvae, aphids (including spotted alfalfa aphids) and climbing cutworms and grasshoppers, use $\frac{1}{2}$ to 1 pint per acre. For alfalfa caterpillars, alfalfa adult weevils, armyworms, clover leaf weevils and webworms, use 1 pint per acre. For Egyptian alfalfa weevils, leafhoppers, Lygus bugs and spider mites, use 1 to 2 pints per acre. Do not treat closer than 15 days before harvest, cutting, or grazing. In California and Nevada do not use more than $\frac{1}{2}$ pint per acre.

COTTON — For control of thrips, use $\frac{1}{4}$ to $\frac{1}{2}$ pint per acre. For cotton leafworms use $\frac{1}{2}$ to 8 pints per acre. For grasshoppers and fall armyworms, use $\frac{1}{2}$ to 6 pints per acre. For boll weevils, aphids and garden webworms, use $\frac{1}{2}$ to 8 pints per acre. For false chinch bugs use 2 pints per acre. For cutworms, use 1 to 2 pints per acre. For bollworms, saltmarsh caterpillars, armyworms, cotton leaf perforators and stink bugs, use $2\frac{1}{2}$ to 8 pints per acre. Applications should be made at 4 to 5 day intervals until control is obtained. At above dosages, application may be made up to the day before harvest, if harvest will be with mechanical pickers.

DANGER — When necessary to enter treated cotton fields within 48 hours after application, protective clothing should be worn. Do not apply within 7 days of handpicking cotton.

GRASS (Forage) — For range caterpillar, use 1 pint per acre. For control of armyworms, crested wheat bugs, false chinch bugs, grasshoppers and leafhoppers, use $1\frac{1}{2}$ pints per acre. Do not apply within 15 days of harvest or grazing.

HOPS — For control of aphids, spider mites, armyworms, and leafhoppers, use 1 quart per acre. Do not treat closer than 15 days before harvest.

SOYBEANS — For control of climbing cutworms, three-cornered alfalfaoppers, garden webworms, aphids, blister beetles, Mexican bean beetles, stink bugs, two-spotted mites and velvet bean caterpillars, use $\frac{1}{4}$ to 2 pints per acre. For bollworms, fall armyworms and green cloverworms use 1 quart per acre. Do not apply closer than 20 days before harvest or grazing.

SUGAR BEETS — For armyworms, cutworms, fall armyworms, flea beetles, grasshoppers, leafhoppers, mites, sugar beet webworms, use $\frac{1}{2}$ to $\frac{3}{4}$ pint per acre. Do not apply within 20 days of harvest, 60 days if tops are to be fed to animals.

TOBACCO — For control of green peach aphids, use $\frac{1}{2}$ pint per acre. For control of surface feeding or climbing cutworms, use $\frac{1}{2}$ to 1 pint per acre.

DANGER — When necessary to enter treated tobacco fields within 48 hours after application, protective clothing should be worn. Do not apply within 5 days of pruning tobacco or within 15 days of cutting tobacco. Avoid contact with plant juices when pruning or cutting tobacco.

VETCH — Alfalfa weevil (larvae) - Apply $\frac{1}{2}$ to 1 pint per acre. Apply when 75% of terminals show feeding, when buds and shoots appear ragged or when the larvae counts reach 25 per sweep. Aphids, leafhoppers - Apply $\frac{1}{2}$ to 1 pint per acre. Climbing cutworms, Lygus bugs, Vetch beetle - 1 pint per acre. Armyworms - $\frac{1}{2}$ to $2\frac{1}{2}$ pints per acre. DO NOT apply within 15 days of harvest through 2 pints. DO NOT apply within 20 days of harvest from 2 pints through $2\frac{1}{2}$ pints.

FRUIT

APPLY AT THE RATES INDICATED FOR THE FOLLOWING CROPS. MAKE NO APPLICATION CLOSER THAN 14 DAYS BEFORE HARVEST. APPLY AS FREQUENTLY AS NEEDED TO CONTROL INSECTS.

APPLES — For control of aphids, including woolly apple aphids, rose apple aphid, green aphid, codling moths, plum curculio, European red mite, scales and red-banded leaf rollers, use $\frac{1}{2}$ pint to 1 quart per 100 gallons of water. For codling moths, plum curculio and red-banded leaf rollers, apply at petal fall and 3 to 4 applications, 8 to 14 days thereafter to maintain control. Do not use more than 6 quarts of this product per acre.

BEST AVAILABLE COPY

APRICOTS (14) - For Woolly apple aphid, Rosy apple aphid, green aphid, aphids, European Red Mite, use 4 oz /100 gals water per acre. Apply twice at 5 day intervals.

CHERRIES (14) - For Black cherry aphid, European Red Mite, use 4 oz /100 gals. water per acre. Apply at 7 to 10 day intervals.

GOOSEBERRIES (15) - For Aphids use 4 oz /100 gals water per acre.

GRAPES - For control of aphids, grape leaf folders and grape leafhoppers, use $\frac{1}{2}$ pint to 1 quart per 100 gallons of water. Do not use more than 1 $\frac{1}{2}$ pints of this product per acre.

PEACHES, PLUMS AND PRUNES - For control of aphids, including woolly apple aphid, rosy apple aphid, green aphid, green peach aphid, mealy plum aphid, peach tree borers, plum curculio, Oriental fruit moths, and European red mite, use $\frac{1}{2}$ pint to 1 quart per 100 gallons of water. Spray tree trunks at time of moth flight to control peach tree borers. For plum curculio and Oriental fruit moths, apply at petal fall or shuck split and 1 to 4 applications, 8 to 14 days thereafter, to maintain control. Do not use more than 1 gallon of this product per acre.

PEARS - For woolly apple aphids, rosy apple aphids, green aphids, aphids, and European red mites, use 4 oz per 100 gallons of water per acre. Apply twice at 5 day intervals.

STRAWBERRIES - For control of aphids, use 1 to 1 $\frac{1}{2}$ pints per acre.

OIL SEED CROPS

SAFFLOWER - For aphids and lygus bugs, use 1 pint per acre before flowering. Do not apply after flowering.

SUNFLOWERS (30) - For sunflower moths, sunflower maggot and sunflower seed weevil, use 2 pints per acre. Do not apply more than 3 times at 5-day intervals per season. Do not feed seeds to birds. Do not apply within 30 days of harvest.

VEGETABLES

APPLY AT THE RATES INDICATED FOR THE FOLLOWING CROPS. MAKE NO APPLICATIONS CLOSER THAN 5 DAYS BEFORE HARVEST.

POTATOES - For control of aphids, armyworms, large chinch bugs, flea beetles, leafhoppers, mites and shield bugs, use 1 $\frac{1}{2}$ quarts per acre.

SWEET POTATOES - For control of aphids, armyworms, surface feeding and climbing cutworms, flea beetles, leafhoppers and mites, use $\frac{1}{2}$ quart per acre.

APPLY AT THE RATES INDICATED FOR THE FOLLOWING CROPS. MAKE NO APPLICATIONS CLOSER THAN 7 DAYS BEFORE HARVEST.

ARTICHOKE - For control of aphids, armyworms, surface feeding and climbing cutworms, flea beetles, leafhoppers, mites and artichoke plume moths, use 1 quart per acre.

BROCCOLI - For control of aphids, armyworms, flea beetles, leafhoppers and mites, use 1 to 3 pints per acre. For stink bugs, use 2 to 3 pints per acre. Rates above 1 pint should not be applied closer than 21 days before harvest.

BRUSSELS SPROUTS - For control of aphids, armyworms, flea beetles, leafhoppers and mites, use 1 to 3 pints per acre. For stink bugs, use 2 to 3 pints per acre. Rates above 1 pint should not be applied closer than 21 days before harvest.

CAULIFLOWER - For control of aphids, armyworms, flea beetles, leafhoppers and mites, use 1 to 3 pints per acre. For stink bugs, use 2 to 3 pints per acre. Rates above 1 pint should not be applied closer than 21 days before harvest.

KOHLRABI - For control of aphids, armyworms, flea beetles, leafhoppers, and mites, use 1 to 3 pints per acre. For stink bugs, use 2 to 3 pints per acre. Rates above 1 pint should not be applied closer than 21 days before harvest.

RUTABAGAS - For control of aphids, armyworms, flea beetles, leafhoppers and mites, use 1 to 3 pints per acre. Rates above 1 pint should not be applied closer than 21 days before harvest.

APPLY AT THE RATES INDICATED FOR THE FOLLOWING CROPS. MAKE NO APPLICATIONS CLOSER THAN 10 DAYS BEFORE HARVEST.

CABBAGE - For control of aphids, armyworms, flea beetles, leafhoppers and mites, use 1 to 3 pints per acre. For stink bugs, use 2 to 3 pints per acre. Rates above 1 pint should not be applied closer than 21 days before harvest.

BEST AVAILABLE COPY

COLLARDS, KALE AND MUSTARD - For control of aphids, armyworms, flea beetles, leafhoppers and mites, use 1 to 3 pints per acre. For stink bugs, use 2 to 3 pints per acre. Rates above 1 pint should not be applied closer than 21 days before harvest.

PEAS - For control of aphids, armyworms, flea beetles, leafhoppers, Lygus bugs and mites, use 1 to 2 pints per acre. For cowpea curculio, surface feeding and climbing cutworms and stink bugs, use 2 pints per acre. Rates above 1 pint should not be applied closer than 15 days before harvest.

TOMATOES - For control of aphids, armyworms, flea beetles, leafhoppers, mites and psyllids, use 1 to 3 pints per acre. Rates above 1 pint should not be applied closer than 15 days before harvest.

APPLY AT THE RATES INDICATED FOR THE FOLLOWING CROPS. MAKE NO APPLICATIONS CLOSER THAN 15 DAYS BEFORE HARVEST.

BEANS (Dry and Green) - For control of aphids, armyworms, flea beetles, leafhoppers, Lygus bugs and mites, use 1 to 3 pints per acre. For cowpea curculio, surface feeding and climbing cutworms and stink bugs, use 2 to 3 pints per acre. Rates above 1 pint should not be applied closer than 21 days before harvest.

BEETS (Red) - For control of aphids, armyworms, flea beetles, leafhoppers, Lygus bugs, mites and stink bugs, use 1 quart per acre. If tops are to be used for food or feed do not apply closer than 21 days before harvest.

CARROTS - For control of aphids, armyworms, surface feeding and climbing cutworms, flea beetles, leafhoppers and mites, use 1 quart per acre. Do not feed tops.

CELERY - For control of aphids, armyworms, flea beetles, leafhoppers, Lygus bugs, mites and stink bugs, use 1 quart per acre.

CUCUMBERS - For control of aphids and two-spotted mites, use 1/2 pint per acre. Do not apply before vining.

ONIONS - For control of thrips, use 1 pint per acre.

PEPPERS - For control of aphids, armyworms, surface feeding and climbing cutworms, flea beetles, leafhoppers and mites, use 1 quart per acre.

SPINACH - For control of aphids, armyworms, flea beetles, leafhoppers, mites, seed corn maggot, cutworms and stink bugs, use 1 to 2 pints per acre. Rates above 1 pint should not be applied closer than 21 days before harvest.

TURNIPS - For control of aphids, armyworms, flea beetles, leafhoppers, leafminers and mites, use 1 to 2 pints per acre. If tops are to be used for food or feed do not apply closer than 21 days before harvest.

APPLY AT THE RATES INDICATED FOR THE FOLLOWING CROPS. MAKE NO APPLICATIONS CLOSER THAN 21 DAYS BEFORE HARVEST.

LETTUCE - For control of aphids, armyworms, flea beetles, leafhoppers and mites, use 1 quart per acre.

FOREST & NONAGRICULTURAL

ORNAMENTAL - (Woody shrubs, tree, and vines, Christmas tree plantation (Pine - Apply 1 pint per acre.) European pine shoot moth - Apply 1 pint per acre. Harknackel pine tip moth - Apply 2 pints per acre.

FOREST, NONAGRICULTURAL LAND, WASTELAND - Mosquitoes (larvae) - Apply 1/4 pint per acre. DO NOT apply within 7 days of harvest. For use by mosquito abatement districts and official governmental agencies only. Repeat as needed. Toxic to fish and wildlife. DO NOT use where shrimp and crabs are an important resource. DO NOT use in mangroves or coastal, riparian areas along any coast line.

PINE FORESTS - European pine shoot moth - Harknackel pine tip moth - Apply 2 pints per acre. Apply in 5 to 10 gallons of water.

BEST AVAILABLE COPY

FORMULATED FOR
PLATTE CHEMICAL CO.
152 So. Main St.
Fremont, NE 68025

11655

CURSE A

BEST AVAILABLE COPY

