



PH 19 34704-10  
UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

MAR 4 1994

File 1046

J. Allen Dunlap III  
William M. Mahlborg  
Agent for: PLATTE CHEMICAL CO., INC.  
P. O. Box 667  
Greeley, CO 80632

OFFICE OF  
PREVENTION, PESTICIDES AND  
TOXIC SUBSTANCES

**Subject: Label Amendment Submission of 10/19/93 in Response to PR Notice 93-7  
EPA Reg. No. 34704-10  
CLEAN CROP METHYL PARATHION 4-E**

**Dear Registrant:**

The labeling cited above and submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA), as amended, is accepted subject to the comments reflected on the enclosed sheet. A copy of your proposed labeling stamped "ACCEPTED WITH COMMENTS" is enclosed.

**WHAT THIS ACCEPTANCE MEANS:**

Based on your certification, the Agency has accepted the labeling changes that are necessary to comply with the Worker Protection Standard (WPS) labeling requirements of 40 CFR part 156, subpart K, described in PR Notices 93-7 and 93-11. Any other labeling changes submitted in connection with this amendment application but not directly related to compliance with the WPS have not been reviewed or accepted by the Agency. If you wish to make such changes, you must submit a separate amendment application proposing them. If your product is currently suspended, the acceptance of this labeling amendment does not affect the suspension in any way.

**WHAT YOU NEED TO DO NEXT:**

By the next label printing make all the specified changes to your labeling. Send to EPA one (1) copy of the final printed labeling:

- BEFORE selling or distributing any product bearing the final printed labeling
- AND
- WITHIN one year from date of this acceptance.

**BEST AVAILABLE COPY**



Recycled/Recyclable  
Printed with SoyCanola Ink on paper that  
contains at least 50% recycled fiber

2 of 6

Page 2


Submit the final printed labeling via the U.S. Postal Service to:

Document Processing Desk (FIN-LABEL)  
Office of Pesticide Programs (7505C)  
U.S. Environmental Protection Agency  
401 M Street, SW  
Washington, D.C. 20460-0001

Hand or courier deliveries of final printed labeling may be made to:

Document Processing Desk (FIN-LABEL)  
Office of Pesticide Programs  
Room 266A, Crystal Mall 2  
1921 Jefferson Davis Highway  
Arlington, VA 22202

Sincerely,

  
Jim Tompkins, Deputy Chief  
Registration Support Branch  
Registration Division (7505W)

Attachment

BEST AVAILABLE COPY

IOPOSED  
BEL

## RESTRICTED USE PESTICIDE

Due to very high acute toxicity to Humans and Birds

For retail sale to and use only by certified applicator or persons under their direct supervision and only for those uses covered by the certified applicator's certification. Direct supervision for this product is defined as the certified applicator being physically present during application, mixing, loading, repair and cleaning of application equipment. Commercial certified applicators must also ensure that all persons involved in these activities are informed of the precautionary statements.



# METHYL PARATHION 4E

### Organophosphate Insecticide

Emulsifiable Liquid - An Agricultural Insecticide

#### ACTIVE INGREDIENTS:

\*O,O-dimethyl O-p-nitrophenyl  
phosphorothioate ..... 43.8%  
Related Products ..... 1.7%

INERT INGREDIENTS: ..... 54.5%

TOTAL 100.0%

\* Methyl Parathion

Contains 4 lbs. Methyl Parathion per Gallon.

KEEP OUT OF REACH OF CHILDREN

**DANGER**  
**PELIGRO**



**POISON**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

### DANGER—POISON—PRECAUTIONS



#### CAN KILL YOU IF SWALLOWED

This product can kill you if swallowed even in small amounts: spray mist or dust may be fatal if swallowed.



#### CAN KILL YOU BY SKIN CONTACT

This product can kill you if touched by hands or spilled or splashed on skin, in eyes, or on clothing (liquid goes through clothes).



#### CAN KILL YOU IF BREATHED

This product can kill you if vapors, spray mist or dust are breathed.

See Below For Statement of Practical Treatment, Antidote, and Additional Precautionary Statements

EPA REG. NO. 34704-10

EPA EST. NO. 34704-MS-1<sup>st</sup>, 2737-KS-1<sup>st</sup>

NET CONTENTS \_\_\_\_\_ GAL. (\_\_\_\_\_ L)

3730

WPSEX008R3

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS & DOMESTIC ANIMALS

## DANGER

The National Pesticide Telecommunication Network is available for recommendations regarding poisoning management, emergency treatment, and other information regarding the toxicity of methyl parathion. The toll free number for the National Pesticide Telecommunication Network is 1-800-858-7378.

If handled indoors provide mechanical exhaust ventilation. Keep all unprotected persons, children, livestock, and pets away from treated area or where there is danger of drift. Do not rub eyes or mouth with hands. If you feel sick in any way, STOP work and get help right away. See First Aid (Statement of Practical Treatment) section.

#### Personal Protective Equipment

Some individuals may be allergic to this product. See below for information on how to avoid allergic reactions. For the EPA chemical safety manual, see the section on "Allergic Reactions."

Handlers who may be exposed to the dust through application or

other tasks must wear

Handers who may be exposed to the concentrate through mixing, loading, application, or other tasks must wear

NEED ENG. CONT. (b)

### USER SAFETY RECOMMENDATIONS

When using this product, wear the following protective clothing and equipment: long-sleeved shirt, long pants, socks, shoes, and gloves. If you are using this product in a confined space, wear a respirator. If you are using this product in a confined space, wear a respirator. If you are using this product in a confined space, wear a respirator.

IMPORTANT! If pesticide comes in contact with skin, wash off with soap and water and contact a physician immediately.

Respirators must be cleaned and filters replaced according to instructions included with the respirators. DURING AERIAL APPLICATION, HUMAN FLAGGERS ARE PROHIBITED.

#### POISON SIGNS (Symptoms)

Methyl Parathion is a very dangerous poison. It rapidly enters the body on contact with all skin surfaces and eyes. Clothing wet with this material must be removed immediately. Exposed persons must receive prompt medical treatment or they may die.

Some of the signs and symptoms of poisoning are: Headache, nausea, vomiting, cramps, weakness, blurred vision, pin-point pupils, tightness in chest, labored breathing, nervousness, sweating, watering of eyes, drooling or frothing of mouth and nose, muscle spasms and coma.

#### + STATEMENT OF PRACTICAL TREATMENT +

Call a doctor (physician), clinic or hospital immediately. Explain that the victim has been exposed to methyl parathion and describe his condition. After first aid is given and if a doctor cannot come, take victim to clinic or hospital.

If breathing has stopped, start artificial respiration immediately and maintain until doctor sees victim.

If swallowed — Drink 1 or 2 glasses of water and induce vomiting by

Chem. Not. & P. not needed with chemical resistant suit.

BEST AVAILABLE COPY

# METHYL PARATHION 4E

EPA REG. NO. 34704-10

touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

In case of contact, immediately flush eyes or skin with plenty of water while removing contaminated clothing and shoes. See doctor immediately.

## NOTE TO PHYSICIAN

Antidote - administer atropine sulfate in large doses. TWO to FOUR mg. intravenously or intramuscularly as soon as cyanosis is overcome. Repeat at 5 to 10 minute intervals until signs of atropinization appear. 2-PAM chloride is also antidotal and may be administered in conjunction with atropine. DO NOT GIVE MORPHINE OR TRANQUILIZERS. Methyl Parathion is a strong cholinesterase inhibitor affecting the central and peripheral nervous systems and producing cardiac and respiratory depression. At first sign of pulmonary edema, the patient should be given supplemental oxygen and treated symptomatically. Continued absorption of the poison may occur and fatal relapses have been reported after initial improvement. VERY CLOSE SUPERVISION OF THE PATIENT IS INDICATED FOR AT LEAST 48 HOURS.

## ENVIRONMENTAL HAZARDS

This pesticide is highly toxic to aquatic invertebrates and wildlife. Birds in treated areas may be killed. Shrimp and other aquatic organisms may be killed at recommended application rates. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff and drift from target areas may be hazardous to aquatic organisms in adjacent aquatic sites. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters.

This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

## PHYSICAL & CHEMICAL HAZARDS

### FLAMMABLE

DO NOT USE OR STORE NEAR HEAT OR OPEN FLAME.

NOT FOR USE OR STORAGE IN OR AROUND THE HOME

## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

### AGRICULTURAL USE REQUIREMENTS

## STORAGE AND DISPOSAL

**PROHIBITIONS:** Do not contaminate water, food, or feed by storage or disposal. Do not store under conditions which might adversely affect the container or its ability to function properly.

**STORAGE:** Do not store below temperature of 0°F. Store in safe manner. Store in original container only. Keep container tightly closed when

not in use. Reduce stacking height where local conditions can affect package strength. Personnel should use clothing and equipment listed under "PRECAUTIONARY STATEMENT" when handling open containers.

**SPILLED MATERIAL:** Block or dike to prevent spreading of spill. Cover with absorbent material such as lime, clay or sawdust. Scoop and sweep into a disposable container. Wash area with strong lye solution, absorb and place into a disposable container.

**PESTICIDE DISPOSAL:** Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

**CONTAINER DISPOSAL:** Metal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or by other procedures approved by state and local authorities. Plastic: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

## DIRECTIONS

Be sure to read the precautionary statements before using!

This product is designed for application after dilution with water and for use by trained operators using airplane or power ground equipment. The hazards and precautions for handling the product in this container are equally applicable to it after dilution with water for spray application. Add the concentrate to the spray tank while filling with water, and mix thoroughly either by means of a tank agitator or pump by-pass. For best results, thoroughly cover all surfaces to be treated with spray. Rates of application given below should not be exceeded. Never apply later than indicated to assure residue levels at harvest are below tolerances established by the Food and Drug Administration.

Consult the State Agricultural Extension Service or Experiment Station for specific recommendations regarding application, dosage and timing of sprays. For application by ground equipment, add the desired amount of concentrate to sufficient water to apply at least 3 gallons of water per acre. For application by aircraft, add the amount of concentrate desired per acre to 1 to 3 gallons of water consistent with crop growth and good coverage. Greater quantities of water may be required to give sufficient coverage of orchard trees.

Do not use in undiluted form.

Do not apply this product through any type of irrigation system.

## CEREAL INSECTS

**BARLEY, OATS, RYE AND WHEAT** — For barley thrips, use 1/2 to 3/4 pints per acre. For aphids, black grass bugs, stink bugs and winter grain mites, use 1/2 to 1 1/2 pints per acre. For climbing cutworms and grasshoppers use 1/2 to 1 pint per acre. For armyworms and leafhoppers use 1 1/2 pints per acre. For chinch bugs, false chinch bugs and Say's plant bugs, use 1 1/2 pints per acre. For western or brown wheat mites and grasshoppers, use 1 pint per acre. Do not use more than 1/2 pint per acre within 15 days of harvest.

**CORN** — For control of armyworms, climbing cutworms, corn leaf aphids, corn rootworm adults, fall armyworms and stink bugs, use 1/2 pint per acre. For grasshoppers, use 1 pint per acre. Do not apply within 12 days of harvest. Do not apply during the pollen shed period.

**RICE** — For rice stink bugs, use 1 to 1 1/2 pints per acre. For grasshoppers, use 1 pint per acre. Do not apply within 15 days of harvest or within 14 days before or after a propanil application. Shrimp, crabs and crayfish may be killed. Do not apply where these are important resources.

## FIELD AND FORAGE CROPS

**ALFALFA AND CLOVER** — For control of alfalfa weevil larvae, aphids (including spotted alfalfa aphids) and climbing cutworms and grasshoppers, use 1/2 to 1 pint per acre. For alfalfa caterpillars, alfalfa adult weevils, armyworms, clover leaf weevils and webworms, use 1 pint per acre. For Egyptian alfalfa weevils, leafhoppers, Lygus bugs and spider mites, use 1 to 2 pints per acre. Do not treat closer than 15 days before harvest, cutting, or grazing. Do not apply to alfalfa if the crop or weeds in the treatment area are in bloom.

**COTTON** — For control of thrips, use 1/4 to 1/2 pint per acre. For cotton leafworms use 1/2 to 6 pints per acre. For grasshoppers and fall armyworms, use 1/2 to 6 pints per acre. For boll weevils, aphids and garden webworms, use 1/2 to 6 pints per acre. For false chinch bugs use 2 pints per acre. For cutworms, use 1 to 2 pints per acre.

For bollworms, saltmarsh caterpillars, armyworms, cotton leaf perforators and stink bugs, use 2 1/2 to 6 pints per acre. Applications should be made at 4 to 5 day intervals until control is obtained. At rates of 1/4 pint through 2 pints, application may be made up to the day before harvest. If harvest will be with mechanical pickers and if hand picking there is a 5 day interval. Mechanical pickers must be used with a 7 day preharvest interval at rates above 2 pints through 6 pints.

**DANGER** — When necessary to enter treated cotton fields within 48 hours after application, protective clothing and equipment must be worn.

**GRASS (Forage)** — For range caterpillar, use 1 pint per acre. For control of armyworms, crested wheat bugs, false chinch bugs, grasshoppers

BEST AVAILABLE COPY

12/20/93 LABEL

## METHYL PARATHION 4E

EPA REG. NO. 34704-10

and leafhoppers, use 1 1/2 pints per acre. Do not apply within 15 days of harvest or grazing.

**SOYBEANS** — For control of climbing cutworms, three-corned alfalfa hoppers, garden webworms, aphids, blister beetles, Mexican bean beetles, stink bugs, two-spotted mites and velvet bean caterpillars, use 3/4 to 2 pints per acre. For bollworms, fall armyworms and green cloverworms, use 1 quart per acre. Do not apply within 15 days of harvest up to and through 1 pint. Do not apply within 20 days of harvest through 1 quart.

**SUGAR BEETS** — For armyworms, cutworms, fall armyworms, flea beetles, grasshoppers, leafhoppers, mites, sugar beet webworms, use 1/2 to 3/4 pint per acre. Do not apply within 20 days of harvest; 60 days if tops are to be fed to animals.

**VETCH** — Alfalfa weevil (larvae) - Apply 1/2 to 1 pint per acre. Apply when 75% of terminals show feeding, when buds and shoots appear ragged or when the larvae counts reach 25 per sweep. Aphids, leafhoppers - Apply 1/2 to 1 pint per acre. Climbing cutworms, Lygus bugs, Vetch bruchid - 1 pint per acre. Armyworms - 1/2 to 2 1/2 pints per acre. DO NOT apply within 15 days of harvest through 2 pints. DO NOT apply within 20 days of harvest from 2 pints through 2 1/2 pints.

### OIL SEED CROPS

**SUNFLOWERS (30)** — For sunflower moths, sunflower maggot and sunflower seed weevil, use 2 pints per acre. Do not apply more than 3 times at 5-day intervals per season. Do not apply within 30 days of harvest.

### VEGETABLES

APPLY AT THE RATES INDICATED FOR THE FOLLOWING CROPS. MAKE NO APPLICATIONS CLOSER THAN 5 DAYS BEFORE HARVEST.

**POTATOES** — For control of aphids, armyworms, false chinch bugs, flea beetles, leafhoppers, mites and shield bugs, use 1 1/2 quarts per acre.

**SWEET POTATOES** — For control of aphids, armyworms, surface feeding and climbing cutworms, flea beetles, leafhoppers and mites, use 3/4 quart per acre.

APPLY AT THE RATES INDICATED FOR THE FOLLOWING CROPS. MAKE NO APPLICATIONS CLOSER THAN 7 DAYS BEFORE HARVEST.

**BROCCOLI** — For control of aphids, armyworms, flea beetles, leafhoppers and mites, use 1 to 3 pints per acre. For stink bugs, use 2 to 3 pints per acre. Rates above 1 pint should not be applied closer than 21 days before harvest.

**BRUSSELS SPROUTS** — For control of aphids, armyworms, flea beetles, leafhoppers and mites, use 1 to 3 pints per acre. For stink bugs, use 2 to 3 pints per acre. Rates above 1 pint should not be applied closer than 21 days before harvest.

**CAULIFLOWER** — For control of aphids, armyworms, flea beetles, leafhoppers and mites, use 1 to 3 pints per acre. For stink bugs, use 2 to 3 pints per acre. Rates above 1 pint should not be applied closer than 21 days before harvest.

**KOHLRABI** — For control of aphids, armyworms, flea beetles, leafhoppers and mites, use 1 to 3 pints per acre. For stink bugs, use 2 to 3 pints per acre. Rates above 1 pint should not be applied closer than 21 days before harvest.

**RUTABAGAS** — For control of aphids, armyworms, flea beetles, leafhoppers and mites, use 1 to 3 pints per acre. Rates above 1 pint should not be applied closer than 21 days before harvest.

APPLY AT THE RATES INDICATED FOR THE FOLLOWING CROPS. MAKE NO APPLICATIONS CLOSER THAN 10 DAYS BEFORE HARVEST.

**CABBAGE** — For control of aphids, armyworms, flea beetles, leafhoppers and mites, use 1 to 3 pints per acre. For stink bugs, use 2 to 3 pints per acre. Rates above 1 pint should not be applied closer than 21 days before harvest.

**COLLARDS, KALE AND MUSTARD GREENS** — For control of aphids, armyworms, flea beetles, leafhoppers and mites, use 1 to 3 pints per acre. For stink bugs, use 2 to 3 pints per acre. Rates above 1 pint should not be applied closer than 21 days before harvest.

**PEAS** — For control of Aphids, Armyworms, Flea Beetles, Leafhoppers, Lygus Bugs and Mites, use 1 to 2 pints per acre. For Cowpea Curculio, Surface Feeding and Climbing Cutworms and Stink Bugs, use 2 pints per acre. Make no application closer than 7 day before harvest at the 1 pint rate and at rates above 1 pint do not apply closer than 15 days before harvest. To avoid injury to bees, do not apply if bees are visiting the areas to be treated when crop or weeds are in bloom.

APPLY AT THE RATES INDICATED FOR THE FOLLOWING CROPS. MAKE NO APPLICATIONS CLOSER THAN FIFTEEN (15) DAYS BEFORE HARVEST.

5046

**BEANS (Dry and Green)** — For control of aphids, armyworms, flea beetles, leafhoppers, Lygus bugs and mites, use 1 to 3 pints per acre. For cowpea curculio, surface feeding and climbing cutworms and stink bugs, use 2 to 3 pints per acre. Dry Bean rates through 3 pints per acre, do not apply closer than 15 days before harvest. Green Bean rates through 1 pint per acre, do not apply closer than 15 days before harvest and for rates through 3 pints per acre, do not apply closer than 21 days before harvest. Do not apply if the crop or weeds in the treatment area are in bloom.

**CARROTS** — For control of aphids, armyworms, surface feeding and climbing cutworms, flea beetles, leafhoppers and mites, use 1 quart per acre. Do not feed tops.

**CELERY** — For control of aphids, armyworms, flea beetles, leafhoppers, Lygus bugs, mites and stink bugs, use 1 quart per acre.

**ONIONS** — For control of thrips, use 1 pint per acre.

**SPINACH** — For control of aphids, armyworms, flea beetles, leafhoppers, mites, seed corn maggots in crown and crown mites, use 1 to 2 pints per acre. Rates through 1 pint should not be applied closer than 15 days before harvest. Rates above 1 pint should not be applied closer than 21 days before harvest.

**TURNIPS** — For control of aphids, armyworms, flea beetles, leafhoppers, leafminers and mites, use 1 to 1 1/2 pints per acre. Rates through 1 pint per acre should not be applied closer than 7 days before harvest. Rates through 1 1/2 pints per acre, if tops are to be used for food or feed should not be applied closer than 21 days before harvest. If tops are NOT being used for food or feed do not apply closer than 15 days before harvest.

APPLY AT THE RATES INDICATED FOR THE FOLLOWING CROPS. MAKE NO APPLICATIONS CLOSER THAN 21 DAYS BEFORE HARVEST.

**BEETS (Red)** — For control of aphids, armyworms, flea beetles, leafhoppers, Lygus bugs, mites and stink bugs, use 1 quart per acre. Do not apply closer than 21 days before harvest.

**LETTUCE** — For control of aphids, armyworms, flea beetles, leafhoppers and mites, use 1 quart per acre.

### ATTENTION

Do not use in any manner other than recommended on this label. To avoid excessive residues of methyl parathion on food or forage crops, always observe the statements found under "Directions for Use," limiting the time before harvest when methyl parathion may be applied.

### NOTICE

PLATTE WARRANTS THAT THIS PRODUCT CONFORMS TO THE CHEMICAL DESCRIPTION ON THE LABEL THEREOF AND IS REASONABLY FIT FOR THE PURPOSES STATED ON SUCH LABEL ONLY WHEN USED IN ACCORDANCE WITH THE DIRECTIONS UNDER NORMAL USE CONDITIONS. IT IS IMPOSSIBLE TO ELIMINATE ALL RISKS INHERENTLY ASSOCIATED WITH THE USE OF THIS PRODUCT. CROP INJURY, INEFFECTIVENESS, OR OTHER UNINTENDED CONSEQUENCES MAY RESULT BECAUSE OF SUCH FACTORS AS WEATHER CONDITIONS, PRESENCE OF OTHER MATERIALS, OR THE MANNER OF USE OR APPLICATION, ALL OF WHICH ARE BEYOND THE CONTROL OF PLATTE. IN NO CASE SHALL PLATTE BE LIABLE FOR CONSEQUENTIAL, SPECIAL OR INDIRECT DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT. ALL SUCH RISKS SHALL BE ASSUMED BY THE BUYER.

EXCEPT AS EXPRESSLY PROVIDED HEREIN, PLATTE MAKES NO WARRANTIES, GUARANTEES, OR REPRESENTATIONS OF ANY KIND, EITHER EXPRESSED OR IMPLIED, OR BY USAGE OF TRADE, STATUTORY OR OTHERWISE, WITH REGARD TO THE PRODUCT SOLD, INCLUDING, BUT NOT LIMITED TO, MERCHANTABILITY, FITNESS FOR A PARTICULAR PURPOSE, USE OR ELIGIBILITY OF THE PRODUCT FOR ANY PARTICULAR TRADE USAGE.

ACCEPTED  
with COMMENTS  
In EPA Letter Dated

MAR 4 1994

Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

34704-10

**BEST AVAILABLE COPY**  
FORMULATED FOR  
PLATTE CHEMICAL CO.

150 SO. MAIN ST. FREMONT, NE 68025-5697

## RESTRICTED USE PESTICIDE

Due to very high acute toxicity to Humans and Birds  
For retail sale to and use only by certified applicator or persons under their direct supervision and only for those uses covered by the certified applicator's certification. Direct supervision for this product is defined as the certified applicator being physically present during application, mixing, loading, repair and cleaning of application equipment. Commercial certified applicators must also ensure that all persons involved in these activities are informed of the precautionary statements.

6076

## SUPPLEMENTAL LABELING

(CONTAINS ADDITIONAL INFORMATION WHICH MAY NOT APPEAR ON PACKAGE LABEL.)



# METHYL PARATHION 4E

EPA REG. NO. 34704-10

### CHANGE IN BEE STATEMENT FOR PEAS

**ALL APPLICABLE DIRECTIONS, RESTRICTIONS, AND PRECAUTIONS  
ON THE EPA REGISTERED LABEL ARE TO BE FOLLOWED.**

**THIS LABEL AND THE FEDERAL LABEL MUST BE IN POSSESSION  
OF USER AT THE TIME OF APPLICATION.**

### DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**PEAS — For control of Aphids, Armyworms, Flea Beetles, Leafhoppers, Lygus Bugs and Mites, use 1 to 2 pints per acre. For Cowpea Curcullo, Surface Feeding and Climbing Cutworms and Stink Bugs, use 2 pints per acre. Make no application closer than 7 days before harvest at the 1 pint rate and at rates above 1 pint do not apply closer than 15 days before harvest. To avoid injury to bees, do not apply if bees are visiting the areas to be treated when crop or weeds are in bloom.**

Read NOTICE on product label for limit of manufacturer's warranty.

**BEST AVAILABLE COPY**

FORMULATED FOR  
PLATTE CHEMICAL CO.

150 SO. MAIN STREET      FREMONT, NE 68025-5697