

WARNING

Keep out of reach of children. Avoid contact with skin, eyes, nose, mouth, or clothing. If contact occurs, wash immediately with plenty of water. If eyes are contacted, flush with water for 15 minutes. If swallowed, do not induce vomiting. If inhaled, get fresh air immediately. Wear protective clothing. During commercial or prolonged use, wear a respirator of a type passed by the National Institute for Occupational Safety and Health (NIOSH) for protection. Do not allow the material to be inhaled by unprotected humans or beneficial animals.

CAUTION

Do not use on treated area within 24 hours of application should wear protective clothing.

DO NOT STORE AT TEMPERATURES BELOW 20° F.

DO NOT USE IN UNLIFTED FORM

DO NOT DRINK, POOR, SPILL, OR STORE NEAR HEAT OR OPEN FLAME

Keep out of reach of children. Avoid contact with skin, eyes, nose, mouth, or clothing. If contact occurs, wash immediately with plenty of water. If eyes are contacted, flush with water for 15 minutes. If swallowed, do not induce vomiting. If inhaled, get fresh air immediately. Wear protective clothing. During commercial or prolonged use, wear a respirator of a type passed by the National Institute for Occupational Safety and Health (NIOSH) for protection. Do not allow the material to be inhaled by unprotected humans or beneficial animals.

THIODAN

Endosulfan (Hexachlorocyclopentadiene dimethoxy) 1, 4, 3-benzodioxane dimethoxy

Do not use on treated area within 24 hours of application should wear protective clothing. Do not use on treated area within 24 hours of application should wear protective clothing.

DO NOT ALLOW THIS MATERIAL TO DRIFT ONTO NEIGHBORING CROPS OR NON-CROP AREAS OR USE IN A MANNER OR AT A TIME OTHER THAN IN ACCORDANCE WITH LABEL DIRECTIONS BECAUSE ANIMAL, PLANT OR CROP INJURY, ILLEGAL RESIDUES OR OTHER UNDESIRABLE RESULTS MAY OCCUR.

DIRECTIONS FOR USE

NOT FOR USE OR STORAGE IN OR AROUND THE HOME.

Apply the listed amount per acre when insects first appear and repeat as required to maintain effective control. Use in sufficient water for thorough coverage. Unless otherwise noted, for listed crops, coverage of upper and lower leaf surfaces is essential for good control. Observe all limitations. If insect control is required beyond the following guidelines, supplement the control program with other suitable insecticides. Observe a 14-day interval between applications. Do not use on crops indicated by a cross (X) in the crop.

ROOT CROPS: Do not plant root crops other than carrots, potatoes, and sweet potatoes in treated areas.

When used on crops, Thiodan should be applied in the following manner: Apply the listed amount per acre in sufficient water for thorough coverage. Unless otherwise noted, for listed crops, coverage of upper and lower leaf surfaces is essential for good control. Observe all limitations. If insect control is required beyond the following guidelines, supplement the control program with other suitable insecticides. Observe a 14-day interval between applications. Do not use on crops indicated by a cross (X) in the crop.

THIODAN SHOULD BE USED IN ORIGINAL PACKAGING ONLY

PRECAUTIONS

Do not use on treated area within 24 hours of application should wear protective clothing. Do not use on treated area within 24 hours of application should wear protective clothing.

Do not use on treated area within 24 hours of application should wear protective clothing. Do not use on treated area within 24 hours of application should wear protective clothing.

CLEAN CROP

THIODAN® 3 E.C

WARNING

KEEP OUT OF REACH OF CHILDREN

See Side Panels For Additional Precautions

ACTIVE INGREDIENTS:

*Endosulfan (Hexachlorocyclopentadiene dimethoxy)

1, 4, 3-benzodioxane dimethoxy

Xylene

INERT INGREDIENTS:

CONTAINS 3 POUNDS ENDOSULFAN PER GALLON

Net Weight 1.75 LBS. (0.79 KG.)

Net Weight 1.75 LBS. (0.79 KG.)

Net Weight 1.75 LBS. (0.79 KG.)

Net Weight 1.75 LBS. (0.79 KG.)

Net Weight 1.75 LBS. (0.79 KG.)

Net Weight 1.75 LBS. (0.79 KG.)

Net Weight 1.75 LBS. (0.79 KG.)

AC
-9
JL
UNDER THE
FUNGICIDE
FOR ECONOMIC
ED UNDER
TO ATTACK

EPA Reg. No. 950-188

WARRANTY

34764-21. 1000-1000 960-188

THIODAN® 3 E.C.

See Side Panels For Additional Precautions.

osulfan (Hexachlorohydromethano-
-benzodiox (thiophin oxide)

INGREDIENTS:	60.7%
	6.0%
	100.0%

The **dm[®]** is a registered trademark of Canadian Yachest Ltd.

* U. S. Patent No. 2,799,685

NET CONTENTS _____ U. S. GALLON(S)

EPA Reg. No. 960-188

WARRANTY

Reveals the warrants that this material conforms to the original description on the offer and is

[illegible]

WARNING

KEEP OUT OF REACH OF CHILDREN
See Side Panel For Additional Precautions.

INGREDIENTS:

Endosulfan (Hexachlorohydromethano- benzodioxathiepin oxide)	33.3%
.....	60.7%
INGREDIENTS:	6.0%
	100.0%

CONTAINS 3 POUNDS ENDOSULFAN PER GALLON.

Thiodan[®] is a registered trademark of Canadian Hoechst, Ltd.


*U. S. Patent No. 2,799, 685

NET CONTENTS _____ U. S. GALLON(S)

EPA Reg. No. 960-188

WARRANTY

Balcom Chemicals, Inc. warrants that this material conforms to the chemical description on the label and is fit for the purposes referred to in the directions for use. The product is sold subject to the understanding that the buyer assumes all risks of use or handling which may result in loss or damage which are beyond the control of the seller, such as for example incompatibility with other products, the manner of its use, the presence of other products or materials in or on the soil or crop. Balcom Chemicals, Inc. makes no other express or implied warranty of fitness or merchantability. The exclusive remedy of the buyer, and the limit of the liability of Balcom Chemicals, Inc. or any other seller, for any and all losses, damages resulting from the use or handling of this product shall be the purchase price paid by the buyer for the quantity of this product involved. The buyer and all users are deemed to have accepted this notice, which may not be varied by any verbal or written agreement.



Tomatoes (Field & Greenhouse) (1)	Stink Bug	Use field application of 1-1/3 to 2 qts./100 gals.	
	Blister Beetle, Colorado	2/3 qt.	
	Blister Beetle, Fila		
	Green Peach Aphid		
	Heliothis	2/3 to 1 1/3 qts.	
	Tomato Russet Mite	1-1/3 qts.	
	Tomato Fruitworm		
	Yellow-striped Armyworm		
	Stink Bugs	1 qt. to 1 1/2 qt.	
	Colorado Potato Beetle	1 qt.	
	Whitefly	2 qt./100 gals. in 100 to 200 gals. of water	
Apples (30)	Apple Aphid, Rusty	2/3 qt./100 gals. of water or 2 2/3 to 3 1/3 qts./acre	Do not feed produce from treated apples to livestock. Do not feed 2 applications during the fruiting period at rate above 1 qt./acre.
	Apple Aphid		
	Apple Rust Mite	1 qt. to 2 2/3 qt./100 gals. of water, or 1 gal. to 5 1/3 qts./acre.	
Cherry (21)	Plum Rust Mite	2/3 qt./100 gals. or 2 2/3 to 3 1/3 qts./acre	Do not exceed 2 applications after shuck split.
	Mineola Moth (Michigan only)	1-1/3 qts./100 gals. applied in delayed dormant period	
Peaches (30), Apricots (30), Nectarines (30)	Black Cherry Aphid	2/3 qt./100 gals. or 2 2/3 to 3 1/3 qts./acre	Do Not exceed 2 applications during fruiting period. For Lesser Peach Tree Borer, thorough wetting of trunks and main branches is important. For foliar application, do not apply more than twice per season.
	Black Peach Aphid, Cat		
	Leafhopper, Green		
	Peach Aphid, Peach		
	Peach Borer, Rusty Plum		
	Aphid		
	Lesser Peach Tree Borer	1 qt./100 gals.	
	Peach Tree Silver Mite	1/3 qt./100 gals. or 2-2/3 to 3-1/3 qt./acre	
Plums (7), Prunes (7)	Lesser Peach Tree Borer	1 qt./100 gals. or 2-2/3 to 3-1/3 qts./acre	Apply spray to thoroughly wet trunks and main branches.
	Peach Tree Borer	1 qt./100 gals. or 2 2/3 to 3-1/3 qts./acre	Spray thoroughly to cover all bark areas from ground to scaffold.
Cherry, Peach, Plum, Nursery Stock Dip	Peach Tree Borer	2/3 qt./100 gals. of water	Thoroughly soak. Immerse the bobbles of trees so that the roots and crowns are covered well above the grafting union. Plant in nursery or dry before returning to storage. Wear rubber gloves when handling.
Grapes (7)	Grape Leafhopper	2 1/3 qt./100 gals. or 1/3 to 2 qts./acre	Do not apply to Concord variety as severe injury is likely to result.
	Rose Chafer	2/3 qt./100 gals.	
Pears (7)	Conspicuous Stink Bug	GROUND TREATMENT Use 2/3 qt./100 gals. 200-400 gals./acre applied to orchard floor & ground around trees prior to bloom.	Do not graze livestock in treated orchard.
		FOLIAR TREATMENT Use 2/3 qt./100 gals. or 2 2/3 to 3 1/3 qts./acre	Stink Bugs may be wet to obtain control. The rate for a foliar treatment on stink bug control is 10 days. Do Not apply more than once per season.
	Pear Rust Mite	2/3 qt./100 gals.	
	Pear Leaf Blister Mite	1/3 - 2/3 qt./100 gals. applied as a post-harvest or dormant treatment	
	Pear Psylla	To control second generation pear psylla use 1 qt./100 gals. not to exceed 330 gals. of spray/acre.	Apply when nymphs are small and repeat if necessary in 7 to 10 days.
Pecans	Black Pecan Aphid	1 qt./100 gals.	Do not exceed 2 applications during the fruiting period.
Ornamental (greenhouse & out of doors)	Aphids, Cyclamen Mite, Whitefly, Rose Chafer	2/3 qt./100 gals.	Do not apply after shuck split. Do not graze livestock in treated groves. On chrysanthemums, best results will be obtained if applied before plants flower. Do not apply to "Tommygold", "Red Chrysanthemum", and "White Knight" chrysanthemums as injury may result. Do not spray on birch trees. In greenhouse applications wear approved protective clothing.
Box Elder	Box Elder Bug	1 qt./100 gals.	Apply in mid-April and again if necessary in late fall. Thoroughly wet bark and main stem, especially where branches join main stem.
Dogwood	Dogwood Borer	2 1/3 qt./100 gals.	Apply to main stem and trunk, when borer is first seen. Do not apply to main stem and trunk, when borer is first seen. Do not apply to main stem and trunk, when borer is first seen.
Lilac	Lilac Borer	2 1/3 qt./100 gals.	Apply to main stem and trunk, when borer is first seen. Do not apply to main stem and trunk, when borer is first seen.
Taxus	Taxus Bud Mite	2 1/3 qt./100 gals.	Apply to main stem and trunk, when borer is first seen. Do not apply to main stem and trunk, when borer is first seen.
Black Vine Weevil	Black Vine Weevil	2 1/3 qt./100 gals.	Apply to main stem and trunk, when borer is first seen. Do not apply to main stem and trunk, when borer is first seen.
Bark Beetles (Pitychus & Scolytus)	Bark Beetles (Pitychus & Scolytus)	2 1/3 qt./100 gals.	Apply to main stem and trunk, when borer is first seen. Do not apply to main stem and trunk, when borer is first seen.
Black Weevils (Hippodamia & Curculio)	Black Weevils (Hippodamia & Curculio)	2 1/3 qt./100 gals.	Apply to main stem and trunk, when borer is first seen. Do not apply to main stem and trunk, when borer is first seen.

COM

CHEMICALS, INC.



P.O. BOX 567

GREELEY, COLORADO 80631

PHONE 303-352-4750