

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS & DOMESTIC ANIMALS  
CAUTION**

Causes eye and skin irritation. Harmful if swallowed, inhaled or absorbed through skin. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Wash with soap and water after handling. Avoid breathing dust. Use with adequate ventilation. Do not contaminate food or foodstuffs.

**ENVIRONMENTAL HAZARDS**

This product is toxic to fish. Do not apply directly to lakes, streams or ponds. Do not contaminate water by cleaning of equipment or disposal of waste. **NOTE TO USER:** Federal Government and most states have seed laws and regulations requiring an appropriate caution statement be affixed to bags containing treated seed such as "Treated Seed—Do Not use for food or feed."

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

**STORAGE AND DISPOSAL**

**PROHIBITIONS:** Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.  
**PESTICIDE DISPOSAL:** Pesticide, spray mixture, or rinseate that cannot be used or chemically reprocessed should be disposed of in a landfill approved for pesticides or buried in a safe place away from water supplies.  
**CONTAINER DISPOSAL:** Dispose of in an incinerator or landfill approved for pesticide containers, or bury in a safe place.  
**GENERAL:** Consult federal, state or local disposal authorities for approved alternative procedures such as limited open burning. Store in a cool, dry place. Protect from excessive heat.


**APPLES:** Apple Scab, Brooks Fruit Spot, Flyspecking, Bitter Rot, Black Pox, Botryosphaena—Use 25 to 40 lbs. per acre. Apply at delayed dormant period. Repeat at weekly intervals as required to maintain control.  
**APRICOTS:** Brown Rot, Jacket Rot—Use 50 to 60 lbs. per acre. Apply at red bud stage and repeat at 75% petal fall.  
**CHERRIES:** Cherry Leaf Spot, Brown Rot—Use 50 to 60 lbs. per acre. Apply at popcorn, blossom and petal fall stages. Continue at 10 to 14 day intervals as required.  
**PEACHES:** Brown Rot, Peach Scab—Use 50 to 60 lbs. per acre. Make 2 to 3 pre-bloom applications. Repeat at petal-fall and 3 to 4 covers as necessary. Make pre harvest application where necessary.  
**PEARS:** Pear Scab (except Pacific Northwest)—Use 25 to 40 lbs. per acre. Apply during early finger and petal-fall stages. Repeat at weekly intervals as necessary to maintain control. Russeting may be reduced on Bosc pears. Do not use on D'Anjou pears.  
**RASPBERRIES:** Fruit Rot—Use 30 lbs. per acre making application 3 to 5 days before harvest starts. At mid harvest and 8 to 10 days later.  
**STRAWBERRIES:** Botrytis Rot—Use 40 lbs. per acre. Make the first application when new growth starts in the spring and before the fruit starts to form. Repeat at weekly intervals. During severe infection periods continue through harvest, treating immediately after each picking.  
**TOMATOES:** Anthracnose, Early Blight, Late Blight, Septoria Leaf Spot—On young plants use 25 to 30 lbs. per acre. On mature plants use 40 to 50 lbs. per acre. Begin applications at first sign of disease. Repeat at 7 to 10 day intervals.  
**POTATOES, SEED PIECES:** Seed Decay (Fusarium), Damping Off, Verticillium Wilt, Brown Eye Disease—Apply sufficient amount of Captan 7.5 Dust to provide a thorough coverage (7 to 10 ounces per bushel of cut seed pieces or 10 ounces to 1 lb. per 100 lbs. of cut seed pieces). Whole or 1-drop seed will need less material. Depending on the number of cuts made in the seed piece, the amount of dust used will vary as a result of the number of surface areas exposed for treatment. Captan 7.5 Dust will NOT protect against fungal inoculum already within the tuber or bacterial diseases. Apply in convenient container or by duster attachment over belt. Cut pieces should be treated within 6 hours of cutting. If planting is to be delayed more than 1 to 2 days the treated pieces should be stored for 2 to 3 days in open crates before bagging. Do not use treated seed for food or feed purposes.  
**YOUNG PLANTS:** Early Blight, Late Blight—Use 30 to 35 lbs. per acre. Begin applications when plants are 2 to 4 inches high. Repeat at 7 to 10 day intervals as long as disease threatens.  
**Mature Plants:** Early Blight, Late Blight—Use 40 lbs. per acre. Begin applications at first sign of disease. Repeat at 7 to 10 day intervals.  
**ORNAMENTALS (ROSES, CARNATIONS, CHRYSANTHEMUMS):** Black Spot, Leaf Spot, Rust Flower Blight—Apply at first sign of disease. Repeat at 7 to 10 day intervals as necessary. Dust plants thoroughly.

**NOTICE**

**DISCLAIMER—**The label instructions for the use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of Platte Chemical Company. All such risks shall be assumed by the user. Platte Chemical Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use thereon described when used in accordance with the directions for use, subject to the risks referred to above. Any damages arising from a breach of this warranty shall be limited to direct damages and shall not include consequential commercial damages such as loss of profits or values or any other special or indirect damages. Platte Chemical Company makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY.

1000

34704-22



ACTIVE INGREDIENT

INERT INGREDIENTS

**CAUTION  
KEEP OUT OF REACH OF CHILDREN.**

**STATEMENT OF PRACTICAL TREATMENT**

**In Case of Eye Contact:** Immediately flush eyes with plenty of water for at least 15 minutes. Call a Physician.

**In Case of Skin Contact:** Wash skin with soap and warm water.

**See Side Panel for Additional Precautionary Statement**

EPA REG. NO. 34704-22

**NET WEIGHT 50 POUNDS**

39

FORMULATED FOR

**PLATTE CHEMICAL CO., Inc.**

STREET

FREMONT, NEBRASKA 68001

P  
CI

P