

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS

DANGER

Causes eye and skin irritation. Harmful if swallowed, inhaled or absorbed through skin. Corrosive to eyes. Causes eye damage. Do not get in eyes, on skin or on clothing. Wear goggles or face shield and rubber gloves when handling. Wash with soap and water after handling. Avoid breathing dust. Use with adequate ventilation. Do not contaminate food or foodstuff.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to water or wetlands. Do not contaminate water by cleaning of equipment or disposal of waste. **NOTE TO USER:** Federal Government and most states have seed laws and regulations requiring an appropriate caution statement be affixed to bags containing treated seed such as "Treated Seed—Do Not use for food or feed."

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

RE-ENTRY STATEMENT

Do not enter treated areas without protective clothing until dusts have settled. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Do not store under conditions which might adversely affect the container or its ability to function properly. Such conditions include, but are not limited to, positioning of the container in storage, storage temperature, potential for crushing or damage due to stacking, and penetration of moisture.

STORAGE: Store in safe manner. Store in original container only. Store in cool, dry place. Reduce stacking height where local conditions, such as humidity or pallet overhang, can affect package strength. Personnel should use clothing and equipment consistent with good pesticide handling.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

APPLES: Apple Scab, Brooks Fruit Spot, Flyspecking, Bitter Rot, Black Pox, Botryosphæria—Use 25 to 40 lbs per acre. Apply at delayed dormant period. Repeat at weekly intervals as required to maintain control.

APRICOTS: Brown Rot, Jacket Rot—Use 50 to 60 lbs per acre. Apply at red bud stage and repeat at 75% petal-fall.

CHERRIES: Cherry Leaf Spot, Brown Rot—Use 50 to 60 lbs per acre. Apply at popcorn, blossom and petal fall stages. Continue at 10 to 14 day intervals as required.

PEACHES: Brown Rot, Peach Scab—Use 50 to 60 lbs per acre. Make 2 to 3 pre-bloom applications. Repeat at petal fall and 3 to 4 covers as necessary. Make pre-harvest application where necessary.

PEARS: Pear Scab (except Pacific Northwest)—Use 25 to 40 lbs per acre. Apply during early finger and petal-fall stages. Repeat at weekly intervals as necessary to maintain control. Russeting may be reduced on Bosc pears. Do not use on D'Anjou pears.

RASPBERRIES: Fruit Rot—Use 30 lbs per acre making application 3 to 5 days before harvest starts, at mid-harvest and 8 to 10 days later.

STRAWBERRIES: Botrytis Rot—Use 40 lbs per acre. Make the first application when new growth starts in the spring and before the fruit starts to form. Repeat at weekly intervals. During severe infection periods continue through harvest, treating immediately after each picking.

TOMATOES: Anthracnose, Early Blight, Late Blight, Septoria Leaf Spot—On young plants use 25 to 30 lbs per acre. On mature plants use 40 to 50 lbs per acre. Begin applications at first sign of disease. Repeat at 7 to 10 day intervals.

POTATOES, SEED PIECES: As an aid in the control of seed piece decay caused by soil born *fusarium*. CAPTAN 7.5 DUST will NOT protect the seed piece against bacterial diseases (such as soft rot or black leg caused by *Erwinia*) in or on the seed piece or in the soil. Captan will NOT protect against *fusarium* already established in or on the seed piece. Apply sufficient amount of CAPTAN 7.5 DUST to provide a thorough coverage (7 to 10 ounces per bushel of cut seed pieces or 10 ounces to 1 lb per 100 lbs of cut seed pieces). Whole or 1-drop seed will need less material. Depending on the number of cuts made in the seed piece, the amount of dust used will vary as a result of the number of surface areas exposed for treatment. Apply in convenient container or by dustler attachment over belt. Cut pieces should be treated within 6 hours of cutting. If planting is to be delayed more than 1 to 2 days the treated pieces should be stored for 2 to 3 days in open crates before bagging. Do not use treated seed for food or feed purposes.

NOTE TO USER: Tuber damage and/or decay may occur on sandy soils under moist soil conditions and higher than normal soil temperatures. Do not use if these conditions can occur or if there has been a past history of seed piece decay due to this problem.

This product will not control bacterial diseases.

Not for use in Jefferson County, Idaho.

Young Plants: Early Blight, Late Blight—Use 30 to 35 lbs per acre. Begin application when plants are 2 to 4 inches high. Repeat at 7 to 10 day intervals as long as disease threatens.

Mature Plants: Early Blight, Late Blight—Use 40 lbs per acre. Begin applications at first sign of disease. Repeat at 7 to 10 day intervals.

ORNAMENTALS (ROSES, CARNATIONS, CHRYSANTHEMUMS): For control of black spot and Botrytis blossom blight of roses. Alternaria leaf spot and rust of carnations and Botrytis flower blight and Septoria leaf spot of chrysanthemums—Apply at first sign of disease. Repeat at 7 to 10 day intervals as necessary. Dust plants thoroughly.



ACTIVE INGREDIENT
Captan (N-trichloro
cyclohexane-1, 2
Related compound)
INERT INGREDIENT

KEEP OUT

STATEMENT
In Case of Eye Contact: Irrigate
15 minutes. Call a Physician.
In Case of Skin Contact: Wash

See Side Panel for
EPA REG. NO. 3470

NET

DISCLAIMER—The label instructions are based on field use and tests. The user should read the label carefully. However, it is important to use this product carefully because of such factors as weather, soil conditions, and method of use or application all of which affect the results. The user assumes all risks. The user's use of this product should conform to the use therein described when the risks referred to above. Any damages or losses of profits or values or any other special damages shall not be recovered from the manufacturer or distributor. No other express or implied warranty is made by the manufacturer or distributor. **FITNESS or MERCHANTABILITY**

* CLEAN CROP is a Registered Trademark

10015

**STATEMENTS
DOMESTIC ANIMALS**

PRECAUTIONS
Do not get on skin. Corrosive to eyes. Causes eye damage. Do not eat or drink when handling. Wash with soap and water after handling food or foodstuff.


HAZARDS
Do not contaminate water by cleaning of equipment or disposal. Federal laws and regulations requiring an appropriate caution statement—Do Not use for food or feed.

PRECAUTIONARY USE
Read and follow all directions on its labeling.

RESTRICTIONS
Because certain states may require more restrictive reentry restrictions of Agriculture for further information. Do not apply this product to other persons. The area being treated must be vacated by un-

DISPOSAL
Do not store under conditions which might adversely affect the product, but are not limited to, positioning of the container in stacking, and penetration of moisture. Store in a cool, dry place. Reduce stacking height where local conditions require. Personnel should use clothing and equipment consistent with the use of the product.
Do not dispose of on site or at an approved waste disposal facility. Dispose of empty bag in a sanitary landfill or by incineration. Stay out of smoke.

Botryosphaeria—Use 25 to 40 lbs. per acre. Apply at delayed bloom stage and repeat at 75% petal fall.
Apply at popcorn, blossom and petal fall stages. Continue at 10 to 15 day intervals.
3 pre bloom applications. Repeat at petal fall and 3 to 4 covers.
Apply during early finger and petal fall stages. Repeat at weekly intervals.
Bosc pears. Do not use on D'Anjou pears.
Apply 10 to 15 days before harvest starts, at mid harvest and 8 to 10 days later.
Apply when new growth starts in the spring and before the fruit begins to ripen through harvest, treating immediately after each picking.
On young plants use 25 to 30 lbs. per acre. On mature plants use 25 to 30 lbs. per acre at 7 to 10 day intervals.
Diseases caused by soil born *Fusarium*. CAPTAN 7.5 DUST will NOT control diseases caused by *Erwinia* in or on the seed piece or in the soil.
Apply sufficient amount of CAPTAN 7.5 DUST to 100 lbs. of cut seed pieces or 10 ounces to 1 lb. per 100 lbs. of cut seed pieces. Whole seed pieces. The amount of dust used will vary as determined by the dust container or by duster attachment over belt. Cut pieces should be stored for 2 to 3 days before use. The treated pieces should be stored for 2 to 3 days before use.
Apply at most soil conditions and higher than normal soil temperatures. The rate of seed piece decay due to this product is dependent on soil conditions.
Apply when plants are 2 to 4 inches high. Repeat at 7 to 10 day intervals.
Applications at first sign of disease. Repeat at 7 to 10 day intervals for control of black spot and Botrytis blossom blight of roses. After application for control of leaf spot of chrysanthemums—Apply at first sign of disease.



**CAPTAN
7.5
DUST
Fungicide**

| ACTIVE INGREDIENT: | BY WEIGHT: |
|---|---------------|
| Captan (N-trichloromethylthio-4-cyclohexane-1, 2-dicarboximide) | 7.1% |
| Related compounds | 0.4% |
| INERT INGREDIENTS: | 92.5% |
| TOTAL | 100.0% |

**KEEP OUT OF REACH OF CHILDREN
DANGER**

STATEMENT OF PRACTICAL TREATMENT

In Case of Eye Contact: Immediately flush eyes with plenty of water for at least 15 minutes. Call a Physician.
In Case of Skin Contact: Wash skin with soap and warm water.

See Side Panel for Additional Precautionary Statement

EPA REG. NO. 34704-22 EPA EST. NO. 38091-ID-1

NET WEIGHT 40 POUNDS

NOTICE

DISCLAIMER—The label instructions for the use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of Platte Chemical Company. All such risks shall be assumed by the user. Platte Chemical Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use, subject to the risks referred to above. Any damages arising from a breach of this warranty shall be limited to direct damages and shall not include consequential damages, such as loss of profits or values or any other special or indirect damages. Platte Chemical Company makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY.

CLEAN CROP is a Registered T.M. of United Agr. Products, Inc.