

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS**

DANGER

Causes irreparable eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Avoid contact with skin and clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Drift and runoff from treated areas may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water by cleaning of equipment or disposal of wastes. In the event that treated seeds are spilled and cannot be salvaged, completely cover them with soil. **NOTE TO USER:** Federal Government and most states have seed laws and regulations requiring an appropriate caution statement be affixed to bags containing treated seed such as: "Treated Seed—Do Not use for food or feed."

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

RE-ENTRY STATEMENT

Do not allow persons to enter treated areas within 4 days following application unless a long-sleeved shirt and long pants or a coverall that covers all parts of the body except the head, hands and feet and chemically resistant gloves are worn. Because certain states may require more restrictive re-entry intervals for various crops treated with this product, consult your State Department of Agriculture for local information. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Conspicuously post reentry information at site of application.

Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. When oral warnings are given, warnings shall be given in a language customarily understood by workers.

Written or oral warnings must include the following information:

DANGER

(Insert area or field description) treated with Captan on (insert date of application) Do not allow persons to enter treated areas within 4 days following application unless a long-sleeved shirt and long pants or a coverall that covers all parts of the body except the head, hands and feet and chemically resistant gloves are worn.

In case of accidental exposure: Call a doctor (physician), clinic or hospital immediately. Explain that the victim has been exposed to Captan and describe his condition. For further information see the "STATEMENT OF PRACTICAL TREATMENT" portion of the pesticide label.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container. Do not store under conditions which might adversely affect the container or its ability to function properly. Such conditions include, but are not limited to, positioning of the container in storage, storage temperature, potential for crushing or damage due to stacking, and penetration of moisture.

STORAGE: Store in safe manner. Store in original container only. Store in cool, dry place. Keep container tightly closed when not in use. Reduce stacking height where local conditions, such as humidity or pallet overhang, can affect package strength. Personnel should use clothing and equipment consistent with good pesticide handling.

PESTICIDE DISPOSAL: Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal Law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

CONTAINER DISPOSAL: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Wear goggles or face shield when handling. Wash thoroughly with soap and water after handling. Remove contaminated clothing and separately launder clothing before reuse. Mixers/loaders when mixing and loading this product must wear a long-sleeved shirt and long pants or a coverall that covers all parts of the body except the head, hands and feet. Wear chemical resistant gloves, goggles or a face shield and a chemically resistant apron.

Fieldworkers and harvesters must wear chemical resistant gloves. Leather or fabric gloves are not acceptable.

A dust mask must be worn when mixing and loading this product.

During application only, from a tractor with a completely enclosed cab or from an aircraft with an enclosed cockpit a long-sleeved shirt and long legged pants must be worn. Chemically resistant gloves must be available in the cab or cockpit and worn while exiting.

AFTER WORK: Take off all clothes and shoes. Shower using soap and water. Do not wear contaminated clothing. Wash protective clothing and equipment with soap and water after each use. Personal clothing worn during use must be laundered separately from household articles. Clothing and protective equipment heavily contaminated or drenched with captan must be destroyed according to state and local regulations.



**CAPTAN
7.5 DUST
Fungicide**

ACTIVE INGREDIENTS:	BY WEIGHT
Captan (N-trichloromethylthio-4-cyclohexene-1,2-dicarboximide)	7.33%
Related derivatives	0.17%
INERT INGREDIENTS	92.50%
TOTAL	100.00%

**KEEP OUT OF REACH OF CHILDREN
DANGER PELIGRO**

PRECAUCION AL USUARIO: Si usted no lee Ingles, no use este producto hasta que la etiqueta le haya sido explicada amplamente.

STATEMENT OF PRACTICAL TREATMENT

If in eyes: Flush with plenty of water. Call a physician.

If swallowed: Call physician or Poison Control Center immediately. Drink promptly a large quantity of milk, egg whites, gelatin solutions, or if these are not available, drink large quantities of water. Avoid alcohol. Do not give anything by mouth to an unconscious person.

If inhaled: Remove victim to fresh air. If not breathing give artificial respiration, preferably mouth-to-mouth. Get medical attention.

In on skin: Wash skin with soap and warm water

See Side Panel for Additional Precautionary Statements

EPA REG. NO. 34704-22 EPA EST. NO. 38091-ID-1

NET WEIGHT 50 POUNDS

CLEAN CROP is a Registered T.M. of United Agri Products, Inc.

17144

EXP-20

APPLES: Apple Scab, Brown Rot, Powdery Mildew, Bitter Rot, Black Rot, Botrytis, Phoma—Use 25 to 40 lbs. per acre. Apply at delayed dormant period. Repeat at weekly intervals as required to maintain control.

APRICOTS: Brown Rot, Jacket Rot—Use 50 to 60 lbs. per acre. Apply at red bud stage and repeat at 75% petal-fall.

CHERRIES: Cherry Leaf Spot, Brown Rot—Use 50 to 60 lbs. per acre. Apply at popcorn, blossom and petal fall stages. Continue at 10 to 14 day intervals as required.

PEACHES: Brown Rot, Peach Scab—Use 50 to 60 lbs. per acre. Make 2 to 3 pre-bloom applications. Repeat at petal-fall and 3 to 4 covers as necessary. Make pre-harvest application where necessary.

PEARS: Pear Scab (except Pacific Northwest)—Use 25 to 40 lbs. per acre. Apply during early leafing and petal-fall stages. Repeat at weekly intervals as necessary to maintain control. Rinsing may be produced on Josc pears. Do not use on D'Anjou pears.

STRAWBERRIES: Botrytis Rot—Use 40 lbs. per acre. Make the first application when new growth starts in the spring and before the 2nd starts to form. Repeat at weekly intervals. During severe infection periods continue through harvest, treating immediately after each picking.

TOMATOES: Anthracnose, Early Blight, Late Blight, Septoria Leaf Spot—On young plants use 25 to 30 lbs. per acre. On mature plants use 40 to 50 lbs. per acre. Begin applications at first sign of disease. Repeat at 7 to 10 day intervals.

POTATOES, SEED PIECES: As an aid in the control of seed piece decay caused by soil borne *fusarium*, CAPTAN 7.5 DUST will NOT protect the seed piece against bacterial diseases (such as soft rot or black leg caused by *Erwinia*) in or on the seed piece or in the soil. Captan will NOT protect against *fusarium* already established in or on the seed piece. Apply sufficient amount of CAPTAN 7.5 DUST to provide a thorough coverage (7 to 10 ounces per bushel of cut seed pieces or 10 ounces to 1 lb. per 100 lbs. of cut seed pieces). Whole or 1-drop seed will need less material. Depending on the number of cuts made in the seed piece, the amount of dust used will vary as a result of the number of surface areas exposed for treatment. Apply in convenient container or by duster attachment over belt. Cut pieces should be treated within 6 hours of cutting. If planting is to be delayed more than 1 to 2 days the treated pieces should be stored for 2 to 3 days in open crates before bagging. Do not use treated seed for food or feed purposes.

NOTE TO USER: Tuber damage and/or decay may occur on sandy soils under moist soil conditions and higher than normal soil temperatures. Do not use if these conditions can occur or if there has been a past history of seed piece decay due to this problem.

This product will not control bacterial diseases.

Not for use in Jefferson County, Idaho.

ORNAMENTALS (ROSES, CARNATIONS, CHRYSANTHEMUMS): For control of black spot and Botrytis blossom blight of roses, Alternaria leaf spot and rust of carnations and Botrytis flower blight and Botrytis leaf spot of chrysanthemums—Apply at first sign of disease. Repeat at 7 to 10 day intervals as necessary. Dust plants thoroughly.

NOTICE

The label instructions for the use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as: weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of Platte Chemical Company. All such risks shall be assumed by the user. Platte Chemical Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use therein described when used in accordance with the directions for use, subject to the risks referred to above. Any damages arising from a breach of this warranty shall be limited to direct damages and shall not include consequential commercial damages such as loss of profits or values or any other special or indirect damages. Platte Chemical Company makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY.

FORMULATED FOR
PLATTE CHEMICAL COMPANY, INC.
150 SO. MAIN STREET FREMONT, NEBRASKA 68025