

CLEAN CROP®
SEVIN®5 BAIT

FOR THE CONTROL OF ANTS, ARMYWORMS,
CRICKETS, CUIWORMS, AND GRASSHOPPERS

ACTIVE INGREDIENT:

Carbaryl (1-Naphthyl methylcarbamate).....	5%
INERT INGREDIENTS:.....	95%
TOTAL.....	100%


SEVIN is the Registered T.M. of Union Carbide Corporation for Carbaryl Insecticide.

CLEAN CROP is a Registered T.M. of United Agri Products, Inc.

KEEP OUT OF REACH OF CHILDREN (12 pt.)
CAUTION (18 pt.)

EPA REG. NO. 34704-23
EPA EST. NO.

NET WEIGHT _____ LBS.



STATEMENTS OF PRACTICAL TREATMENTS

If on Skin: Wash thoroughly with soap and water.
If Swallowed: Induce vomiting by drinking 1 or 2 glasses of water and touching back of throat with finger. Call a physician or Poison Control Center immediately. Do not induce vomiting or give anything by mouth to an unconscious person.
If in eyes: Flush with water for at least 15 minutes and get medical attention.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. Do not use 2-PAM, opiates, or cholinesterase-inhibiting drugs.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

KEEP OUT OF REACH OF CHILDREN AND ANIMALS

Caution: May be harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly after handling.
Wear regular long-sleeved work clothing. Change to clean clothing daily.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water or wetlands.

except for rice uses. Discharge from rice fields may kill aquatic and estuarine invertebrates. Do not contaminate water by cleaning of equipment or disposal of wastes. Do not apply when weather conditions favor drift from areas treated. Do not apply where runoff is likely to occur.

Carbaryl is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

RE-ENTRY STATEMENT

Do not enter treated areas without protective clothing until dusts have settled. Because certain states may require more restrictive reentry intervals for various crops treated with this product, consult your State Department of Agriculture for further information. Do not apply this product in such a manner as to directly or through drift expose workers or other persons. The area being treated must be vacated by unprotected persons. Written or oral warnings must be given to workers who are expected to be in a treated area or in an area about to be treated with this product. Oral warnings must be given if there is reason to believe that written warnings cannot be understood by workers. When oral warnings are given, warnings shall be given in a language customarily understood by workers. Written or oral warnings must include the following information:

CAUTION

(Insert area or field description) treated with carbaryl on (insert date of application). In case of accidental exposure: Call a doctor (physician), clinic or hospital immediately. Explain that the victim has been exposed to carbaryl and describe his condition. For further information see the "STATEMENTS OF PRACTICAL TREATMENTS" portion of the pesticide label. Do not enter treated areas without protective clothing until dusts have settled.

STORAGE AND DISPOSAL (12 pt.)

PROHIBITIONS: Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Do not reuse empty container. Do not store under conditions which might adversely affect the container or its ability to function properly. Such conditions include, but are not limited to, positioning of the container in storage, storage temperature, potential for crushing or damage due to stacking, and penetration of moisture.

CLEAN CROP SEVIN 5 BAIT
EPA Reg. No. 34704-23
Page three

STORAGE: Store in safe manner. Store in original container only. Store in cool, dry place. Keep container tightly closed when not in use. Reduce stacking height where local conditions, such as humidity or pallet overhang, can affect package strength. Personnel should use clothing and equipment consistent with good pesticide handling.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Plastic: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Paper Bag: Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

HOUSEHOLD USE DISPOSAL

Do not reuse container (bottle, can, jars). Rinse thoroughly before discarding in trash. Securely wrap original container in several layers of newspaper and discard in trash.

GENERAL INFORMATION

READ ENTIRE LABEL. USE IN STRICT ACCORDANCE WITH LABEL DIRECTIONS AND CAUTIONS.

IMPORTANT PRE-HARVEST AND GRAZING USE INFORMATION

Tolerances established under the Federal Food, Drug and Cosmetic Act permit the sale of crops bearing probable SEVIN residues when SEVIN is used in accordance with label directions. If SEVIN insecticide is used in accordance with label directions, these crops including bean vines, carrot tops, pea vines, sugar beet tops, corn forage and cotton forage may be grazed or harvested for use as feed for dairy and meat animals without resulting in residues in milk or meat in excess of established tolerances.

APPLICATION INSTRUCTIONS

Broadcast application may be made with either ground equipment or aircraft, under conventional, minimal and no-till practices. Treatments may be repeated as necessary. Where a range of rates is provided, use lower rates for light infestation, early instars or when foliage is not too dense; use higher rates for moderate to heavy infestation, advanced instars or when foliage is thick. For best results on lawn and turf, cut lawn, remove clippings and water thoroughly before treatment. Dosages from 20 to 48 pounds

CLEAN CROP SEVIN 5 BAIT

EPA Reg. No. 34704-23

Page four

per acre of bait will control cutworms, armyworms, field crickets, house crickets, mole crickets, darkling beetles, grasshoppers, sowbugs and ants.

Subsequent plantings of food and feed crops in treated soil must be made with crops on this or other carbaryl label(s).

CROP	RATE/ ACRE	RATE/ 1000 SQ. FT.	DAYS BETWEEN FINAL APPLICATION AND HARVEST
Cucumbers, melons, peanuts, pumpkins, squash	20 Lbs	7 1/2 Ozs	0
Birdsfoot trefoil (forage and hay), clover, wheat (pasture, forage)	30 Lbs	11 Ozs	0
Lawn and turf — GROUND APPLICATION ONLY	30 Lbs	11 Ozs	7
Beans, carrots, corn (field, sweet, pop), eggplant, fence row and non-crop areas, forage grasses, okra, pasture (rangeland and wasteland) peppers, potatoes, sorghum forage, soybeans, sweet potatoes, tobacco beds, tomatoes	20-40 Lbs	7 1/2-14 1/2 Ozs	0
Asparagus, strawberries	40 Lbs	14 1/2 Ozs	1
Broccoli, Brussels sprouts, cabbage, cauliflower, garden beets (roots), head lettuce, horseradish, kohlrabi, parsnips, radishes, rutabagas, salsify (roots), tobacco (field), turnips (roots)	20-40 Lbs	7 1/2-14 1/2 Ozs	3
Alfalfa, Peas	30 Lbs	11 Ozs	3
Cotton	30 Lbs	11 Ozs	7
Celery, Chinese cabbage, collards, dandelion, endive (escarole), garden beet tops, Hanover salad, kale, leaf lettuce, mustard greens, parsley, rice*, salsify (tops), spinach, sugar beets, Swiss chard, turnip tops * Do not use on rice fields in which crayfish and/or catfish farming is included in the cultural practice	20-40 Lbs	7 1/2-14 1/2 Ozs	14
Wheat (grain — Fall or Spring)	30 Lbs	11 Ozs	21
Sorghums (milo, grain sorghum, sweet sorghum and hybrids)	20-40 Lbs	7 1/2-14 1/2 Ozs	21 (grain)
Oats, barley and rye (fall treatments)	20-40 Lbs	7 1/2-14 1/2 Ozs	Do not feed treated fodder to dairy animals or animals being finished for slaughter or allow them to graze on crops treated with this product
Sunflowers	20-40 Lbs	7 1/2-14 1/2 Ozs	60 (crop for oil, food or feed purposes) 0 (crop for seeding purpose only)
(all insects listed above except darkling beetles) for the following crops.			
annual plants — herbaceous plants and shrubs, trees, woody plants (ornamental), grapes	40-48 Lbs	14 1/2-17 1/2 Ozs	0
Apples, cherries, peaches, pears, plums	40-48 Lbs	14 1/2-17 1/2 Ozs	1
Nectarines	40-48 Lbs	14 1/2-17 1/2 Ozs	3
Citrus	40-48 Lbs	14 1/2-17 1/2 Ozs	5
Apricots	40-48 Lbs	14 1/2-17 1/2 Ozs	28

Note: Do not make more than two applications after grain heads of wheat, alfalfa and clovers emerge from boot. Do not allow animals to graze on treated sunflower crops

NOTICE

Platte Chemical Co., Inc. warrants that this material conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the directions for use. This product is sold subject to the understanding that the buyer assumes all risks of use or handling which may result in loss or damage which are beyond the control of the seller, such as for example incompatibility with other products, the manner of its use or application, or the presence of other products or materials in or on the soil or crop. Platte Chemical Co., Inc. makes no other express or implied warranty of fitness or merchantability. The exclusive remedy of the buyer or user, and the limit of the liability of Platte Chemical Co., Inc., or any other seller, for any and all losses, injuries or damages resulting from the use or handling of this product shall be the purchase price paid by the user or buyer for the quantity of this product involved. The buyer and all users are deemed to have accepted the terms of this notice, which may not be varied by any verbal or written agreement.

CLEAN CROP SEVIN 5 BAIT
EPA Reg. No. 34704-23
Page five

Formulated for

PLATTE CHEMICAL COMPANY
150 SOUTH MAIN ST.
FREEMONT, NEBRASKA 68025

4 9
5 2
6 3
7 4
8 5