

SEVIN* 10 DUST

INSECTICIDE

ACCEPTED
34704-25

DIRECTIONS FOR USE

Apply dust when insects first appear and repeat at 7 to 10 day intervals as required to maintain control. Apply thoroughly to upper and lower leaf surfaces. Sevin 10 Dust may be applied up to the day of harvest except where noted.

ASPARAGUS: Asparagus Beetles - Use 10 to 20 pounds dust per acre for use on seedlings or spears. Do not apply more than once every 3 days. Allow 1 day between application and harvesting asparagus spears. Use 20 to 40 pounds dust per acre on ferns at post-harvest.

BEANS: Mexican Bean Beetle - Use 5 pounds dust per acre. Bean Leaf Beetle, Leafhoppers, Velvet Bean Caterpillar, Western Bean Cut Worm - Use 10 pounds of dust per acre. Armyworms, Corn Earworm, Plant Bugs - Use 10 to 15 pounds of dust per acre.

CORN: Corn Earworm, European Corn Borer, Fall Armyworm, Sap Beetles, Flea Beetles - Use 10 to 20 pounds dust per acre. For larvae in whorl (budworm damage) and foliage feeders, apply to entire plant. For insects attacking silks and ears, apply 2 to 3 day intervals starting when first silks begin to dry. Three or more applications may be required. Timing and good coverage are absolutely essential for effective control.

CUCUMBERS, MELONS, PUMPKINS, SQUASH: Pickleworm, Melonworm - Use 5 to 10 pounds of dust per acre. Cucumber Beetles, Squash Bugs - Use 10 pounds of dust per acre. Avoid excessive applications. Do not treat when tender foliage is wet or in presence of prolonged high humidity.

EGGPLANT, PEPPER, POTATO, TOMATO: Colorado Potato Beetle, Flea Beetles, Leafhoppers - Use 5 to 10 pounds of dust per acre. European Corn Borer, Tomato Fruitworm, Tomato Hornworm, Stink Bugs - Use 10 to 20 pounds of dust per acre.

ALFALFA, CLOVERS, COWPEAS, SOYBEANS, PEANUTS: Blister Beetle, Mexican Bean Beetle - Use 5 to 10 pounds of dust per acre. Alfalfa Caterpillar, Green Cloverworm, Leafhoppers, Thrips, Velvet Bean Caterpillar - Use 10 pounds of dust per acre. Armyworms, Stink Bugs, Webworms - Use 10 to 15 pounds of dust per acre. Some leaf injury may occur if applications are made when tender foliage is wet.

FORAGE, GRASSES AND PASTURE: Armyworms, Stink Bugs, Thrips - Use 10 to 15 pounds of dust per acre. For optimum thrips control in grasses grown for seed, direct dust into boot.

ACTIVE INGREDIENTS:

*Carbaryl (1-Naphthyl N-methylcarbamate) - - - -	10.0%
INERT INGREDIENTS: - - - - -	90.0%
Total - - - - -	100.0%

*Sevin

*Trademark of Union Carbide Corporation
for the active ingredient carbaryl.

CAUTION: Keep Out of Reach of Children.

HARMFUL IF SWALLOWED OR INHALED. Avoid breathing of dust. Do not take internally. Avoid contact with skin and eyes. Wear regular long-sleeved work clothing. Change to clean clothing daily. Wash hands and face before eating. Wash thoroughly after handling.

AVOID CONTAMINATION OF FOOD, FEED, WATER SUPPLIES, STREAMS AND PONDS DURING APPLICATION OR WHEN CLEANING EQUIPMENT.

NOTE FOR PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. Do not use 2-PAM, opiates, or cholinesterase-inhibiting drugs.

BEE CAUTION

May Kill Honeybees in Substantial Numbers.

This product is highly toxic to BEES exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

Do Not Use when value of Bees as pollinators is more important than insect control. Before applying, warn beekeepers to locate hives beyond bee flight range until one week after application or to take other equally effective precautions.

DISPOSE OF WASTES by burying in non-crop lands away from water supplies. Container should be disposed of by burying with wastes or by burning. Keep out of smoke.

EPA Reg. No. 960-193

EPA Est. No. 00960-CO-02

NET WEIGHT 50 LBS.

DIRECTIONS FOR USE

SORGHUMS: Armyworms, Corn Earworm, Stink Bug, Webworms - Use 10 to 20 pounds of dust per acre. Sorghum Midge - Use 15 pounds of dust per acre. Direct dust into the forming heads. To control sorghum midge, apply 3 to 4 days after heads have emerged from boot. Do not apply within 21 days of harvest of grain. No time limitation on forage.

SUGAR BEETS: Armyworms, Leafhoppers, Webworms - Use 10 to 20 pounds of dust per acre. Do not apply within 14 days of harvest.

STRAWBERRIES: Meadow Spittlebug, Strawberry Leaf Roller, Strawberry Weevil - Use 10 to 20 pounds of dust per acre. Do not apply within 1 day of harvest.

LETTUCE, CARROTS: Flea Beetles, Harlequin Bug, Leafhoppers - Use 5 to 10 pounds of dust per acre. Six-spotted Leafhopper - Use 10 to 15 pounds of dust per acre. Use in a 5 to 7 day schedule. Armyworms, Imported Cabbage Worm, Corn Earworm, Spittlebugs, Tarnish Plant Bugs, Stink Bugs - Use 10 to 20 pounds of dust per acre. For optimum control of corn earworm after lettuce heads begin to form, use 20 pounds in a 5 to 7 day schedule. Do not use for cabbage looper control. Do not apply within 3 days of harvest of head lettuce or within 14 days of harvest of leaf lettuce.

CHICKENS, PARTRIDGES, PHEASANTS, PIGEONS, DUCKS, GEESE, TURKEYS, GAME BIRDS: Chicken Mite, Fleas, Lice, Northern Fowl Mite. **ON BIRDS:** Use 1/2 pound dust per 100 birds. Repeat in 4 weeks if necessary. Apply thoroughly by means of a shaker can or hand duster. Do not apply within 7 days of slaughter. Do not contaminate feed or drinking water. **Bedbugs, Chicken Mite, Fleas - IN OR AROUND POULTRY HOUSES:** Apply 1/2 pound per 40 square feet of litter or floor space, and to roosts and adjacent areas by means of a shaker can, hand duster or power duster. Force dust into cracks and crevices. Repeat as necessary. Do not treat nestlitter. Avoid direct contamination of eggs. Do not house birds in treated houses within 7 days of slaughter.

AVOID CONTAMINATION OF FOOD, FEED, FEEDING TROUGHS AND WATERING RECEPTACLES.

NOTICE: Buyer assumes all risk of use, storage, or handling of this product when not in strict accordance with directions given herewith.



BALCOM CHEMICALS, INC.



240 22nd STREET

GREELEY, COLORADO 80631

P. O. BOX 667