

## STATEMENT OF PRACTICAL TREATMENT

If swallowed—Call a physician or Poison Control Center immediately. Drink one or two glasses of water and induce vomiting by touching the back of throat with finger. Repeat until vomit fluid is clear. Do not induce vomiting or give anything by mouth to an unconscious person.

If inhaled—Remove victim to fresh air. Apply artificial respiration if indicated.

If on skin—Remove contaminated clothing and wash affected area with soap and water.

In Case of Eye Contact: Immediately flush eyes with plenty of water for at least 15 minutes. Call a physician.

## ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not contaminate water by cleaning of equipment or disposal of wastes. In the event that treated seeds are spilled and cannot be salvaged, completely cover them with soil. **NOTE TO USER:** Federal Government and most states have seed laws and regulations requiring an appropriate caution statement be affixed to bags containing treated seed such as: "Treated Seed - Do Not use for food or feed."

## ENDANGERED SPECIES RESTRICTIONS

The use of this product in a manner which may harm endangered species or adversely modify their habitat is prohibited by Federal law. The following endangered species and the general location of their habitats have been identified by county:

|                           |  |
|---------------------------|--|
| Blackwater Darter         | Alabama (Lauderdale, Limestone and Madison)          |
|                           | Tennessee (Lawrence and Wayne)                       |
| Pecos gambusia            | New Mexico (Chaves and Eddy)                         |
|                           | Texas (Reeves, Jeff Davis and Pecos)                 |
| Freshwater Mussel         | Tennessee (Hancock, Claiborne, Hawkins and Sullivan) |
|                           | Virginia (Smyth, Scott, Lee, Washington and Grayson) |
| Commanche Springs Pupfish | Texas (Reeves and Jeff Davis)                        |

Prior to use of this product in the above counties you must determine that these species are not located in, or immediately adjacent to, the areas to be treated with captan. Consult your regional U.S. Fish and Wildlife Service office (Endangered Species Specialist) or State Wildlife Agency if there is any doubt as to the areas where the outdoor use of captan is prohibited.

## DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

## STORAGE AND DISPOSAL

**PROHIBITIONS:** Do not contaminate water, food, or feed by storage or disposal. Do not store under conditions which might adversely affect the container or its ability to function properly. Such conditions include, but are not limited to, positioning of the container in storage, storage temperature, potential for crushing or damage due to stacking, and penetration of moisture.

**STORAGE:** Store in safe manner. Store in original container only. Store in cool, dry place. Reduce stacking height where local conditions, such as humidity or pallet overhang, can effect package strength. Personnel should use driving and equipment

by incineration, or allowed by State and local authorities, by burning. If burned, stay out of smoke.

Mixers, loaders, and applicators, when mixing, loading, and applying, must wear mid-forearm to elbow-length natural or synthetic rubber, vinyl, or plastic gloves impermeable to captan; boots or overshoes; one-piece overalls which have long sleeves and long pants; face shield or goggles and a hat or other appropriate head covering. A dust mask must be worn when mixing and loading this product.

Clothing worn while loading, mixing, and applying this product must be laundered separately from other clothing before reuse. Clothing that may have been drenched or heavily contaminated must be disposed of in accordance with State and local regulations.

**POTATO SEED PIECE TREATMENT**—To control Brown Eye, Verticillium Wilt, Seed Decay (Fusarium), Damping-Off: Apply sufficient amount of CAPTAN 5 DUST to provide thorough coverage. Use 1 to 2 pounds of this product per bushel of cut seed. Whole or 1-drop seed will need less material. Depending on the number of cuts made in the seed piece, the amount of dust used will vary as a result of the number of surface areas exposed for treatment. CAPTAN 5 DUST will NOT protect against fungal inoculum already within the tuber or bacterial diseases. Dust seed pieces before planting, but pieces should be treated within 6 hours of cutting. If planting is delayed more than 1 or 2 days, treated pieces should be stored 2 days in open crates before bagging.

**NOTE TO USER:** Tuber damage and/or decay may occur on sandy soils under moist soil conditions and higher than normal soil temperatures. Do not use if these conditions can occur or if there has been any past history of seed piece decay due to this problem. This product will not control bacterial soft rot. Not for use in Jefferson County, Idaho.

## DO NOT USE TREATED SEED FOR FEED OR FOOD

## NOTICE

**DISCLAIMER**—The label instructions for the use of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and should be followed carefully. However, it is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application all of which are beyond the control of Platte Chemical Company. All such risks shall be assumed by the user. Platte Chemical Company warrants only that the material contained herein conforms to the chemical description on the label and is reasonably fit for the use herein described when used in accordance with the directions for use, subject to the risk referred to above. Any damages arising from a breach of this warranty shall be limited to direct damages and shall not include consequential commercial damages such as loss of profits or values or any other special or indirect damages. Platte Chemical Company makes no other express or implied warranty, including any other express or implied warranty of FITNESS or of MERCHANTABILITY.

FORMULATED FOR  
PLATTE CHEMICAL COMPANY, INC.

# CAPTAN

SEP 18 1986

Under the Federal Insecticide, Fungicide, and Rodenticide Act, CAPTAN 5 DUST is registered under EPA Reg. No. 34704-30

# 5%

## Potato Seed Treater

### ACTIVE INGREDIENT

### BY WEIGHT

|  |        |
|--|--------|
| Captan (N-trichloromethylthio-4-cyclohexene-1,2-dicarboximide) | 4.7%   |
| Related (Derivatives)  | 0.3%   |
| INERT INGREDIENTS:   | 95.0%  |
| TOTAL  | 100.0% |

A FUNGICIDE FOR THE PROTECTION OF  
POTATO SEED PIECES AGAINST  
CERTAIN DISEASES.

KEEP OUT OF REACH OF CHILDREN  
**DANGER**

SEE SIDE PANEL FOR ADDITIONAL  
PRECAUTIONARY STATEMENTS

EPA REG. NO. 34704-30

EPA EST. NO.

NET WEIGHT \_\_\_\_\_ POUNDS