

8-20-75  
Handwritten  
34704-34

# CLEAN CROP®

## CAUTION:

KEEP OUT OF REACH OF CHILDREN.

Harmful if swallowed.

Do not get in eyes, on skin, or on clothing.

Wash hands with soap and water after handling and before eating or smoking.

Wear clean clothing daily.

Do not contaminate water, feed or food stuffs.

Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

Do not allow spray to drift to other crops, trees or plants that might be defoliated.

Destroy used containers. Do not re-use for any purpose. Rinse and drain empty containers into spray tank at least three successive times, then puncture or crush, and bury in an isolated area away from water supplies. Wash and clean spray equipment after use. Do not allow wash water to contact desirable plants.

NOTICE: Buyer assumes all risks of use, handling and storage of this product when not in accordance with directions given herewith.

# CHLORATE - 3 HARVEST AID

## CAUTION:

KEEP OUT OF REACH OF CHILDREN

See Other Precautions on Side Panels

ACTIVE INGREDIENT:	BY WEIGHT:
Sodium Chlorate .....	28%
INERT INGREDIENTS:.....	72%
	100%

CONTAINS 3 POUNDS OF SODIUM CHLORATE  
PER GALLON IN A FIRE-RETARDANT SOLUTION

NET CONTENTS \_\_\_\_\_ GALLONS

34704-34  
EPA Reg. No. ~~960-284~~  
EPA Est. No. 960-CO-01

READ ENTIRE LABEL. FOLLOW DIRECTIONS AND PRECAUTIONS.  
APPLICABLE STATE AND FEDERAL REGULATIONS.  
STORE IN COOL, DRY PLACE.

Weather and plant conditions are high temperature, high humidity, and defoliation are good plant maturity. Active vegetative growth, little nitrogen, and a healthy condition are favorable for good defoliation. Apply 7 to 10 days before anticipated harvest.

Thorough coverage is essential.

COTTON: Apply 1 to 1 1/3 gallons per acre. Use the lower rate for good defoliation. Apply more than 7 days before harvest. Do not apply more than 7 days before harvest.

GRAIN SORGHUM DESICCATION: Apply 1 to 1 1/3 gallons per acre. Use the lower rate when temperatures require more time for desiccation.

Use 1 1/2 to 2 gallons of this product 7 to 10 days before anticipated harvest.

For airplane application apply 1 to 1 1/3 gallons per acre. Use the lower rate when temperatures require more time for desiccation. Swath width should be regulated to give best spray coverage and avoid drift to susceptible nearby crops.

# CHLORATE - 3 HARVEST AID

## CAUTION:

KEEP OUT OF REACH OF CHILDREN

See Other Precautions on Side Panels

ACTIVE INGREDIENT:	BY WEIGHT:
Sodium Chlorate .....	28%
INERT INGREDIENTS: .....	72%
	100%

CONTAINS 3 POUNDS OF SODIUM CHLORATE  
PER GALLON IN A FIRE-RETARDANT SOLUTION

NET CONTENTS \_\_\_\_\_ GALLONS

34704-34  
EPA Reg. No. ~~960-204~~  
EPA Est. No. 960-CO-01

## GENERAL INFORMATION

READ ENTIRE LABEL. USE ONLY IN ACCORDANCE WITH DIRECTIONS AND APPLICABLE STATE AND FEDERAL REGULATIONS.

STORE IN COOL, DRY PLACE.

Weather and plant conditions affect defoliation. Weather conditions favoring good defoliation are high temperature, high humidity and good soil moisture. Plant conditions favoring good defoliation are good plant maturity, (e.g. for cotton, boll load heavy with all bolls mature, no active vegetative growth, little or no secondary growth), good fertility without excessive nitrogen, and a healthy condition of the plants. For best results, leaves should be green and turgid at the time of application, not damaged by insects, wilted or toughened by poor fertilization or other cultural practices.

Thorough coverage is essential.

## DIRECTIONS FOR USE

**COTTON:** Apply 1 to 1 1/3 gallons of CHLORATE-3 HARVEST AID in 4 to 6 gallons of water per acre. Use the lower rate of 1 gallon in 4 to 6 gallons of water when conditions are favorable for good defoliation. Use the higher rate of 1 1/3 gallons when all conditions may not favor good defoliation. Apply 2 to 3 weeks prior to anticipated picking date, but no later than 7 days before harvest. Do not graze treated areas for feeding waste to livestock.

**GRAIN SORGHUM DESICCATION:** To reduce the moisture content in grain sorghum prior to harvest, apply by aircraft while crop is standing. Thorough coverage is essential. Low temperatures require more time for this product to act, and the higher rate is required.

Use 1 1/2 to 2 gallons of this product per acre in sufficient water to give good coverage. Apply 7 to 10 days before anticipated harvesting date.

For airplane application apply the required amount in 5 to 10 gallons of total solution per acre. Use the lower rate when grain sorghum and weather conditions are favorable for rapid desiccation. Swath width should not exceed wing span; altitude and speed of aircraft should be regulated to give best spray coverage. Do not apply during windy weather as spray may drift to susceptible nearby crops and cause damage. Do not apply during very high temperatures.



**BALCOM CHEMICALS, INC.**  
240 22ND STREET P.O. BOX 667 GREELEY COLORADO 80631 PHONE 303-352-4750

